

Thermochemical and Rheological Dynamics of Fruit Preserve Production: Integrating Beet Sugar and Extracted Pectin for Optimal Gelation

Introduction to Viscoelastic Fruit Preserves

The formulation and commercial production of high-quality fruit preserves, specifically jams and marmalades, reside at a highly complex intersection of carbohydrate chemistry, polymer rheology, and thermodynamic processing. At its foundational level, the manufacturing of a shelf-stable, organoleptically superior jam relies on the precise calibration of three primary components: a pectin-rich fruit substrate, a high-purity sucrose matrix, and a meticulously controlled acidic aqueous environment. When processed under optimal thermal conditions, these elements undergo a rigorous sol-gel transition. This transition yields a semi-solid viscoelastic matrix capable of immobilizing water, arresting microbial proliferation, and suspending macroscopic fruit particulates evenly throughout the continuous liquid phase. Regulatory standards and microbiological safety paradigms dictate that standard commercial and high-grade artisanal jams must be concentrated to a minimum of 65% soluble solids, universally quantified as 65° Brix¹. At this exact concentration threshold, the water activity of the system drops to approximately 0.80 to 0.82¹. This thermodynamic depression of unbound water creates an impenetrable hurdle against the proliferation of spoilage bacteria and suppresses the metabolic activity of most xerophilic molds and osmophilic yeasts, ensuring ambient shelf stability².

Historically, preserve manufacturing relied on the inherent pectin content of the primary fruit. However, modern culinary and industrial methodologies necessitate the exogenous addition of extracted pectins to guarantee consistent yield stress and textural rigidity, particularly when working with low-pectin botanical varieties. The structural integrity of the final gel is heavily dictated by the molecular weight, the degree of esterification, and the spatial conformation of the pectin polymer, alongside the colligative properties of the sucrose co-solute. This comprehensive report elucidates the physical chemistry and industrial mechanics of jam and marmalade production, with a specific focus on utilizing highly refined beet sugar and naturally extracted High Methoxyl pectin sourced from under-ripe apples and citrus albedo. Furthermore, it details the localized logistical framework for acquiring these raw materials within the Berlin, Germany, commercial ecosystem, providing an end-to-end analysis from agricultural sourcing to the rheological validation of the final macromolecular gel.

Agronomic Sourcing and Supply Chain Architecture in Berlin

The architectural foundation of a premium fruit preserve requires uncompromising quality and

traceability in its foundational ingredients. For a production facility or high-end artisanal operation situated in Berlin, the regional supply chain offers a robust framework for sourcing organic, ecologically sustainable, and highly refined raw materials that align with stringent European Union organic regulations.

The global sugar market is predominantly supplied by tropical sugarcane, but the European agricultural sector is overwhelmingly dominated by the cultivation of the sugar beet. For production in Berlin, premium organic beet sugar can be procured directly through regional agricultural cooperatives and specialized organic wholesalers. Südzucker AG, a primary European producer, operates processing facilities in southern Germany, such as the plant in Rain am Lech, where over 80% of the processed organic sugar beets are cultivated in surrounding Bavarian soils⁵. This facility supplies certified Bioland and Biokreis organic beet sugar⁵. Bioland standards significantly exceed baseline EU organic requirements, mandating strict multiannual crop rotations, the preservation of soil organic carbon, and the absolute prohibition of synthetic agrochemicals and neonicotinoid seed treatments⁵.

For commercial operations in Berlin, this Bioland-certified, EU Category II fine crystalline beet sugar is typically acquired in robust, recyclable 25-kilogram multi-wall paper sacks⁸. Purchasing through regional distributors or direct-to-trade supply networks yields significant economies of scale, with prices ranging from 1.96 EUR per kilogram for single sacks down to 1.39 EUR per kilogram for palletized bulk orders exceeding 36 sacks⁸. Smaller-scale or pilot productions can source Alnatura or Südzucker organic beet sugar directly from extensive local bio-supermarket chains. The LPG Biomarkt GmbH operates multiple large-scale locations across Berlin, including facilities in Neukölln, Steglitz (Albrechtstraße), and Moabit, prioritizing goods produced within a 200-kilometer radius of the capital to minimize carbon emissions¹⁰.

To extract natural pectin, the raw biomass must possess a high concentration of protopectin, the water-insoluble parent pectic substance found in the primary cell walls and middle lamella of plant tissues¹⁴. Apple pomace, the fibrous byproduct of apple juice pressing, is an exceptional source, consisting of 10% to 17% pectin on a dry matter basis¹⁶. If raw pomace from a cider press is unavailable, under-ripe whole apples represent the superior alternative. In under-ripe fruit, the protopectin has not yet been enzymatically depolymerized into soluble pectin by endogenous polygalacturonases during the natural ripening process. The Berlin region benefits immensely from its proximity to the Werder (Havel) agricultural district in Brandenburg, a historically significant fruit-growing region. Bulk quantities of under-ripe, high-acid apple cultivars can be sourced directly from Werder orchards or via the extensive network of Berlin's weekly farmers' markets, such as the bi-weekly market at Maybachufer in Neukölln¹⁹.

Citrus peels, specifically the inner white spongy tissue known as the albedo, are the most globally abundant natural source of pectin, yielding 20% to 30% on a dry matter basis¹⁶. While citrus is not cultivated in the temperate German climate, wholesale markets in Berlin, such as the Berliner Großmarkt, provide continuous access to untreated, unwaxed organic lemons and bitter oranges imported from Spain or Italy²⁰. Utilizing certified organic citrus is mandatory, as the peel of conventionally grown citrus heavily accumulates fungicidal residues, such as imazalil or thiabendazole, which would be aggressively concentrated during the acid extraction of the pectin.

Physicochemical Distinctions: Beet Sugar Versus Cane Sugar

While beet sugar and cane sugar are chemically identical in their highly refined states, both yielding greater than 99.9% pure sucrose and conforming to the ICUMSA 45 standard for brilliant white, highly crystalline product, their integration into boiling fruit matrices introduces distinct thermochemical phenomena⁷. These operational differences are primarily driven by trace impurities inherent to the disparate biological sources.

Cane sugar often retains microscopic traces of cane molasses, which imparts a subtle, complex flavor profile that is frequently described as rich or caramelized, even in its highly refined white state²². Conversely, beet sugar is completely devoid of molasses, rendering its flavor profile exceptionally neutral and providing a clean, unadulterated sweetness that allows delicate fruit aromatics to dominate the sensory experience of the preserve²². However, experienced confectioners and preserve manufacturers frequently report that beet sugar exhibits a higher susceptibility to rapid caramelization and burning when subjected to aggressive, high-temperature boiling²⁵. This necessitates precise thermal regulation during the final stages of the jam concentration process.

The most critical operational challenge introduced by beet sugar is the aggressive generation of surface foam. This foam is a highly stable matrix of trapped air bubbles, described rheologically as a metastable thermodynamic system²⁸. The primary chemical agents responsible for this foaming are saponins, specifically bidesmosidic triterpenoid glycosides endogenous to the root tissue of the sugar beet³⁰. Saponins possess a highly amphiphilic structure, comprising a hydrophobic aglycone base covalently bonded to a hydrophilic oligosaccharide moiety²⁹. Even highly refined beet sugar retains trace parts-per-million concentrations of these surface-active molecules³¹.

When beet sugar is dissolved and subjected to vigorous boiling, the saponins migrate rapidly to the air-water interface. They significantly depress the surface tension of the aqueous phase, frequently reducing it to approximately 40 mN/m³¹. This powerful surfactant activity allows steam and mechanically entrapped air to form a persistent, resilient foam layer on the surface of the boiling masecuite³¹. The stability of this foam is further exacerbated by the presence of pectin and divalent cations, which act synergistically to reinforce the lamellae of the bubbles³³.

Interfacial Thermodynamics and the Imperative of Skimming

In commercial and artisanal preserve manufacturing, the thorough removal, or skimming, of this saponin-stabilized foam is a non-negotiable processing step. Beyond the visual defect of introducing cloudy, opaque, sea-foam-like streaks into an otherwise translucent, jewel-like gel, foam introduces severe structural and microbiological risks²⁸.

Foam represents a localized matrix of exceptionally high air entrapment. If foamy jam is decanted into glass jars, the entrained air mimics liquid volume, leading to a phenomenon known as false headspace²⁸. Upon cooling and the establishment of a vacuum seal, the

microscopic air bubbles collapse, increasing the absolute headspace volume well beyond the standard 0.25-inch margin²⁸. Excessive oxygen trapped within the jar accelerates the oxidative degradation of ascorbic acid and fruit anthocyanins, inducing premature browning and off-flavors during long-term storage²⁸. Furthermore, the insulative thermodynamic properties of the foam can disrupt the thermal lethality kinetics during hot-water bath processing²⁸. This disruption potentially allows xerophilic fungal spores to survive the pasteurization phase and subsequently colonize the headspace boundary²⁸.

To mitigate foaming without relying on synthetic polydimethylsiloxane antifoaming agents, manufacturers employ physical maceration and prolonged osmotic resting²⁸. By mechanically chopping the fruit and macerating it in the beet sugar for 12 to 24 hours prior to thermal processing, a powerful osmotic gradient draws intracellular moisture out of the fruit matrix²⁸. This gradual plasmolysis physically collapses the intercellular air pockets within the parenchyma tissue of the fruit, drastically reducing the total volume of air available for subsequent saponin entrapment during the aggressive boiling phase²⁸.

Structural Biochemistry of Pectin Domains

To engineer a stable, high-yield stress jam from fruits with natively low pectin content, exogenous pectin must be extracted from the procured apple pomace or citrus albedo and introduced into the primary formulation. The structural complexity of the extracted pectin dictates its ultimate utility in a sugar-acid gel. Pectin is a highly heterogeneous plant cell wall macromolecule comprising multiple distinct domains, predominantly Homogalacturonan, Rhamnogalacturonan-I, and Rhamnogalacturonan-II¹⁴.

The Homogalacturonan domain is a linear, unbranched polymer consisting of alpha-1,4-linked D-galacturonic acid residues¹⁴. This smooth backbone is the primary functional component responsible for junction-zone formation during gelation¹⁴. Conversely, the Rhamnogalacturonan-I domain consists of a repeating disaccharide backbone of alpha-1,2-L-rhamnose and alpha-1,4-D-galacturonic acid¹⁴. Crucially, the rhamnose residues within the RG-I domain are highly substituted at the O-4 position with complex, bulky neutral sugar side chains, primarily composed of alpha-1,5-linked arabinans and beta-1,4-linked galactans¹⁴.

Historically, the RG-I domain was considered an impediment to industrial gelation. The insertion of rhamnose into the backbone produces physical kinks in the otherwise linear galacturonan chain, while the massive, bushy arabinan and galactan side chains generate severe steric hindrance³⁴. This steric repulsion physically obstructs adjacent Homogalacturonan chains from aligning closely enough to establish the hydrogen bonds and hydrophobic interactions required for a three-dimensional network¹⁴. However, recent rheological advancements reveal that while the side chains induce steric hindrance, they simultaneously execute a vital function in water-binding and synergistic structural reinforcement³⁶. When blended with exogenous sucrose, the lengthy galactan branches intertwine, creating a pliable, highly resilient continuous network³⁶. This entanglement allows the resulting gel to exhibit high fracture strain and recoverability, moving the gel away from a purely brittle state toward an elastic, spreadable

morphology ideal for fruit preserves³⁶.

Pectin Domain	Structural Composition	Rheological Function
Homogalacturonan (HG)	Linear α - -(1,4)-D-galacturonic acid backbone.	Primary site for intermolecular hydrogen bonding and hydrophobic junction zone formation. Drives gel rigidity.
Rhamnogalacturonan-I (RG-I)	Alternating α - -(1,2)-L-rhamnose and α - -(1,4)-D-GalA backbone.	Introduces backbone "kinks" disrupting continuous rigid alignment.
RG-I Side Chains	α -(1,5)-arabinans and β - -(1,4)-galactans attached to rhamnose.	Generates steric hindrance, but intertwines to provide extreme elasticity, fracture strain, and massive water-binding capacity.
Rhamnogalacturonan-II (RG-II)	Highly branched complex featuring apiose, fucose, and aceric acid.	Facilitates borate-ester cross-linking in the primary cell wall; minor role in food gelation.

Table 1: Biochemical domains of the pectin macromolecule and their respective rheological implications.

[cite: 14, 34, 36, 37]

Methodologies of Natural Pectin Extraction

The objective of extraction is the hydrolytic cleavage of the insoluble protopectin macromolecule, freeing the water-soluble pectic hydrocolloids while meticulously preserving their polymeric backbone length and degree of methyl esterification. The process typically begins with a physical pretreatment. For citrus albedo, subjecting the peels to steam or hot water blanching effectively denatures endogenous pectin methylesterase and polygalacturonase enzymes, preventing them from prematurely depolymerizing the protopectin²¹. Furthermore, blanching softens the cellulosic cell wall architecture, facilitating deeper solvent penetration²¹. The raw material is then dried and milled to an optimized particle size; extractions utilizing apple flour with a particle diameter between 125 and 250 micrometers yield significantly higher pectin

recoveries than roughly chopped pomace, owing to the exponential increase in solvent contact surface area⁴¹.

Pectin extraction is universally executed via hot acid hydrolysis. The selection of the acidic solvent, alongside the precise modulation of temperature and reaction time, determines both the quantitative yield and the qualitative rheological power of the resulting extract. Industrial environments frequently employ harsh mineral acids, such as hydrochloric, sulfuric, or nitric acid, due to their aggressive hydrolytic power. These reagents can achieve absolute mass yields approaching 30% to 35% from citrus peel⁴⁴. However, these harsh conditions precipitate severe structural collateral damage to the pectin polymer⁴⁴. Mineral acids unselectively cleave the glycosidic bonds of the homogalacturonan backbone, drastically reducing the intrinsic viscosity and molecular weight of the hydrocolloid⁴⁴. Furthermore, they induce aggressive de-esterification and trigger runaway Maillard browning reactions with residual neutral sugars, producing dark, visually unappealing extracts with poor solubility⁴⁴.

Consequently, the extraction of premium culinary pectin necessitates the use of milder organic aliphatic acids, specifically citric acid or L-(+)-tartaric acid⁴⁴. While organic acids typically deliver lower total yields (ranging from 10% to 15%), their mild hydrolytic action preserves the integrity of the polymer backbone and protects the critical esterified functional groups⁴⁴. Experimental response surface methodology kinetics indicate that a citric acid solution calibrated to a pH of 2.0 to 2.5, operated at 90°C to 97°C for a duration of 60 to 150 minutes, optimizes the extraction envelope for apple pomace⁴¹. Under these controlled parameters, the hydronium ions selectively target the non-covalent calcium bridges and weak ester linkages anchoring the protopectin to the cellulosic matrix, while leaving the primary covalent backbone intact⁴⁸. Advanced extraction methodologies are increasingly utilizing ultrasonic probe intervention. Exposing the acidic citrus albedo slurry to high-intensity ultrasound (e.g., using a Qsonica transducer) induces acoustic cavitation. The implosion of microscopic cavitation bubbles generates extreme localized shear forces that mechanically fracture the plant cell walls⁴⁹. This drastically accelerates mass transfer, allowing extractions to achieve theoretical maximum yields (up to 34.6%) in an unprecedented 3 to 7 minutes of total processing time, bypassing the thermal degradation inherent to hours-long boiling⁴⁹.

Following the extraction phase, the hot acidic slurry is rapidly filtered. The soluble pectin is subsequently precipitated out of the aqueous phase via the introduction of absolute ethanol or isopropanol⁴³. The alcohol acts as an anti-solvent, drastically depressing the dielectric constant of the medium. This strips the hydration spheres from the highly hydrophilic pectin polymers, forcing them to coagulate into a buoyant, recoverable gelatinous floc that is subsequently pressed, washed, and dried⁵¹.

Analytical Quantification: Degree of Esterification via FTIR Spectroscopy

The single most critical metric defining pectin functionality is its Degree of Esterification, defined as the molar percentage of the D-galacturonic acid carboxyl groups that have been naturally esterified with a methyl group¹⁴. Pectins with a Degree of Esterification greater than 50% are

classified as High Methoxyl pectins, whereas those below 50% are Low Methoxyl pectins¹⁴. Traditional high-sugar preserve manufacturing relies exclusively on High Methoxyl pectin, as it dictates the specific sugar-acid mechanism required for product setting. To assure the extracted batch meets the High Methoxyl criteria without resorting to laborious, chemical-intensive, and time-consuming titrimetric or gas chromatography assays, Fourier-Transform Infrared spectroscopy is employed as a highly accurate, non-destructive analytical tool⁵³. The Degree of Esterification is calculated by analyzing the specific absorption intensities within the characteristic carbonyl region, spanning 1800 to 1500 cm⁻¹, of the infrared spectrum⁵⁶. The characteristic absorption band for the esterified carbonyl group, representing the C=O stretching vibration in the methyl esters, manifests at approximately 1740 to 1747 cm⁻¹⁵³. Conversely, the free, unesterified carboxylate ion, representing the COO⁻ asymmetric stretching, exhibits a strong, broad absorption peak at 1600 to 1639 cm⁻¹⁵³. The degree of esterification can be mathematically derived by taking the ratio of the integrated peak area, or the baseline-corrected absorbance intensity, of the esterified groups relative to the total carboxyl population:

$$DE(\%) = \left(\frac{A_{1740}}{A_{1740} + A_{1630}} \right) \times 100$$

[cite: 53, 58, 59]

Functional Group	Chemical Shift / Wavenumber	Spectral Significance
Esterified Carbonyl (C=O)	1740 - 1747 cm ⁻¹	Stretching vibration of methyl ester groups. High intensity indicates HM pectin.
Free Carboxylate (COO ⁻)	1600 - 1639 cm ⁻¹	Asymmetric stretching of free galacturonic acid. High intensity indicates LM pectin.
Pyranose Ring (C-O-C)	~1100 cm ⁻¹	Carbohydrate fingerprint region; frequently used to normalize spectral data.

Table 2: Key Fourier-Transform Infrared spectral assignments for the quantification of pectin esterification.

[cite: 53, 57, 58, 59]

In complex matrices containing high levels of residual protein, spectral deconvolution software must be utilized to separate overlapping amide absorption bands from the primary carboxylate signals⁵³. However, for purified apple pomace and citrus albedo extracted under controlled, mild citric acid conditions, direct peak ratio analysis routinely yields highly accurate Degree of Esterification values ranging from 68% to 75%⁴¹. This securely positions the extracted biopolymer in the rapid-set High Methoxyl pectin category, mechanically ideal for high-sugar fruit preserve applications³⁵.

Thermochemical Kinetics of the Boiling Phase

The introduction of the fruit, beet sugar, and extracted pectin into the boiling kettle initiates a cascade of complex, interconnected chemical reactions. The application of thermal energy must be rigorously controlled to achieve the target soluble solids concentration while actively managing sucrose inversion, the degradation of native anthocyanin pigments, and the preservation of the pectin backbone.

Kinetics of Sucrose Inversion

During the boiling of a highly acidic jam, operating strictly between pH 2.8 and 3.4, the disaccharide sucrose undergoes a continuous hydrolytic cleavage into an equimolar mixture of its constituent monosaccharides: D-glucose and D-fructose⁶¹. This process is universally known as inversion, as the specific optical rotation of the solution, measurable via a polarimeter, shifts from strongly dextrorotatory (+66.5°) to mildly levorotatory (-20.0°)⁶². This optical shift occurs because the powerful levorotatory specific rotation of the liberated fructose (-92.0°) mathematically overpowers the dextrorotatory nature of the liberated glucose (+52.0°)⁶².

The reaction is catalyzed by hydronium ions provided by the endogenous organic acids of the fruit and the exogenous pectin-extraction acids. Because water is present in vast stoichiometric excess, the reaction is functionally independent of water concentration and rigidly follows pseudo-first-order kinetics⁶¹. The rate of inversion is expressed by the differential equation:

$$-\frac{d[\text{Sucrose}]}{dt} = k_{eff} [\text{Sucrose}]$$

[cite: 64]

Where the effective rate constant (k_{eff}) is exponentially dependent on both temperature and hydronium ion concentration. Empirical kinetics calculate the base rate constant at ambient temperatures to be approximately $1.23 \times 10^{-2} \text{ min}^{-1} \text{ M}^{-1}$, which accelerates drastically

at the boiling point⁶¹. Integrated, the concentration of sucrose at any time t follows standard exponential decay⁶³.

Inversion is absolutely vital to the thermodynamic stability of the jam. Pure sucrose has a rigid solubility limit of approximately 67 g/100 mL at room temperature. If a jam is boiled to 65% or 68% soluble solids using exclusively un-inverted sucrose, the product will inevitably cross the boundary of supersaturation and undergo catastrophic macroscopic crystallization during shelf storage, rendering the texture gritty and commercially unacceptable⁶⁶. The generated invert sugar, however, boasts a significantly higher aggregate solubility and acts as a powerful humectant, interfering with the sucrose crystal lattice and ensuring a smooth, glossy, amorphous texture⁶².

However, inversion must be strictly arrested before completion. Fructose is highly hygroscopic and highly reactive in high-temperature environments. Excessive inversion, exceeding 30% to 40% conversion of the total sugars, results in a jam that is overly fluid due to the depression of the mixture's average molecular weight⁶⁷. More critically, high concentrations of free fructose severely darken the product via accelerated Maillard browning and spontaneous fructose caramelization under hot acidic conditions⁶⁷. The manufacturer must therefore carefully balance the boiling duration to achieve adequate, but not total, sucrose inversion.

Anthocyanin Degradation Kinetics

For preserves featuring deep red or purple fruits, such as cherries, blackberries, or specific citrus varieties, the preservation of visual color is a critical quality parameter. The primary pigments, anthocyanins, prominently including cyanidin-3-glucoside and cyanidin-3-rutinoside, are highly unstable and sensitive to thermal degradation⁶⁹.

The thermal destruction of cyanidin-3-glucoside during boiling follows strict first-order reaction kinetics. In high-sugar environments, the massive concentration of sucrose significantly influences the thermodynamics of the aqueous phase, frequently exerting a minor protective effect against pigment degradation by reducing water mobility and reactant collision frequency. However, under standard atmospheric boiling temperatures, reaching 100°C to 105°C depending on boiling point elevation, the activation energy for the degradation of total anthocyanins is approximately 48.0 to 57.4 kJ/mol⁶⁹. The half-life of these vital pigments at aggressive boiling temperatures shrinks to a mere 6.6 to 15.9 minutes⁶⁹. Consequently, extended boiling phases drastically bleach the fruit matrix, transitioning the brilliant purples and reds into dull, polymerized brown melanoidins⁷¹. This reinforces the necessity for rapid, high-efficiency evaporative boiling.

Thermal Depolymerization of the Pectin Backbone

Simultaneous to sucrose inversion and pigment degradation, the dissolved High Methoxyl pectin polymer is subjected to severe hydrolytic stress. Pectin is inherently unstable at elevated temperatures, undergoing massive, irreversible depolymerization that directly destroys its capacity to form a gel matrix upon cooling⁷². The non-enzymatic thermal degradation of pectin is governed by two competing thermochemical mechanisms: acid hydrolysis and beta-elimination⁷³.

In the highly acidic environment of a boiling jam, typically below pH 3.5, the predominant mechanism of chain scission is acid hydrolysis⁷³. Acid hydrolysis targets the alpha-1,4-glycosidic linkages between the galacturonic acid monomers, directly fracturing the Homogalacturonan backbone⁷⁴. The rate of acid hydrolysis increases exponentially with decreasing pH and has a calculated activation energy of approximately 95 kJ/mol⁷⁴. Interestingly, acid hydrolysis is sterically hindered by high degrees of esterification; therefore, High Methoxyl pectin is marginally more resilient to pure acid hydrolysis than Low Methoxyl pectins⁷⁴.

Conversely, beta-elimination represents the cleavage of the glycosidic bond localized specifically at the C-4 position, concomitant with the removal of a hydrogen atom at the C-5 position of an esterified galacturonic acid unit⁷³. This reaction creates a double bond and shatters the polymer chain⁷⁷. While beta-elimination is universally recognized as the dominant mode of degradation in neutral or alkaline conditions, it still actively and aggressively occurs in heated acidic environments, particularly because High Methoxyl pectin inherently possesses the massive concentration of esterified residues required to drive the reaction⁷⁴. The activation energy for beta-elimination is exceptionally high, calculated at 136 kJ/mol⁷⁴.

Degradation Mechanism	Chemical Action	Activation Energy (Ea)	Primary Drivers
Acid Hydrolysis	Cleavage of α -(1,4)-glycosidic linkages.	~ 95 kJ/mol	Low pH. Rate is hindered by high DE.
β-Elimination	Cleavage at C-4, creating a double bond via proton loss at C-5.	~ 136 kJ/mol	Neutral/alkaline pH, but occurs in acid. Requires esterified groups (high DE).
Demethoxylation	Hydrolysis of methyl esters to free carboxyl groups.	~ 98 kJ/mol	High temperatures. Converts HM pectin to LM pectin over time.

Table 3: Thermodynamic parameters governing the non-enzymatic thermal degradation of pectin during prolonged boiling.

[cite: 73, 74, 76]

Because the activation energy of beta-elimination (136 kJ/mol) is significantly higher than that of acid hydrolysis (95 kJ/mol), the predominant mechanism of pectin destruction is highly

temperature-dependent⁷⁴. At the elevated temperatures required to boil off water and achieve 65° Brix, often reaching 104°C to 105°C, the thermal energy easily surpasses the threshold to aggressively drive both reactions simultaneously⁷⁴.

If the boiling process is protracted, the continuous depolymerization fractures the long High Methoxyl pectin chains into short, rheologically useless oligomers. Once the molecular weight drops below a critical structural threshold, typically below 50 to 60 kDa, the truncated chains lack the spatial geometry, entanglements, and interaction sites required to form continuous junction zones⁷⁸. The inevitable result is a total rheological failure—a runny, liquid syrup that cannot be salvaged, regardless of subsequent cooling⁷⁸.

Rheology of the Sol-Gel Transition

Assuming the boiling phase is executed rapidly, preserving the structural integrity of the High Methoxyl pectin and achieving 65% soluble solids, the ultimate phase of production is the sol-gel transition upon cooling. The gelation of High Methoxyl pectin represents a massive physical, rather than chemical, cross-linking process.

In an aqueous solution at a neutral pH, the unesterified carboxyl groups on the homogalacturonan backbone are fully ionized, carrying strong negative charges⁸⁰. The resulting electrostatic repulsion forces the polymer chains apart, maintaining a highly fluid, solubilized state⁸⁰. To collapse this system into a rigid gel matrix, two exact physiochemical conditions must be met: the suppression of electrostatic repulsion and the severe dehydration of the polymer⁸⁰.

First, the pH of the system must be lowered below the pK_a of pectin, which rests at approximately 3.5⁸⁰. The massive influx of hydronium ions protonates the ionized carboxylate groups, converting them to neutral carboxylic acids⁸⁰. With the repulsive forces neutralized, the long chains can physically approach one another without electrostatic deflection⁸⁰. Second, even when neutralized, pectin chains are heavily sheathed by a tight hydration sphere of highly ordered water molecules bound via hydrogen bonding. The addition of massive quantities of a co-solute, specifically the 65% beet sugar sucrose matrix, is required to break this hydration⁸⁰. Sucrose acts as an aggressively powerful dehydrating agent, outcompeting the pectin for available free water and effectively stripping the hydration spheres from the polymer backbone¹⁴.

Forced into extremely close proximity in a highly thermodynamically unfavorable, low-water environment, the hydrophobic methyl ester groups of adjacent pectin chains seek to minimize their contact with the remaining polar aqueous phase⁸¹. They associate tightly via hydrophobic interactions, while the newly protonated carboxylic acid groups and available hydroxyl groups lock together via dense intermolecular hydrogen bonding⁸¹. These localized areas of intense, zipper-like association are termed junction zones³⁵. When sufficient junction zones form, a continuous three-dimensional polymer network traps the viscous sucrose and water solvent, completing the transition from a liquid sol to a solid gel³⁵.

Viscoelasticity and the Winter-Chambon Criterion

The physical manifestation of this macroscopic gelation can be tracked and quantified via

dynamic oscillatory rheology. The material's mechanical response is characterized by its complex shear modulus, encompassing the elastic storage modulus (G') and the viscous loss modulus (G'')⁸³.

In the hot, pre-gel state, the boiling jam behaves as a viscoelastic liquid. The energy dissipated as heat via viscous flow exceeds the energy stored elastically, meaning G'' is strictly greater than G' ⁸⁶. As the temperature drops and the massive network of hydrogen bonds and hydrophobic junction zones rapidly nucleates, the storage modulus undergoes a sudden, exponential divergence, violently crossing over the loss modulus⁸⁷.

The precise, definitive moment of the liquid-to-solid transition is defined rheologically by the Winter-Chambon criterion⁸⁸. Winter and Chambon established mathematically that at the critical gel point, the storage and loss moduli exhibit identical power-law frequency dependence,

scaling uniformly with the relaxation exponent n ⁸⁶. Mathematically, at the gel point, the phase shift becomes absolutely independent of the oscillation frequency, and the loss tangent ($\tan \delta$) converges to a constant value:

$$\tan \delta = \frac{G''}{G'} = \tan \left(\frac{n\pi}{2} \right)$$

[cite: 86, 88, 91]

For an ideal, stoichiometrically balanced polymer network where the relaxation exponent

$n = 0.5$, the loss tangent equals exactly 1, representing the exact crossover point where

$G' = G''$ ⁸⁸. The critical gel at this specific threshold exhibits a self-similar fractal geometry, possessing infinite steady-shear viscosity yet a zero equilibrium elastic modulus, acting as the precise evolutionary boundary between the fluid state and the semi-solid matrix⁹¹.

The Manifestation of Yield Stress

Upon full thermal equilibration, the mature jam exhibits a defined yield stress. Yield stress is the critical threshold of applied mechanical force below which the material behaves as a rigid, elastic solid, and above which the hydrophobic junction zones fracture, causing the material to shear and flow like a liquid⁹³.

From a consumer and functional perspective, the yield stress governs spreadability and suspension dynamics. A meticulously extracted High Methoxyl pectin, with protected molecular weight and optimized sucrose concentration, will generate a yield stress high enough to suspend heavy fruit particulates uniformly in the jar against the force of gravity, yet low enough to shear smoothly under the pressure of a knife without crumbling or weeping syneresis⁹³.

Utilizing a rheometer, dynamic yield stress is typically quantified via an amplitude sweep,

measuring the exact stress required to initiate the sharp drop in the elastic modulus⁹³.

Process Control: Refractometry and Water Activity

To guarantee that the thermodynamic hurdle of an a_w near 0.80 is met, and to trigger the dehydration mechanisms required for High Methoxyl pectin junction-zone formation, the soluble solids must be rigorously quantified during the boiling process¹. This is universally executed using a refractometer, an optical instrument that measures the refractive index of the syrup, translating the optical bending of light into a standard °Brix value⁹⁶.

Because refractive index is highly sensitive to the density of the medium, and fluid density is inversely proportional to thermal expansion, accurate Brix measurement is strictly temperature-dependent⁹⁶. Refractometers are universally calibrated to a baseline standard of 20°C⁹⁸. Attempting to measure boiling jam directly from the kettle introduces massive mathematical deviations. At elevated temperatures, the expanded, less-dense fluid bends light less severely, resulting in a falsely low Brix reading⁹⁶.

Modern digital refractometers employ Automatic Temperature Compensation, utilizing internal thermistors and algorithms to mathematically correct the reading back to a 20°C baseline⁹⁶. However, Automatic Temperature Compensation systems are generally bounded between 10°C and 30°C¹⁰⁰. A sample pulled directly from a 104°C boiling kettle possesses thermal mass far exceeding the compensatory limits of the instrument's microprocessors¹⁰¹. Consequently, to prevent catastrophic over-boiling—which would irreparably degrade the anthocyanins and irreversibly depolymerize the pectin—the manufacturer must rapidly quench a minute sample of the jam on a room-temperature optical prism. This allows the sample to equilibrate into the compensation system's operational window prior to optical assessment, ensuring a precise 65° Brix measurement¹⁰⁰.

Conclusion

The transformation of raw fruit, water, and sucrose into a shelf-stable, structurally perfect fruit preserve represents a masterclass in controlled polymer chemistry and thermodynamics. Utilizing locally sourced organic beet sugar from the Berlin agricultural sector mandates careful attention to the powerful surfactant properties of inherent triterpenoid saponins. This requires deliberate, physical skimming protocols to prevent foam-induced oxidative degradation, false-headspace generation, and thermal insulation during pasteurization.

Simultaneously, the extraction of High Methoxyl pectin from under-ripe regional apples or citrus albedo demands rigorous control over acidic parameters. By employing mild organic acids like citric acid, manufacturers can successfully liberate the homogalacturonan backbone while preserving the critical degree of esterification and Rhamnogalacturonan-I side branches. This guarantees synergistic water-binding and yields a robust, high-molecular-weight hydrocolloid capable of forming strong junction zones.

The thermal processing phase must function as a rapid, highly optimized intervention. The boil must be brief enough to outpace the exponential kinetics of pectin beta-elimination and anthocyanin degradation, yet long enough to facilitate pseudo-first-order sucrose inversion, preventing post-production crystallization and generating necessary invert humectants. Finally,

the successful formulation culminates in the Winter-Chambon sol-gel transition, where the strategic manipulation of pH and massive sucrose dehydration forces the pectin polymers into an interlocking lattice of hydrophobic junction zones. The mastery of these interconnected physical, chemical, and rheological parameters ensures the reliable production of an optimal, highly structured gel matrix.

Works cited

1. Introduction to Sugar-Based Fruit Products - Agriculture Institute, <https://agriculture.institute/food-processing-and-engineering-i/introduction-to-sugar-based-fruit-products/>
2. Water Activity (Aw) in Beverage Formulation: The Complete Guide, <https://foodscientistforhire.com/water-activity-aw-in-beverage-formulation-the-complete-guide/>
3. Shelf Life Simplified: A Water Activity Based Approach - Neutec Group, <https://www.neutecgroup.com/resource-library/water-activity/white-papers/228-shelf-life-simplified-a-water-activity-based-approach/>
4. Water Activity in Food - Pathogen Modeling Program (PMP) Online, <https://pmp.errc.ars.usda.gov/WaterActivity.aspx>
5. Bio-Rübenzucker für industrielle Anwendungen - suedzucker.com, <https://www.suedzucker.com/de/produkte/bio-ruebenzucker-zucker/>
6. Rüben Zucker in Bio-Qualität I Mein Südzucker, <https://suedzuckermomente.de/produkte/zucker-in-bio-qualitaet/>
7. 31 Growing And Processing Sugar Beets.pdf
8. BIO-Rübenzucker 25 kg von Südzucker - thehoneyfactory, <https://thehoneyfactory.de/bienenfutter/futter-in-selbsterstellung/bio-ruebenzucker-bioland-fein-im-papiersack-25-kg/>
9. 25 kg Bioland Bio-Rübenzucker (Lebensmittelqualität) im Sack ab, <https://bienenzuchtbedarf-geller.de/25-kg-Bioland-Bio-Ruebenzucker-Lebensmittel-qualitaet-im-Sack-ab-Wuerselen-unsere-OEko-Kontrollnr.-DE-OEKO-034/263119>
10. LPG BioMarkt - Kollwitzstraße - LoveNexus, <https://lovenexus.com/de/map/laden/lpg-biomarkt-kollwitzstrae/>
11. LPG BioMarkt / Bio erleben in Berlin und Brandenburg, <https://www.bio-berlin-brandenburg.de/adressen/lpg-biomarkt-gmbh-hauptstrasse-78/>
12. Our stores: Welcome! - LPG Biomarkt, <https://www.lpg-biomarkt.de/en/our-stores-welcome/>
13. LPG Biomarkt in Berlin, Öffnungszeiten & Adressen der Filialen, https://www.in-berlin-brandenburg.com/Einkaufen_Berlin/Supermarkt/Bio/LPG-Biomarkt/
14. Structure-Related Gelling of Pectins and Linking with Other Natural, <https://www.mdpi.com/2073-4360/10/7/762>
15. EXTRACTION OF PECTIN FROM APPLE POMACE, <https://www.iitf.lbtu.lv/conference/proceedings2020/Papers/TF549.pdf>
16. Extraction and Characterization of Pectins From Peels of Criolla,

- <https://www.intechopen.com/chapters/69850>
17. Role of Pectin in Food Processing and Food Packaging - IntechOpen, <https://www.intechopen.com/chapters/65793>
 18. Structure-Related Gelling of Pectins and Linking with Other Natural, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6404037/>
 19. Top 10 Farmers' Markets in Berlin, <https://www.top10berlin.de/en/cat/shopping-261/farmers-markets-2840>
 20. Extraction, characterisation, and enzymatic degradation of lemon, <https://research.wur.nl/en/publications/extraction-characterisation-and-enzymatic-degradation-of-lemon-pe/>
 21. Effects of Blanching on Flavanones and Microstructure of Citrus, <https://www.semanticscholar.org/paper/Effects-of-Blanching-on-Flavanones-and-of-Citrus-Zid-Dhuique-Mayer/82212f8e56bb26ad572468c000fd3b5c2516f132>
 22. How to Choose the Right Sugar Grade – ICUMSA Buyer Guide 2026, <https://theworldmerchants.com/how-to-choose-the-right-sugar-grade/>
 23. ICUMSA Sugar Grades Explained: 45, 100-150, VHP & Raw Cane, <https://mcispcoltd.com/blog/icumsa-sugar-grades-buyers-guide>
 24. Cane vs Beet, the battle of the sugars - Portland Syrups, <https://beveragemixers.com/blogs/psw-blog-posts/cane-vs-beet-the-battle-of-the-sugars>
 25. Cane vs. Beet Sugar: A Difference? - WhatSugar, <https://www.whatsugar.com/post/difference-between-cane-and-beet-sugar>
 26. Beet Sugar vs Cane Sugar | What's the Difference - Indiana Sugars, <https://www.sugars.com/beet-sugar-vs-cane-sugar>
 27. Sugar for Marmalade | Ask Nigella.com, <https://www.nigella.com/ask/sugar-for-marmalade>
 28. Skimming Foam From Jam: Everything You Need To Know, <https://nwedible.com/skimming-jam-foam-questions/>
 29. Foam Properties and Detergent Abilities of the Saponins from, <https://www.mdpi.com/1422-0067/11/11/4417>
 30. Molecularization of Foam-Active Saponins from Sugar Beet Side, <https://pubmed.ncbi.nlm.nih.gov/32881499/>
 31. Sugar Beet Extract (Beta vulgaris L.) as New Natural Emulsifier, https://www.researchgate.net/publication/316576983_Sugar_Beet_Extract_Beta_vulgaris_L_as_New_Natural_Emulsifier_Emulsion_Formation
 32. US2727009A - Antifoam compositions and method of foam inhibition, <https://patents.google.com/patent/US2727009A/en>
 33. Foam formation in biogas plants caused by anaerobic digestion of, https://fing.htwk-leipzig.de/fileadmin/portal/f_i/Fakult%C3%A4t/Professoren/Schenk/Publikationen/Schaumbildung.pdf
 34. Insights into pectin O-acetylation in the plant cell wall - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9922974/>
 35. Pectin Basics - Gelling properties and applications - CyberColloids, <https://cybercolloids.net/information/technical-articles/pectin-basics-gelling-properties-and-applications/>

36. Synergistic gelling mechanism of RG-I rich citrus pectic,
https://eprints.whiterose.ac.uk/id/eprint/171496/3/Zheng_FOODCHEM_2021.pdf
37. Pectins, <https://www.chem.gla.ac.uk/staff/mikej/Pectin.htm>
38. The impact of rhamnogalacturonan-I side chain monosaccharides,
https://www.researchgate.net/publication/271684931_The_impact_of_rhamnogalacturonan-I_side_chain_monosaccharides_on_the_rheological_properties_of_citrus_pectin
39. Structural, rheological and emulsifying properties of RG-I enriched,
https://www.researchgate.net/publication/366676420_Structural_rheological_and_emulsifying_properties_of_RG-I_enriched_pectins_from_sweet_and_sour_cherry_pomaces
40. Effect of steam blanching on peelability and quality of Citrus ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8357887/>
41. Extraction of pectin from apple pomace - SciSpace,
<https://scispace.com/pdf/extraction-of-pectin-from-apple-pomace-1nm5qeb1sv.pdf>
42. Extraction of Pectin From Apple Pomace - SciELO,
<https://www.scielo.br/j/babt/a/xS3NhspJ4CDv3Vbswc5zRRC/?format=pdf&lang=en>
43. The Influence of Extraction Conditions on the Yield and Physico,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9002691/>
44. Pectin Extraction from Citrus Waste: Structural Quality and Yield with,
<https://www.mdpi.com/2673-7167/5/3/32>
45. pH-DEPENDENT YIELD AND PHYSICO-CHEMICAL PROPERTIES,
<https://ijtech.eng.ui.ac.id/download/article/3595>
46. Deciphering Pectin: A Comprehensive Overview of Its Origins ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11740133/>
47. Optimization of Pectin Enzymatic Extraction from Malus domestica,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6600438/>
48. WO2016202986A1 - Process for extraction of pectin - Google Patents,
<https://patents.google.com/patent/WO2016202986A1/en>
49. Optimization of Pectin Extraction Process from Albedo of Citrus,
<https://industria.ub.ac.id/index.php/industri/article/download/5573/803>
50. Optimization of Pectin Extraction Process from Albedo of Citrus,
https://www.researchgate.net/publication/399391374_Optimization_of_Pectin_Extraction_Process_from_Albedo_of_Citrus_Lemon_Citrus_limon_Using_Ultrasonic_Method
51. Extraction of pectin from albedo of lemon peels for preparation of,
https://www.researchgate.net/publication/340977378_Extraction_of_pectin_from_albedo_of_lemon_peels_for_preparation_of_tissue_engineering_scaffolds
52. Extraction of Pectin from lemon peel: Technology development,
<https://scispace.com/pdf/extraction-of-pectin-from-lemon-peel-technology-development-e4kcbfso0y.pdf>
53. FT-IR spectroscopy, a reliable method for routine analysis of ... - Lirias,
<https://lirias.kuleuven.be/retrieve/ce99fb4b-aa3d-48d4-ad19-3864617e1a25>
54. FT-IR spectroscopy, a reliable method for routine analysis of the,

- <https://pubmed.ncbi.nlm.nih.gov/25624209/>
55. Improved method for determination of pectin degree of esterification,
<https://pubmed.ncbi.nlm.nih.gov/11409962/>
56. Determination of pectin degree of esterification by diffuse reflectance,
<https://www.semanticscholar.org/paper/Determination-of-pectin-degree-of-esterification-by-Gnanasambandam-Proctor/4e808146cab92e7c567aae1ccdf9215be7ff3338>
57. Determination of the esterification degree of the pectin of different,
https://www.researchgate.net/publication/368858635_Determination_of_the_esterification_degree_of_the_pectin_of_different_origin_and_composition_by_the_method_of_infrared_spectroscopy
58. Chemical and instrumental characterization of pectin from dried,
<https://www.redalyc.org/pdf/3030/303026597001.pdf>
59. Comparison of Analytical Methods for Determining ... - MDPI,
<https://www.mdpi.com/2076-3417/11/10/4461>
60. Optimization of method for extraction of pectin from apple pomace,
https://www.researchgate.net/publication/286409531_Optimization_of_method_for_extraction_of_pectin_from_apple_pomace
61. (PDF) KINETICS OF SUCROSE INVERSION - ResearchGate,
https://www.researchgate.net/publication/343970485_KINETICS_OF_SUCROSE_INVERSION
62. polarographic investigation of kinetics of inversion of sucrose,
<https://rasayanjournal.co.in/vol-3/issue-2/10.pdf>
63. Kinetics of Inversion of sugar cane or Sucrose - YouTube,
<https://www.youtube.com/watch?v=VR78Q6c3uk4>
64. The rate of reaction between sucrose and water catalyzed by,
https://www.uobabylon.edu.iq/eprints/publication_3_25208_250.pdf
65. Kinetics of Sucrose Inversion in Mixed Solvents - SciSpace,
<https://scispace.com/pdf/kinetics-of-sucrose-inversion-in-mixed-solvents-z8evwjiljh.pdf>
66. The science and magic of jam-making - The Guardian,
<https://www.theguardian.com/science/blog/2013/oct/03/science-magic-jam-making>
67. The Chemical Destruction of Sucrose, Fructose, and Glucose in Hot,
<https://assbt.org/wp-content/uploads/2024/02/JSBRVol13No5P406to414TheChemicalDestructionofSucroseFructoseandGlucoseinHotAlkalineProcessJuicesandLiquors.pdf>
68. The effect of sucrose inversion on boiling point elevation,
https://www.researchgate.net/publication/274964128_The_effect_of_sucrose_inversion_on_boiling_point_elevation
69. thermal degradation kinetics of total polyphenols,
<https://www.tf.ni.ac.rs/wp-content/uploads/casopis/2018/v1/c3.pdf>
70. Kinetics of the Degradation of Anthocyanins, Phenolic Acids and,
https://www.ftb.com.hr/images/pdfarticles/2014/January-March/ftb_52-1_101-108.pdf
71. Degradation Kinetics of Anthocyanin of Traditional and Low-Sugar,

- https://www.researchgate.net/publication/251246332_Degradation_Kinetics_of_Anthocyanin_of_Traditional_and_Low-Sugar_Blackberry_Jam
72. Influence of pectin properties and processing conditions on thermal, https://www.researchgate.net/publication/229423665_Influence_of_pectin_properties_and_processing_conditions_on_thermal_pectin_degradation
73. Influence of pectin properties and processing conditions on thermal, <https://lirias.kuleuven.be/312235>
74. Nonenzymatic Degradation of Citrus Pectin and Pectate during, https://www.academia.edu/33822534/Nonenzymatic_Degradation_of_Citrus_Pectin_and_Pectate_during_Prolonged_Heating_Effects_of_pH_Temperature_and_Degree_of_Methyl_Esterification
75. Nonenzymatic Degradation of Citrus Pectin and Pectate during, https://www.researchgate.net/publication/6286904_Nonenzymatic_Degradation_of_Citrus_Pectin_and_Pectate_during_Prolonged_Heating_Effects_of_pH_Temperature_and_Degree_of_Methyl_Esterification
76. [PDF] Nonenzymatic degradation of citrus pectin and pectate during, <https://www.semanticscholar.org/paper/Nonenzymatic-degradation-of-citrus-pectin-and-of-of-Diaz-Anthon/a8ad5935cb87bd5bbd37f4cfd471b28952158f19>
77. Current Advancements in Pectin: Extraction, Properties and ... - MDPI, <https://www.mdpi.com/2304-8158/11/17/2683>
78. Pectin depolymerization in jam #887942 - Ask Extension, <https://ask.extension.org/kb/faq.php?id=887942>
79. Effects of pectin with different molecular weight on gelatinization, https://www.researchgate.net/publication/324916384_Effects_of_pectin_with_different_molecular_weight_on_gelatinization_behavior_textural_properties_retrogradation_and_in_vitro_digestibility_of_corn_starch
80. Role of Sucrose in Pectin Gelation: Static and Dynamic Light, https://www.researchgate.net/publication/231694472_Role_of_Sucrose_in_Pectin_Gelation_Static_and_Dynamic_Light_Scattering_Experiments
81. Structure formation in sugar containing pectin gels, <https://d-nb.info/1205804226/34>
82. The complex mechanism of HM pectin gelation - Publications CNR API, <https://publications.cnr.it/api/v1/documents/download/186058>
83. (PDF) Role of cosolutes in gelation of high-methoxy pectin. Part 1, https://www.researchgate.net/publication/248425019_Role_of_cosolutes_in_gelation_of_high-methoxy_pectin_Part_1_Comparison_of_sugars_and_polyols
84. Microstructure and Kinetic Rheological Behavior of Amidated and, <https://pubs.acs.org/bomaf6/article/7/1/114/3503038/Microstructure-and-Kinetic-Rheological-Behavior-of>
85. Microstructure and Gelation Behaviour of High Methoxyl and Low, <https://research.chalmers.se/en/publication/6987>
86. Rheological behavior of high acyl gellan gum solution at gel point, <https://www.tandfonline.com/doi/full/10.1080/10942912.2017.1362431>
87. Colloidal Gelation Under Dynamic Perturbation: A Departure ... - MDPI, <https://www.mdpi.com/2310-2861/11/12/985>

88. "Gel Point". In: Encyclopedia of Polymer Science and Technology,
<http://nguyen.hong.hai.free.fr/EBOOKS/SCIENCE%20AND%20ENGINEERING/MECANIQUE/MATERIAUX/COMPOSITES/Encyclopedia%20of%20Polymer%20Science%20and%20Technology/Vol.10/Gel%20Point.pdf>
89. Gelation Kinetics from Rheological Experiments - TA Instruments,
<https://www.tainstruments.com/applications-notes/gelation-kinetics-from-rheological-experiments/>
90. (PDF) Analysis of Linear Viscoelasticity of a Crosslinking Polymer at,
https://www.researchgate.net/publication/243529348_Analysis_of_Linear_Viscoelasticity_of_a_Crosslinking_Polymer_at_the_Gel_Point
91. Evolution of Linear Viscoelasticity across the Critical Gelation ... - arXiv,
<https://arxiv.org/pdf/2603.26082>
92. Linear viscoelasticity of crosslinking polymers at the gel point,
<https://scispace.com/pdf/linear-viscoelasticity-of-crosslinking-polymers-at-the-gel-3zud49a0nz.pdf>
93. Ask the Expert: Using Rheology to Design Better Products—Yield,
<https://www.americanlaboratory.com/914-Application-Notes/117719-Ask-the-Expert-Using-Rheology-to-Design-Better-Products-Yield-Stress-and-How-to-Measure-It/>
94. Rheological investigation of food - Anton Paar Wiki,
<https://wiki.anton-paar.com/en/basics-of-rheology/rheological-investigation-of-food/>
95. Characterizing Gel Properties Using Oscillation Testing on a,
<https://analyzing-testing.netzsch.com/en/application-literature/characterizing-gel-properties-using-oscillation-testing-on-a-rotational-rheometer>
96. Temperature Compensation in Refractometers FAQ,
<https://www.misco.com/temperature-compensation-faq/>
97. Sensors and Instruments for Brix Measurement: A Review - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8951823/>
98. Brix Temperature Correction - UC Davis Viticulture & Enology,
<https://wineserver.ucdavis.edu/industry-info/enology/methods-and-techniques/winery-lab-techniques/brix-temperature-correction>
99. Refractometer Temperature Correction Vs Control - Rudolph Research,
<https://rudolphresearch.com/refractometer-temperature-correction/>
100. Refractometer | Brewfather - docs,
<https://docs.brewfather.app/tools/refractometer>
101. Do I need to adjust for temperature with a refractometer?,
<https://homebrew.stackexchange.com/questions/7494/do-i-need-to-adjust-for-temperature-with-a-refractometer>