

# The Science and Praxis of Traditional Barrel-Fermented Dill Pickles: Osmotic Dynamics, Microbial Succession, and Defect Prevention

The preservation of the cucumber (*Cucumis sativus*) via lactic acid fermentation represents a sophisticated biochemical transformation that merges microbial ecology, thermodynamic water activity, and cellular mechanics. Historically exemplified by the traditional practices of the Spreewald region in Brandenburg, Germany, the art of pickling in wooden barrels has been meticulously refined over centuries<sup>1</sup>. Cultivation in the Spreewald was largely established by Dutch settlers in the 14th century, leveraging the region's humid microclimate, iron oxide-rich water, and humus-dense soils to produce cucumbers of exceptional quality<sup>3</sup>. The resulting product achieved legendary status in the 19th century, famously championed by author Theodor Fontane, and today holds a Protected Geographical Indication (PGI) from the European Union<sup>2</sup>. While historic Spreewald fermentations required several weeks in large 500-gallon wooden barrels, transitioning this heritage practice into a highly reproducible, expert-level protocol requires the meticulous management of endogenous plant enzymes, precise osmotic calibration

via sodium chloride (NaCl) and calcium chloride ( $\text{CaCl}_2$ ), and the strict control of microbial succession<sup>2</sup>. The ultimate objective is to synthesize a product that retains exceptional cellular crispness while completely averting structural defects, particularly the catastrophic formation of internal gas cavities known as bloaters<sup>6</sup>.

## The Botanical Matrix: Cellular Architecture and Cultivar Selection

The physical texture of a fermented pickle is dictated entirely by the structural integrity of its cellular scaffolding. The primary cell wall of the cucumber consists of a complex network of crystalline cellulose microfibrils, formed by  $\beta$ -(1,4)-linked glucan chains, embedded within a highly hydrated matrix of hemicelluloses and pectins<sup>9</sup>.

### The Role of Pectin and the Middle Lamella

The most critical component for textural crispness is the middle lamella, a pectin-rich layer that acts as the primary adhesive binding adjacent plant cells together into a cohesive tissue

structure<sup>11</sup>. Pectin is a highly heterogeneous polysaccharide characterized by a backbone of  $\alpha$ -(1,4)-linked D-galacturonic acid residues, which can be heavily methyl-esterified<sup>9</sup>. During the growth phase of the cucumber, the methylesterification status of pectin controls cell wall porosity and plasticity, accommodating volumetric expansion driven by cellular turgor pressure<sup>9</sup>. In the

context of fermentation, the degree to which these pectin chains remain intact, or are successfully cross-linked by external divalent cations, defines the mechanical resistance and sensory juiciness of the final product<sup>11</sup>.

### Cultivar Influence on BLOATER Resistance

Not all *Cucumis sativus* varieties possess the genetic or structural predispositions necessary to survive the intense physical and osmotic stresses of fermentation. The susceptibility to balloon bloating—a condition where internal gas pressure separates the carpel walls—is negatively correlated with fruit size and heavily dependent on specific cultivar genetics<sup>16</sup>.

Extensive biomechanical testing, including compressed air injection protocols designed to measure the necessary force required to separate internal carpels, demonstrates significant variation among commercial pickling varieties<sup>16</sup>. To optimize fermentation outcomes, practitioners must select cultivars explicitly bred for dense internal structures and high carpel adhesion.

| Cultivar                          | Structural Characteristics                                             | Pickling Suitability                                                                                                                 |
|-----------------------------------|------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Vorgebirgstrauben                 | Thick, highly prickled German heritage variety; structurally dense.    | Optimal at lengths under four inches; highly resistant to carpel separation <sup>18</sup> .                                          |
| County Fair                       | Deep-green, arrow-straight hybrid; predominantly seedless if isolated. | Exceptional firmness; delayed maturation of the seed cavity reduces susceptibility to internal enzymatic degradation <sup>18</sup> . |
| Parisienne Cornichon de Bourbonne | Slender, numerous prickles, French origin.                             | Highly bloater-resistant even at extended lengths; ideal for high-density packing <sup>18</sup> .                                    |
| Wautoma                           | Open-pollinated, dark green, highly productive.                        | Demonstrates robust resistance to physical deformation during initial plasmolysis <sup>18</sup> .                                    |

Table 1: Structural evaluation of prominent *Cucumis sativus* cultivars regarding their suitability for hypertonic brine fermentation and resistance to carpel separation<sup>16</sup>.

### Enzymatic Degradation: Pectin Methylesterase and Polygalacturonase

Upon harvesting, mechanical stress triggers an immediate and pronounced physiological response within the cucumber tissue. This stress induces a transient burst of ethylene production, which acts as a signaling molecule to significantly upregulate the activity of cell-wall-degrading enzymes<sup>19</sup>. The two primary enzymatic agents responsible for the catastrophic loss of firmness in fermenting vegetables are pectin methylesterase (PME) and polygalacturonase (PG)<sup>9</sup>.

## **The Mechanism of Pectin Depolymerization**

The degradation of the middle lamella is a coordinated, sequential biochemical process. PME initiates the cascade by catalyzing the demethylesterification of the homogalacturonan chains within the pectin matrix<sup>14</sup>. This reaction removes methyl ester groups, releasing methanol and exposing negatively charged carboxyl groups along the polygalacturonic acid backbone<sup>10</sup>. While PME alone merely alters the electrostatic charge of the cell wall, its action is a prerequisite for the subsequent, far more destructive activity of polygalacturonase<sup>9</sup>. PG

functions by hydrolyzing the  $\alpha$ -(1,4)-glycosidic bonds between the newly demethylesterified galacturonic acid residues<sup>6</sup>. This outright cleavage depolymerizes the structural matrix, leading to cellular detachment, tissue maceration, and the irreversible softening of the cucumber<sup>11</sup>. Research indicates that PG activity exhibits distinct physiological optima, demonstrating maximum activity peaks at both pH 4.6 and pH 6.2, making it exceptionally dangerous during the initial and intermediate phases of lactic acid fermentation<sup>20</sup>.

## **Enzymatic Concentration and Mitigation Strategies**

The highest endogenous concentrations of PME and PG are localized within the blossom end of the cucumber<sup>23</sup>. The biological function of this enzymatic clustering is to eventually break down the distal tissue of the fruit to release mature seeds into the surrounding environment<sup>9</sup>. In commercial and expert artisanal pickling, the absolute removal of a 1/16-inch slice from the blossom tip of every cucumber is a mandatory processing step to physically excise the primary source of pectinolytic enzymes<sup>23</sup>.

Because these enzymes rapidly diffuse into the surrounding brine following mechanical injury, secondary mitigation strategies are essential<sup>26</sup>. Immediate hypothermic shock, achieved by submerging the harvested and trimmed cucumbers in an ice-water bath, drastically reduces the kinetic thermal energy of the system, temporarily suppressing PG activity until the fermentation environment becomes suitably hostile<sup>19</sup>.

## **The Vessel Matrix: Oak Cooperage and Phenolic Chemistry**

The utilization of traditional oak barrels for fermentation is not merely an adherence to historical aesthetics; the lignocellulosic structure of the wood is a biologically and chemically active participant in the maturation of the brine.

### **Stave Hydration and Osmotic Sealing**

Barrels constructed from air-dried, fine-grain European or American white oak inherently exist in

a state of fluctuating moisture<sup>28</sup>. When stored dry, the cellular structure of the wood contracts, resulting in macroscopic gaps between the staves<sup>29</sup>. Prior to the introduction of the primary fermentation matrix, the barrel must undergo a rigorous swelling protocol<sup>31</sup>.

This is accomplished by filling the vessel with hot, chlorine-free water (approximately 70°C to 82°C). The elevated thermal energy accelerates capillary action, forcing the wood fibers to rapidly absorb moisture<sup>30</sup>. As the staves swell, they generate immense structural tension against the galvanized steel hoops, hydro-mechanically sealing the intercellular gaps and rendering the barrel watertight<sup>30</sup>. Extended soaking with stagnant water must be strictly avoided, as it encourages the proliferation of undesirable molds or spoilage organisms prior to the acidification phase<sup>31</sup>. Instead, sterilization is typically achieved via high-pressure steam injection (maintaining internal temperatures of at least 85°C for 20 minutes) or through an acidic sulfur dioxide ( $SO_2$ ) wash utilizing tartaric acid and sulfur strips<sup>34</sup>.

## Phenolic Leaching and Tannin-Mediated Enzyme Inhibition

During the cooperage process, the interior of the barrel is subjected to thermal toasting, which partially pyrolyzes the wood's structural polymers. The thermal degradation of lignin produces volatile phenolic aldehydes such as vanillin and guaiacol, while the decomposition of specific

wood lipids forms oak lactones (e.g.,  $\beta$ -methyl- $\gamma$ -octalactone)<sup>36</sup>. As the hypertonic brine interacts with the toasted wood, these compounds slowly extract into the liquid, contributing profound aromatic depth to the fermented product<sup>36</sup>.

More crucially for the structural preservation of the cucumber, oak wood is exceptionally rich in hydrolyzable tannins<sup>36</sup>. Tannins are high-molecular-weight polyphenols characterized by their ability to precipitate proteins<sup>40</sup>. When these tannins diffuse from the oak staves into the fermentation brine, they actively bind to and denature the residual pectinolytic enzymes secreted by the cucumbers<sup>40</sup>. The introduction of auxiliary tannin sources—such as fresh grape leaves, oak leaves, or horseradish root layered among the cucumbers—provides an immediate, aggressive concentration of polyphenols that neutralize the active sites of polygalacturonase, serving as a passive, continuous biochemical defense against tissue softening<sup>27</sup>.

## Osmotic Dynamics: Precision Salinity and Calcium Fortification

The introduction of sodium chloride (NaCl) to the aqueous environment serves as the primary environmental master-switch for the entire biological trajectory of the fermentation<sup>6</sup>. The precise calibration of the brine establishes the necessary thermodynamic conditions to select for highly specific, aciduric microbiota while simultaneously reinforcing the cellular architecture of the fruit against degradation<sup>6</sup>.

## Mathematical Calibration of Total Salt Concentration (TSC)

A fundamental error in rudimentary pickling methodologies is the formulation of brine concentrations strictly by liquid volume (e.g., mixing a "5% brine solution"), failing to account for the intrinsic water content of the cucumbers, which typically exceeds 93%<sup>6</sup>. When placed in a

5% liquid brine, rapid osmotic equilibration forces unbound water out of the cucumber cells and salt into the tissue. Within 48 hours, the localized brine concentration experiences massive dilution, frequently falling below the threshold necessary for pathogen inhibition<sup>6</sup>. For absolute microbiological control and consistent product quality, salinity must be calculated as a Total Salt Concentration (TSC), encompassing the combined mass of both the vegetable matrix and the added water<sup>42</sup>. The optimal TSC for cucumber fermentation lies in a precise biological compromise between 2.5% and 3.0%<sup>6</sup>.

$$Salt_{mass} = (Water_{mass} + Cucumber_{mass}) \times \text{Target TSC (e.g., 0.025)}$$

Employing a rigorous, mass-based mathematical approach eliminates the volumetric discrepancies caused by the variable density of different salt crystal structures, whether dense, subterranean-mined vacuum pan salt or low-density, hollow flake salt<sup>6</sup>.

### The Salometer Scale and Brine Density

In commercial operations, brine density is continuously validated using a hydrometer specifically calibrated to the Salometer (or Salinometer) scale<sup>45</sup>. By definition, 0 degrees Salometer (0°S) represents pure water at 15°C (60°F), while 100°S represents a fully saturated sodium chloride solution containing 26.395% salt by weight<sup>45</sup>.

| Salometer Degrees (°S) | Specific Gravity (15°C) | NaCl % by Weight | Grams of NaCl per Liter | Fermentation Phase               |
|------------------------|-------------------------|------------------|-------------------------|----------------------------------|
| 10°S                   | 1.019                   | 2.64%            | 26.4g                   | Rapid half-sour initiation       |
| 15°S                   | 1.029                   | 3.96%            | 39.6g                   | Standard full-sour equilibrium   |
| 20°S                   | 1.039                   | 5.28%            | 52.8g                   | Traditional high-salt initiation |
| 25°S                   | 1.050                   | 6.60%            | 66.0g                   | Pre-commercial storage phase     |

Table 2: Salometer conversions demonstrating the precise relationship between specific gravity, salt saturation, and their application phases in cucumber pickling<sup>45</sup>.

## Calcium Chloride ( $CaCl_2$ ) and the Egg-Box Model

While NaCl establishes the primary osmotic pressure gradient, relying upon it entirely often fails to preserve maximal crispness during prolonged storage in highly acidic conditions<sup>43</sup>. Modern preservation chemistry mandates the supplementation of the brine with divalent cations,

specifically calcium ( $Ca^{2+}$ ), introduced via calcium chloride ( $CaCl_2$ )<sup>6</sup>.

The introduction of  $Ca^{2+}$  ions biochemically capitalizes on the preliminary action of pectin methylesterase. As PME demethylates the pectin chains, it exposes the negatively charged carboxyl groups<sup>9</sup>. The free  $Ca^{2+}$  ions migrate deep into the cell wall matrix and form highly stable electrostatic bonds between adjacent parallel pectin chains<sup>6</sup>. This phenomenon, formally recognized as the "egg-box" model of calcium-pectate cross-linking, physically reinforces the middle lamella, creating an impenetrable structural barrier against further polygalacturonase degradation<sup>6</sup>. Diffusion kinetics demonstrate that calcium uptake into the cucumber tissue is

significantly accelerated when the brine is supra-osmotic (containing both NaCl and  $CaCl_2$ ) compared to iso-osmotic solutions, ensuring rapid and even distribution of the calcium throughout the mesocarp and endocarp<sup>51</sup>.

However, the application of  $CaCl_2$  requires strict biochemical titration. Elevated levels of calcium chloride impart highly objectionable bitter, metallic, and astringent organoleptic defects to the final product<sup>52</sup>. Sensory threshold analyses indicate that the 50% detection threshold for  $CaCl_2$  bitterness in dill pickles is  $61.8 \pm 7.6$  mM<sup>52</sup>. Therefore, optimal structural firming

without sensory degradation is achieved by targeting an equilibrated  $CaCl_2$  concentration of 0.05% to 0.1% (approximately 10 mM to 40 mM) within the final fermentation matrix<sup>43</sup>.

## Microbial Succession and Acidification Kinetics

The introduction of the hypertonic, calcium-fortified brine triggers immediate cellular plasmolysis in the cucumbers, forcefully drawing out unbound water, soluble hexose carbohydrates (glucose and fructose), amino acids, and micronutrients<sup>6</sup>. This nutrient-dense exudate becomes the immediate substrate for a fierce, highly sequential microbial succession<sup>6</sup>.

### Phase 1: Osmotic Shock and the Enterobacteriaceae Purge

In the initial 24 to 48 hours of fermentation, the brine maintains residual dissolved oxygen and a near-neutral pH (approximately 6.5)<sup>6</sup>. Under these conditions, transient, indigenous populations of Gram-negative *Enterobacteriaceae* (e.g., *Enterobacter cancerogenus*, *Enterobacter nimipressuralis*) attempt to aggressively metabolize the leached sugars<sup>55</sup>.

However, the Gram-negative cellular architecture—characterized by a thin peptidoglycan layer enclosed between two lipid membranes—is exceptionally susceptible to osmotic shock<sup>6</sup>. The hypertonic 2.5% to 3.0% TSC environment forces a massive, instantaneous efflux of intracellular water from their cytoplasm, causing a sudden and lethal collapse in turgor

pressure<sup>6</sup>. As the surviving, halotolerant lactic acid bacteria (LAB) begin generating early organic acids, the rapidly declining pH permanently suppresses and eradicates these transient coliforms, rendering the matrix microbiologically secure and degrading the initial "nitrite peak" associated with their brief metabolic activity<sup>6</sup>.

## **Phase 2: Heterofermentative Dominance (*Leuconostoc mesenteroides*)**

As the initial respiration depletes the available oxygen, the environment transitions to an obligate anaerobic state. The ecological niche is swiftly dominated by *Leuconostoc mesenteroides*, an obligately heterofermentative lactic acid bacterium uniquely adapted to outcompete competitors at high salinities and lower temperatures (15°C - 20°C)<sup>6</sup>.

Because *L. mesenteroides* lacks the critical enzyme aldolase, it cannot utilize the standard Embden-Meyerhof-Parnas glycolytic pathway. Instead, it metabolizes hexose sugars via the phosphoketolase pathway, producing an equimolar ratio of D-lactic acid, acetic acid, ethanol,

and large volumes of carbon dioxide ( $CO_2$ )<sup>6</sup>. The vigorous evolution of  $CO_2$  actively purges any remaining dissolved oxygen from the brine, solidifying the strict anaerobic conditions required for subsequent maturation<sup>6</sup>.

Simultaneously, in environments rich in sucrose, *L. mesenteroides* deploys extracellular dextransucrase enzymes (such as glucose glucosyltransferase) to synthesize massive, branched exopolysaccharide polymers known as dextrans<sup>6</sup>. While excessive dextran production can lead to a transiently viscous or "ropy" brine, controlled amounts act as functional prebiotics and enhance the rheological mouthfeel of the liquid<sup>6</sup>. The combination of lactic and acetic acids synthesized by *L. mesenteroides* rapidly depresses the ambient pH from 6.5 to roughly 4.5 within 3 to 5 days<sup>6</sup>.

## **Phase 3: Homofermentative Climax (*Lactiplantibacillus plantarum*)**

As the pH falls below 4.5, *L. mesenteroides* becomes increasingly inhibited by the toxicity of its own acidic exudates, eventually ceasing growth entirely<sup>6</sup>. The microbiome abruptly transitions to a homofermentative climax, dominated overwhelmingly by highly aciduric species, principally *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) and *Lactiplantibacillus pentosus*<sup>6</sup>.

*L. plantarum* employs the highly efficient Embden-Meyerhof-Parnas (EMP) pathway, converting the remaining freely available carbohydrates almost exclusively into lactic acid<sup>6</sup>. This aggressive homofermentative action drives the pH down to a terminal, shelf-stable range of 3.2 to 3.8<sup>6</sup>.

To survive the extreme, self-generated acidity—where the external proton gradient exerts immense pressure on the cellular membrane—*L. plantarum* actively regulates its intracellular

pH. It achieves this by deploying  $F_1F_0$ -ATPase proton pumps located in its cytoplasmic

membrane. These multi-subunit complexes actively extrude invading hydrogen ions ( $H^+$ ) back into the extracellular brine, a mechanism powered directly by the continuous hydrolysis of cellular ATP, allowing the bacteria to maintain a nearly neutral internal pH despite the hostile

external environment<sup>6</sup>.

| Fermentation Phase                     | Dominant Microbiota                                       | Primary Metabolic Pathway    | Key Exudates                                          | pH Shift              |
|----------------------------------------|-----------------------------------------------------------|------------------------------|-------------------------------------------------------|-----------------------|
| Phase 1: Initiation (Days 1-2)         | <i>Enterobacteriaceae</i> (Transient)                     | Diverse heterofermentative   | $CO_2$ , minor organic acids                          | 6.5 $\rightarrow$ 5.5 |
| Phase 2: Heterofermentative (Days 3-7) | <i>Leuconostoc mesenteroides</i>                          | Phosphoketolase              | D-Lactic acid, Acetic acid, Ethanol, $CO_2$ , Dextran | 5.5 $\rightarrow$ 4.5 |
| Phase 3: Climax (Days 8-30+)           | <i>Lactiplantibacillus plantarum</i> , <i>L. pentosus</i> | Embden-Meyerhof-Parnas (EMP) | L- and D-Lactic acid (racemic mixture)                | 4.5 $\rightarrow$ 3.2 |

Table 3: The chronological microbial succession and corresponding biochemical outputs defining the lactic acid fermentation of cucumbers<sup>6</sup>.

## Defect Mitigation: The Etiology and Prevention of BLOATER Damage

The most catastrophic economic and structural failure in the bulk cucumber fermentation industry is the development of "bloaters" damage<sup>8</sup>. Bloaters are cucumbers that develop massive internal hollow cavities, classified morphologically as balloon, lens, or honeycomb bloaters, depending on the severity and pattern of the internal tissue rupture<sup>7</sup>.

### The Physics of Internal Gas Pressure

The fundamental etiology of the bloater defect lies in the accumulation of microbiologically generated carbon dioxide ( $CO_2$ ) within the highly restrictive geometry of the pickling vessel<sup>8</sup>.

During the early stages of fermentation, the rapid evolution of  $CO_2$  by surviving *Enterobacteriaceae* and the explosive heterofermentative activity of *L. mesenteroides* saturates the external brine<sup>55</sup>.

Concurrently, *L. plantarum*—despite being homofermentative regarding hexose sugars—possesses a highly active malic acid decarboxylase (MDC) enzyme. This enzyme actively decarboxylates the naturally occurring malic acid locked within the cucumber tissue,

generating supplemental massive volumes of  $CO_2$  and lactic acid<sup>7</sup>. Because carbon dioxide is highly soluble in aqueous solutions, it rapidly diffuses across the semi-permeable epidermis of the cucumber and infiltrates the plant's internal intercellular spaces, which are naturally filled with highly insoluble nitrogen gas<sup>8</sup>.

As the external  $CO_2$  moves into the fruit at a rate far exceeding the outward diffusion of the internal nitrogen, an immense internal hydrostatic pressure gradient develops<sup>8</sup>. Once this internal pressure exceeds the mechanical tensile strength of the mesocarp and endocarp tissues, the carpels violently separate, physically tearing the interior of the cucumber apart to create large, hollow gas pockets<sup>8</sup>. Research clearly demonstrates that bloater incidence is

directly correlated to the percentage of  $CO_2$  saturation in the brine, with severe damage occurring rapidly when saturation exceeds 50%<sup>70</sup>.

### Strategic Interventions for BLOATER Prevention

To preserve the dense, crisp architecture of the cucumber against this thermodynamic pressure, expert practitioners deploy an array of precise physical, chemical, and microbiological interventions to manage gas equilibrium.

**Mechanical Venting (Needle Pricking):** The most direct physical intervention involves the micro-perforation of the cucumber epidermis using specialized stainless steel needle arrays. Puncturing the skin longitudinally immediately establishes a mechanical exhaust pathway,

allowing trapped  $CO_2$  and insoluble nitrogen to easily vent into the external brine before fatal internal pressure gradients can accumulate<sup>71</sup>.

**Nitrogen Purging:** Adapted directly from industrial United States Department of Agriculture (USDA) Agricultural Research Service (ARS) protocols developed by Etchells and Fleming, advanced artisanal and commercial producers utilize a continuous or intermittent bubbling of nitrogen ( $N_2$ ) gas through the fermentation matrix<sup>8</sup>. The mechanical agitation of the  $N_2$

bubbles physically sweeps the dissolved  $CO_2$  out of the brine, driving the  $CO_2$  concentration safely below the critical threshold (less than 20 mg/100 mL) required to induce bloater damage<sup>17</sup>. It is critical to note that air purging is actively discouraged; the introduction of oxygen heavily promotes the proliferation of highly oxidative surface molds and film yeasts (kahm yeast). These aerobic organisms secrete aggressive pectin depolymerase enzymes that subsequently diffuse downward into the brine, leading to devastating top-down tissue softening<sup>73</sup>.

**Biological Control via Starter Cultures:** Advanced bioprocessing relies on the targeted inoculation of the brine with specific bacterial strains designed to outcompete natural gas producers or eliminate specific enzymatic pathways. The application of isolated *L. plantarum* mutant strains that are entirely deficient in malic acid decarboxylase (MDC- strains) prevents the secondary evolution of  $CO_2$  from the fruit's endogenous malic acid reserves<sup>56</sup>. Alternatively, co-inoculation with *Lactococcus lactis*, an organism possessing an exceptionally rapid doubling

time, quickly establishes competitive exclusion against early  $CO_2$ -producing *Enterobacteriaceae* and *Leuconostoc* species, dramatically lowering the overall bloater index of the batch<sup>75</sup>.

## Biochemical Safety and Pathogen Eradication

The lacto-fermentation of cucumbers is inherently self-sterilizing, providing an impeccable food safety record that spans millennia. The preservation mechanism relies not merely on the measurement of ambient hydrogen ion concentration (ambient pH), but specifically on the accumulation of weak organic acids through a biological phenomenon known as the "proton effect"<sup>6</sup>.

### The Proton Effect and Weak Organic Acids

Unlike strong mineral acids (e.g., hydrochloric acid) that fully dissociate in aqueous solutions, weak organic acids like lactic acid ( $pK_a = 3.86$ ) and acetic acid ( $pK_a = 4.76$ ) only partially dissociate<sup>6</sup>. Governed by the Henderson-Hasselbalch equation, when the brine pH falls near or below these specific  $pK_a$  values, a significant proportion of the acid molecules exist in an undissociated, lipophilic (uncharged) state<sup>6</sup>.

These lipophilic molecules easily diffuse across the lipid bilayer membranes of invading pathogenic bacteria, such as *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella* species<sup>6</sup>. Once inside the pathogen's cytoplasm, where the internal pH is maintained near neutrality (~7.0), the organic acids instantly encounter an environment well above their  $pK_a$ .

Consequently, they rapidly dissociate, dumping massive amounts of free protons ( $H^+$ ) and acid anions (lactate and acetate) directly into the cell interior<sup>6</sup>. This catastrophic internal acidification destroys the transmembrane electrochemical gradient (the proton motive force), paralyzing the pathogen's ability to synthesize ATP or transport nutrients, resulting in rapid metabolic exhaustion and cell death<sup>6</sup>.

### Absolute Mitigation of *Clostridium botulinum*

The primary, systemic microbiological hazard universally associated with anaerobic food preservation is the proliferation of *Clostridium botulinum*, an obligate anaerobic, spore-forming bacterium responsible for synthesizing the lethal botulinum neurotoxin<sup>77</sup>.

However, the fundamental biochemistry of *C. botulinum* dictates a strict, absolute minimum pH threshold of 4.6 for both cellular growth and toxin synthesis<sup>77</sup>. The immense safety profile of lacto-fermentation is derived directly from the rapid acidification kinetics of *L. mesenteroides* and *L. plantarum*, which reliably drive the ambient pH of the brine well below 4.6 within the first 48 to 72 hours<sup>76</sup>. Because the hypertonic saline environment significantly delays spore germination during the initial hours of fermentation, by the time the *C. botulinum* spores attempt to transition into vegetative cells, the matrix has already crossed the fatal pH 4.6 threshold, rendering neurotoxin production biologically impossible<sup>76</sup>.

# Praxis: The Master Protocol for Barrel-Fermented Pickles

The following protocol synthesizes the thermodynamic, enzymatic, and microbiological principles detailed above into a rigorous, reproducible methodology for producing traditional, hollow-free dill pickles of exceptional crispness.

## Phase 1: Hydration and Sanitation of the Oak Matrix

The process commences with the procurement of a traditional, untreated white oak barrel, typically ranging from 10 to 50 liters in volume. Because the staves arrive in a contracted state, the vessel must be completely filled with unchlorinated, hot water (approximately 75°C). The barrel is allowed to rest undisturbed for 24 to 48 hours. During this period, the intense thermal energy forces the lignocellulosic fibers to rapidly absorb moisture, expanding against the galvanized hoops until all external weeping ceases and the joints are hydro-mechanically sealed<sup>29</sup>.

Following successful hydration, the water is entirely expelled. To eliminate residual, competitive fungi and wild *Brettanomyces* yeasts residing within the porous wood grain, the interior is sanitized. This is achieved via high-pressure steam injection, maintaining an internal surface temperature of at least 85°C for 20 minutes, or alternatively, by employing an acidic sulfur dioxide wash comprising 10 grams of tartaric acid per liter of water, followed by the combustion of a standard sulfur strip<sup>33</sup>. The barrel is then thoroughly flushed with sterile, cold water to remove any chemical residue that might interfere with the LAB.

## Phase 2: Enzymatic Deactivation and Substrate Assembly

Freshly harvested, small-to-medium pickling cucumbers, specifically selected from bloater-resistant cultivars such as Vorgebirgstrauben or Parisienne Cornichon de Bourbonne, are assembled<sup>18</sup>. Uniformity in size is critical to ensure predictable osmotic equilibration across the entire batch.

Using a sanitized blade, exactly 1/16th of an inch is physically severed from the blossom end of every cucumber. This non-negotiable action eradicates the concentrated nodes of pectin methylesterase and polygalacturonase enzymes, decisively halting the primary biological trigger for structural degradation<sup>23</sup>. The trimmed cucumbers are immediately submerged in an ice-water bath (0°C to 4°C) for a minimum of two hours. This profound hypothermic shock dramatically retards any residual enzymatic action and maximizes cellular turgor pressure prior to encountering the hypertonic brine<sup>19</sup>.

To preclude the accumulation of microbiologically generated carbon dioxide and the subsequent formation of internal gas cavities, each cucumber is systematically pierced two to three times longitudinally using a sterile needle array. These micro-perforations serve as critical mechanical exhaust vents, equalizing the internal and external hydrostatic pressures<sup>71</sup>.

## Phase 3: Osmotic Calibration and Brine Formulation

The precise formulation of the brine demands rigorous mathematical calculation. The total mass of the cucumbers to be packed into the barrel is determined, followed by a volumetric calculation of the water required to fully submerge them, thereby yielding the total matrix weight.

To achieve optimal microbiological control while initiating cellular plasmolysis, the Total Salt Concentration (TSC) is targeted at exactly 2.5%. The combined mass of the cucumbers and water is multiplied by 0.025 to dictate the precise mass (in grams) of unrefined, non-iodized sodium chloride required<sup>6</sup>.

To exploit the "egg-box" model of cross-linking and guarantee maximal crispness, food-grade calcium chloride ( $CaCl_2$ ) is integrated into the solution. The concentration is strictly limited to 0.05% by mass of the total matrix. This precise titration ensures significant stabilization of the demethylated pectin chains while remaining safely below the 61.8 mM threshold that would otherwise impart severe metallic and bitter organoleptic defects<sup>52</sup>.

#### Phase 4: Packing and Atmospheric Sealing

The foundation of the barrel is layered with fresh, tannin-rich oak or grape leaves. As the tannins leach into the brine, they actively precipitate and permanently bind to any surviving polygalacturonase, reinforcing the structural defense<sup>27</sup>. Auxiliary aromatics, including mature dill umbels, whole garlic cloves, horseradish root, and mustard seeds, are distributed throughout the vessel to establish the classic flavor profile.

The chilled, mechanically vented cucumbers are packed tightly into the oak barrel, intentionally minimizing interstitial void space. The thoroughly dissolved, ambient-temperature NaCl and  $CaCl_2$  brine is poured over the assembly until complete submergence is achieved. An anaerobic seal is finalized by placing a sanitized wooden follower board or traditional ceramic weight over the vegetables, ensuring they remain entirely submerged below the brine surface. Any exposure to atmospheric oxygen will immediately promote the growth of highly oxidative surface molds that actively digest lactic acid, thereby driving the pH upward and precipitating systemic putrefaction<sup>6</sup>.

#### Phase 5: Fermentation Kinetics and Maturation

The ambient environment is strictly controlled, maintaining a temperature band between 20°C and 22°C. Temperatures exceeding 25°C dangerously accelerate the heterofermentative *L.*

*mesenteroides* phase, generating uncontrollable volumes of  $CO_2$  that exacerbate the risk of bloater damage and induce undesirable off-flavors<sup>6</sup>.

Throughout the initial 7 to 10 days, vigorous effervescence indicates the active purging of carbon dioxide. The barrel bung is fitted with a proper airlock mechanism to permit gas escape while entirely preventing atmospheric oxygen ingress<sup>8</sup>. Acidification is continuously monitored utilizing a calibrated digital pH meter. The brine must cross the absolute safety threshold of pH 4.6 within 72 hours, effectively neutralizing any threat of *Clostridium botulinum* spore germination<sup>76</sup>.

The primary homofermentative climax is typically realized within three to four weeks, identifiable when the pH definitively stabilizes between 3.2 and 3.8. This specific metric signals the successful, comprehensive conversion of available hexose sugars into preservative lactic acid<sup>6</sup>. Finally, the sealed barrel is transferred to a hypothermic cellar environment (4°C to 10°C). Immersed in profound acidity, fortified by complex oak-derived lactones, and bound by a robust

calcium-pectate matrix, the cucumbers remain in a state of crisp, microbiologically sterile suspension, capable of enduring up to 12 months of storage without degradation.

## Works cited

1. Spreewälder Gurken | Local Pickled Cucumber From Spreewald, <https://www.tasteatlas.com/spreewalder-gurken>
2. Spreewald gherkins - Wikipedia, [https://en.wikipedia.org/wiki/Spreewald\\_gherkins](https://en.wikipedia.org/wiki/Spreewald_gherkins)
3. A short history of the Spreewald Pickle - Berlin Journal, <https://www.berlinjournal.biz/en/short-history-spreewald-pickle/>
4. On the gherkin trail - The Berliner, <https://www.the-berliner.com/berlin/on-the-gherkin-trail/>
5. Gurkenmuseum (Cucumber Museum) in Lübbenau/Spreewald, <https://www.atlasobscura.com/places/das-gurkenmuseum-the-cucumber-museum>
6. 93 Vegetable Lacto-Fermentation Process.pdf
7. Fermenting Cucumbers: A Traditional Method for Pickles, <https://agriculture.institute/food-chemistry-and-physiology/fermenting-cucumbers-traditional-pickle-method/>
8. Research Perspectives - USDA ARS, <https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p172.pdf>
9. Restoration of Mature Etiolated Cucumber Hypocotyl Cell Wall, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2492596/>
10. Biochemical and Molecular Characterization of Cell Wall Degrading, <https://scialert.net/fulltext/?doi=ajbkr.2017.1.23>
11. Pectinesterase Activity, Pectin Methylation, and Texture Changes, [https://www.researchgate.net/publication/240662717\\_Pectinesterase\\_Activity\\_Pectin\\_Methylation\\_and\\_Texture\\_Changes\\_During\\_Storage\\_of\\_Blanched\\_Cucumber\\_Slices](https://www.researchgate.net/publication/240662717_Pectinesterase_Activity_Pectin_Methylation_and_Texture_Changes_During_Storage_of_Blanched_Cucumber_Slices)
12. Plant Cell Wall Degrading Enzymes, Pectinase and Cellulase, in the, <https://pps.agriculturejournals.cz/pdfs/pps/2014/04/06.pdf>
13. Influences on Cell Wall Assembly and Structure, Plant Development, <https://www.annualreviews.org/content/journals/10.1146/annurev-arplant-083023-034055?crawler=true&mimetype=application/pdf>
14. Pectin Methylesterases: Cell Wall Remodeling Proteins Are, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2018.01612/full>
15. Understanding pectin cross-linking in plant cell walls - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11748717/>
16. A compressed air test for carpel adhesion in pickling cucumbers, <https://cucurbitbreeding.wordpress.ncsu.edu/files/2016/04/a-compressed-air-test-for-carpel-adhesion-in-pickling-cucumbers.pdf>
17. Evaluation of Bloat Resistance in Pickling Cucumbers Using a, <https://journals.ashs.org/downloadpdf/view/journals/jashs/109/2/article-p261.pdf>
18. Cucumbers for Pickling: They're Not All Alike - A Gardener's Table, <https://agardenerstable.com/cucumbers-for-pickling-theyre-not-all-alike/>

19. Mechanical Stress, Storage Time, and Temperature Influence Cell,  
<https://journals.ashs.org/downloadpdf/view/journals/jashs/112/4/article-p666.pdf>
20. Polygalacturonase Activity and Ethylene Synthesis ... - USDA ARS,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p146.pdf>
21. Different Activities of Pectin-degrading Enzymes Affecting the,  
<https://www.hst-j.org/articles/xml/JzDR/>
22. Potential Role of Native Pickling Cucumber Polygalacturonase in,  
[https://www.researchgate.net/publication/224767741\\_Potential\\_Role\\_of\\_Native\\_Pickling\\_Cucumber\\_Polygalacturonase\\_in\\_Softening\\_of\\_Fresh\\_Pack\\_Pickles](https://www.researchgate.net/publication/224767741_Potential_Role_of_Native_Pickling_Cucumber_Polygalacturonase_in_Softening_of_Fresh_Pack_Pickles)
23. The Part Of The Cucumber You Must Remove When Making Pickles,  
<https://www.tastingtable.com/1079573/the-part-of-the-cucumber-you-must-remove-when-making-pickles/>
24. Soft pickles? Here's a fix. - YouTube,  
<https://www.youtube.com/shorts/-SNpwdFkSiQ>
25. Growing Cucumbers for Pickling? Cut Off Those Blossom Ends!,  
<https://www.groworganic.com/blogs/articles/cut-those-blossom-ends-off-your-cucumbers>
26. Inhibition of Pectinolytic and Cellulolytic Enzymes in Cucumber,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p79.pdf>
27. If I want crispy cucumbers, how do I cut them? : r/fermentation - Reddit,  
[https://www.reddit.com/r/fermentation/comments/1k1n256/if\\_i\\_want\\_crispy\\_cucumbers\\_how\\_do\\_i\\_cut\\_them/](https://www.reddit.com/r/fermentation/comments/1k1n256/if_i_want_crispy_cucumbers_how_do_i_cut_them/)
28. EDER – FassStolz® 1000 l Moselfuder - German Oak - barrel-shop,  
<https://www.barrel-shop.com/EDER-FassStolz-1000-l-Moselfuder-German-Oak/2275>
29. How to fix a leaking wine barrel — causes, fixes and when to call for,  
<https://www.liquidlux.com.au/blogs/liqlux-academy/fixing-a-leaking-home-barrel-a-step-by-step-guide>
30. Curing Oak Keg - Essential Distilling,  
<https://www.essentialdistilling.com/resources/Curing-Oak-Keg>
31. Step-by-Step Professional Protocols (Fall Edition) - Wooden Barrels,  
<https://barrelswood.com/blog/barrel-seasoning-preparation-step-by-step-professional-protocols-fall-edition/>
32. The 10 mistakes when using a wooden barrel - Barriles La Arboleda,  
<https://www.barrileslaarboleda.com/2019/10/09/the-10-mistakes-when-using-a-wooden-barrel/>
33. Barrel Care | Ensure Barrel Longevity - Pinnacle Oak,  
<https://www.pinnacleoak.com.au/barrel-care>
34. Cleaning, storage and maintenance of barrels,  
[https://www.awri.com.au/industry\\_support/winemaking\\_resources/storage-and-packaging/packaging-operations/barrel-cleaning-storage-and-maintenance/](https://www.awri.com.au/industry_support/winemaking_resources/storage-and-packaging/packaging-operations/barrel-cleaning-storage-and-maintenance/)
35. OAK BARREL MAINTENANCE GUIDE - FermeTech,  
[https://www.fermetech.eu/media/cidermill/product/cdn-old.erply.com/1323-1323\\_644b7dc0a48531.10193075\\_oak-barrel-maintenance-guide.pdf](https://www.fermetech.eu/media/cidermill/product/cdn-old.erply.com/1323-1323_644b7dc0a48531.10193075_oak-barrel-maintenance-guide.pdf)
36. Red Wine Aging in Oak Barrels: The Influence of Toasting Levels on,

- <https://www.mdpi.com/2306-5710/11/6/165>
37. How oak barrels give flavour to spirits, wines and beers - Luso Barrel,  
<https://lusobarrel.com/2023/03/22/how-oak-barrels-give-flavour-to-spirits-wines-and-beers/>
  38. Vanillin — Phenolic Aldehyde from Lignin Degradation,  
<https://www.winewithseth.com/winewiki/vanillin/>
  39. The Science of Aging: Understanding Phenolic and Flavor ... - PMC,  
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12346643/>
  40. The science on tannins from grape leaves in fermented cucumbers,  
[https://www.reddit.com/r/fermentation/comments/vja112/the\\_science\\_on\\_tannins\\_from\\_grape\\_leaves\\_in/](https://www.reddit.com/r/fermentation/comments/vja112/the_science_on_tannins_from_grape_leaves_in/)
  41. Evaluation of enzymatic and non-enzymatic softening in low salt,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p358.pdf>
  42. Brine calculator — salt percentage for pickles, peppers, and lacto,  
<https://fermentcalc.com/brine/>
  43. Enhancing the Texture and Sensory Properties of Pickled ... - MDPI,  
<https://www.mdpi.com/2304-8158/14/3/336>
  44. Salt Concentration Chart for Vegetable Pickles - MakeSauerkraut,  
<https://www.makesauerkraut.com/salt-brine-for-pickles/>
  45. Salometer Guide for Brine Measurement | PDF | Weight - Scribd,  
<https://fr.scribd.com/document/73567264/Salometer-Usage>
  46. Tools for Measuring Brine Strength - A Gardener's Table,  
<https://agardenerstable.com/tools-for-measuring-brine-strength/>
  47. Salt Per Water Brine Calculator - The CheeseMaker,  
<https://www.thecheesemaker.com/salt-per-water-brine-calculator/>
  48. Sodium Chloride Table for 60°F (15.5°C) - BrineMaker,  
<https://www.brinemaker.com/wp-content/uploads/SodiumChlorideTable.pdf>
  49. Fermentation of Cucumbers Brined with Calcium Chloride Instead of,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p366.pdf>
  50. Inhibitory Effects of CaCl<sub>2</sub> and Pectin Methylesterase on Fruit ... - MDPI,  
<https://www.mdpi.com/2311-7524/8/1/1>
  51. WM WALTER, JR.<sup>2</sup>, HP FLEMING, RL THOMPSON and TI FINE<sup>3</sup>,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Sweetpotato%20Publications/s99.pdf>
  52. Consumer Acceptability of Cucumber Pickles Produced by,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p395.pdf>
  53. use of calcium chloride to improve the texture of pickles<sup>1,2</sup>,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p133.pdf>
  54. Buffer Models for pH and Acid Changes Occurring in Cucumber,  
[https://www.researchgate.net/publication/361225563\\_Buffer\\_models\\_for\\_pH\\_and\\_acid\\_changes\\_occurring\\_in\\_cucumber\\_juice\\_fermented\\_with\\_Lactiplantibacillus\\_pentosus\\_and\\_Leuconostoc\\_mesenteroides](https://www.researchgate.net/publication/361225563_Buffer_models_for_pH_and_acid_changes_occurring_in_cucumber_juice_fermented_with_Lactiplantibacillus_pentosus_and_Leuconostoc_mesenteroides)
  55. CO<sub>2</sub>-mediated bloater defect can be induced by the uncontrolled,  
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10563668/>
  56. Bloater formation in brined cucumbers by *Lactobacillus plantarum*,

- [https://www.researchgate.net/publication/230061895\\_Bloater\\_formation\\_in\\_brined\\_cucumbers\\_by\\_Lactobacillus\\_plantarum](https://www.researchgate.net/publication/230061895_Bloater_formation_in_brined_cucumbers_by_Lactobacillus_plantarum)
57. Wo bekommt man die besten Gurken in der Stadt? - Berlin - Reddit,  
[https://www.reddit.com/r/berlin/comments/wg9mjx/where\\_to\\_get\\_the\\_best\\_pickles\\_in\\_the\\_city/?tl=de](https://www.reddit.com/r/berlin/comments/wg9mjx/where_to_get_the_best_pickles_in_the_city/?tl=de)
  58. Acid Tolerance of *Leuconostoc mesenteroides* and *Lactobacillus*,  
<https://pmc.ncbi.nlm.nih.gov/articles/PMC184570/>
  59. (PDF) Lactic Acid Bacteria Profile of Fermenting Cucumber in 7,  
[https://www.researchgate.net/publication/333148956\\_Lactic\\_Acid\\_Bacteria\\_Profile\\_of\\_Fermenting\\_Cucumber\\_in\\_7\\_Brine\\_Solution](https://www.researchgate.net/publication/333148956_Lactic_Acid_Bacteria_Profile_of_Fermenting_Cucumber_in_7_Brine_Solution)
  60. Fermentation kinetics of *Lactobacillus plantarum*, *Leuconostoc*,  
[https://www.researchgate.net/figure/Fermentation-kinetics-of-Lactobacillus-plantarum-Leuconostoc-mesenteroides-and\\_fig1\\_353893574](https://www.researchgate.net/figure/Fermentation-kinetics-of-Lactobacillus-plantarum-Leuconostoc-mesenteroides-and_fig1_353893574)
  61. Causes and Possible Solutions for Problems with Fermented Pickles,  
<https://nchfp.uga.edu/how/ferment/general-information-on-fermenting/causes-and-possible-solutions-for-problems-with-fermented-pickles/>
  62. Optimization and characterization of dextranucrase production by,  
[https://www.researchgate.net/publication/344461429\\_Optimization\\_and\\_characterization\\_of\\_dextranucrase\\_production\\_by\\_local\\_Leuconostoc\\_mesenteroides](https://www.researchgate.net/publication/344461429_Optimization_and_characterization_of_dextranucrase_production_by_local_Leuconostoc_mesenteroides)
  63. Characterization of the Different Dextranucrase Activities Excreted,  
<https://pmc.ncbi.nlm.nih.gov/articles/PMC106144/>
  64. Purging of natural salt-stock pickle fermentation to reduce bloater,  
[https://www.researchgate.net/publication/230061876\\_Purging\\_of\\_natural\\_salt-stock\\_pickle\\_fermentation\\_to\\_reduce\\_bloater\\_damage](https://www.researchgate.net/publication/230061876_Purging_of_natural_salt-stock_pickle_fermentation_to_reduce_bloater_damage)
  65. Factors influencing bloater damage in brined cucumbers during,  
[https://www.researchgate.net/publication/230061900\\_Factors\\_influencing\\_bloater\\_damage\\_in\\_brined\\_cucumbers\\_during\\_controlled\\_fermentation](https://www.researchgate.net/publication/230061900_Factors_influencing_bloater_damage_in_brined_cucumbers_during_controlled_fermentation)
  66. Process for reducing pickle bloating with dehydroacetic acid,  
<https://patents.google.com/patent/US3849579A/en>
  67. BLOATER FORMATION IN BRINED CUCUMBERS FERMENTED,  
<https://www.semanticscholar.org/paper/2c0af8d7d922febb40c5acee9ada3540ceb04e0f>
  68. Contribution of *Leuconostocaceae* to CO<sub>2</sub>-mediated bloater defect,  
<https://www.ars.usda.gov/ARSUserFiles/60702500/Fermented%20Veg%20Manuscripts/p434.pdf>
  69. Evaluation of malolactic-deficient strains of *Lactobacillus plantarum*,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p246.pdf>
  70. susceptibility of pickling cucumbers - USDA ARS,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p137.pdf>
  71. What causes a pickling cucumber to hollow out like this ... - Reddit,  
[https://www.reddit.com/r/fermentation/comments/x0d2qo/what\\_causes\\_a\\_pickling\\_cucumber\\_to\\_hollow\\_out/](https://www.reddit.com/r/fermentation/comments/x0d2qo/what_causes_a_pickling_cucumber_to_hollow_out/)
  72. bloaters during cucumber fermentation - USDA ARS,  
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p127.pdf>
  73. Prevention of Mold-Induced Softening in Air-Purged, Brined,

- <https://www.ars.usda.gov/ARSTUserFiles/60701000/Pickle%20Pubs/p151.pdf>
74. Purging CO<sub>2</sub> from cucumber brines to reduce bloater damage,  
[https://www.researchgate.net/publication/230062383\\_Purging\\_CO2\\_from\\_cucumber\\_brines\\_to\\_reduce\\_bloater\\_damage](https://www.researchgate.net/publication/230062383_Purging_CO2_from_cucumber_brines_to_reduce_bloater_damage)
  75. Publication : USDA ARS,  
<https://www.ars.usda.gov/research/publications/publication/?seqNo115=425891>
  76. Fermentation Safety Guide: Signs, Rules & When to Toss,  
<https://bloomcooking.com/learn/fermentation/fermentation-safety/>
  77. Is My Ferment Safe? A Decision Tree - SourChad,  
<https://sourchad.com/safety/is-my-ferment-safe>
  78. Clostridium botulinum & Botulism | Food Safety and Inspection Service,  
<https://www.fsis.usda.gov/food-safety/foodborne-illness-and-disease/illnesses-and-pathogens/botulism>
  79. Manufacture of Traditionally Fermented Vegetable Products,  
<https://www.foodprotection.org/members/fpt-archive-articles/2020-07-manufacture-of-traditionally-fermented-vegetable-products-best-practice-for-small-businesses/>
  80. Preservation of vegetables and herbs in oil - CSIRO,  
<https://www.csiro.au/en/research/health-medical/nutrition/Vegetable-preservation>
  81. The Importance of Food pH in Commercial Canning Operations,  
<https://extension.okstate.edu/fact-sheets/the-importance-of-food-ph-in-commercial-canning-operations>
  82. Growth Kinetics of Probiotic Lactobacillus Strains in the Alternative,  
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7760101/>
  83. Oak Barrel Care and Cleaning 101,  
<https://1000oaksbarrel.com/blog/oak-barrel-cleaning/>