

The Forage-Fed Heritage Bovine: An Integrated Mass-Balance Analysis of Milk Yield and Artisanal Dairy Production

The Dual-Purpose Paradigm in Pasture-Based Systems

The contemporary global dairy industry is overwhelmingly dominated by specialized high-input breeds, primarily the Holstein-Friesian, managed within Total Mixed Ration (TMR) confinement systems designed to maximize volumetric output. However, a profound structural and economic reset is occurring within the artisanal and organic dairy sectors. This shift is driven by the valuation of milk solids, genetic resilience, and the premiumization of grass-fed dairy products¹. Within this framework, dual-purpose heritage breeds—such as the Fleckvieh (German Simmental), Braunvieh (Brown Swiss), and Normande—offer a highly sophisticated biological engine for converting human-inedible perennial forages into high-value lipids and proteins².

These heritage breeds are characterized by immense metabolic plasticity. While a modern Holstein prioritizes lactogenesis to the detriment of body condition, often requiring massive starch and concentrate inputs to survive negative energy balance, dual-purpose breeds effectively partition energy to maintain muscle mass and reproductive health alongside lactation⁵. In an optimized, 100% pasture-based or forage-only system, the volumetric milk yield of these animals is naturally constrained by the physical limits of Dry Matter Intake (DMI) associated with bulky, high-fiber forages².

While a Fleckvieh cow may produce up to 11,000 kg of milk per lactation in a highly supplemented TMR system, the physiological realities of a strictly forage-fed diet yield significantly different outcomes⁵. Comprehensive agronomic tracking demonstrates that heritage cows fed wholly on grazed pasture and conserved winter forage yield between 3,250 kg and 7,000 kg of milk annually, depending on the intensity of rotational grazing and sward quality¹⁰. The utilization of intensive rotational grazing systems, incorporating deep-rooting multispecies swards like ribwort plantain and chicory, enhances protein bypass in the rumen and provides profound drought resilience⁵. For the purpose of establishing a definitive mass-balance and yield mapping model in this report, an optimized baseline of 6,000 kg of milk per lactation will be utilized. This figure represents the established physiological capability of a well-managed grazing Fleckvieh or Braunvieh cow operating within a seasonal, low-input grass-fed system⁹.

Reproductive Physiology and Herd Replenishment Dynamics

The successful realization of a consistent 6,000 kg annual milk yield is entirely predicated on immaculate reproductive management. The bovine female is a non-seasonal polyestrous

animal, maintaining continuous reproductive cyclicity throughout the year post-puberty, regulated by a complex hypothalamic-pituitary-ovarian axis⁵.

Ovarian Follicular Waves and Estrous Endocrine Shifts

The bovine estrous cycle averages 21 days in length and is characterized by distinct biological waves of follicular growth. Transrectal ultrasonography reveals that these cycles are composed of either two or three distinct follicular waves, initiated by transient elevations in follicle-stimulating hormone (FSH)⁵. During a phenomenon termed deviation, one follicle secures dominance via localized insulin-like growth factor signaling and enhanced aromatase activity, secreting massive quantities of estradiol to force the subordinate cohort into atresia⁵. Estrus is initiated by peak concentrations of estradiol, leading to a massive preovulatory surge of luteinizing hormone (LH), triggering ovulation 24 to 32 hours later⁵. Following ovulation, the corpus luteum forms and secretes progesterone, holding the uterus in a state of pregnancy preparedness. If maternal recognition of pregnancy does not occur, prostaglandin F2-alpha induces rapid luteolysis, initiating the next cycle⁵. Mastery of these ovarian dynamics allows producers to accurately time insemination protocols.

Venereal Pathogens and Low-Technology Artificial Insemination

Historically, pasture-based reproduction relied on natural service bulls, inherently posing severe biosecurity risks. The two most economically destructive venereal pathogens in cattle are the protozoan *Tritrichomonas foetus* and the bacterium *Campylobacter fetus* subsp. *venerealis*⁵. These obligate extracellular organisms persistently colonize the prepuce of mature bulls asymptotically, mechanically transferring to the female during coitus to induce endometritis, mid-term abortions, and plummeting pregnancy rates⁵.

The transition to artificial insemination (AI) eradicates these biosecurity threats while granting access to elite genetics capable of boosting lactation yields⁵. In low-resource, pasture-based settings lacking liquid nitrogen cryopreservation, ambient-temperature liquid semen preservation remains highly effective. Utilizing extenders such as the Illini Variable Temperature (IVT) buffer, which bubbles carbon dioxide through the solution to lower the pH to 6.3, exploits the glyoxylate mechanism to reversibly arrest sperm respiration⁵. Alternatively, Caprogen extenders utilize nitrogen gassing and the enzyme catalase to neutralize toxic hydrogen peroxide generated during prolonged storage⁵. These buffered solutions permit the storage of highly viable semen at room temperatures for up to six days, facilitating immaculate recto-vaginal insemination using sterilized, reusable glass pipettes to deposit semen precisely into the short uterine body⁵.

Obstetric Biomechanics and Uterine Involution

Parturition demands exacting obstetrical management to ensure maternal survival and subsequent lactation. Dystocia, caused frequently by fetopelvic disproportion, requires meticulous mechanical extraction. Obstetrical chains must be applied using a "double half-hitch" configuration to distribute tension across the calf's limbs, preventing severe metacarpal fractures⁵. Traction must never exceed the force generated by two to three adults (approximately 95.4 kg of mean peak force), as motorized extraction routinely exceeds 188.6 kg of force, precipitating fatal maternal nerve damage⁵. Aggressive compression of the obturator

nerve against the maternal pelvis induces calving paralysis, destroying the innervation of the adductor muscles and resulting in devastating hip dislocations⁵.

Following successful delivery, the massive 10 kg hypertrophied uterus undergoes involution, requiring 40 to 50 days for complete histological regeneration and the avascular necrosis of the maternal caruncles⁵. Concurrently, the immediate survival of the agammaglobulinemic neonate relies entirely on the ingestion of maternal colostrum within the first hours of life, exploiting a transient physiological window where enterocyte pinocytosis allows intact Immunoglobulin G macromolecules to traverse the intestinal barrier before total gut closure occurs at 24 hours postpartum⁵.

Metabolic Management and Trace Mineral Antagonisms

Ensuring a steady supply of milk from dual-purpose breeds requires exacting nutritional precision, particularly during the transition period. A mature 650 kg dual-purpose cow consumes between 3.0% and 3.5% of her body weight in dry matter daily, equating to approximately 19.5 to 22.75 kg of forage⁵.

The onset of high-volume lactogenesis places an unprecedented instantaneous drain on the cow's calcium reserves. Synthesizing colostrum demands nearly eight times the total normal calcium circulating in the plasma pool⁵. When parathyroid hormone and 1,25-dihydroxyvitamin D3 fail to mobilize skeletal reserves rapidly enough, total serum calcium plummets below 2.14 mmol/L, inducing subclinical hypocalcemia⁵. This intracellular calcium deficit severely impairs the phagocytic capabilities of polymorphonuclear leukocytes, rendering the cow highly susceptible to metritis and retained placentas⁵.

Furthermore, pastured herds face complex trace mineral antagonisms dictated by soil geochemistry. Excess dietary sulfur and molybdenum in lush pastures lead to the formation of thiomolybdates in the rumen, which bind irreversibly to copper, inducing severe secondary deficiencies⁵. Concurrently, high potassium influxes from spring grasses depolarize the ruminal epithelium, paralyzing active magnesium transport and precipitating hypomagnesemia (grass tetany)⁵. Meticulous agronomic management and targeted mineral supplementation are therefore mandatory to sustain the 6,000 kg lactation curve safely.

Compositional Biochemistry of Forage-Fed Milk

The true economic and functional value of milk derived from heritage breeds on pasture lies not in its volume, but in its dense compositional architecture and altered lipidomic profile.

Macronutrient Baselines and Genetic Polymorphisms

Milk from robust dual-purpose breeds exhibits superior concentrations of fat and protein. The Braunvieh breed, for instance, routinely delivers milk comprising 4.0% to 4.22% butterfat and 3.50% to 3.80% protein²². The Fleckvieh demonstrates a similar profile, averaging 4.13% to 4.50% fat and 3.50% protein in grass-fed systems¹¹. This analysis will utilize a standardized baseline composition of 4.20% butterfat and 3.50% true protein for the 6,000 kg annual yield model.

Beyond gross macronutrient percentages, the genetic polymorphisms of milk proteins in these breeds fundamentally alter their cheesemaking suitability. Bovine milk protein consists of approximately 80% casein (subdivided into α_{s1} , α_{s2} , β , and κ fractions) and 20% whey proteins (β -lactoglobulin and α -lactalbumin)²⁴. The κ -casein (CSN3) gene is of paramount importance, as κ -casein stabilizes the casein micelle in suspension, dictating rennet coagulation efficiency²⁶.

Heritage breeds possess an exceptionally high allelic frequency of the κ -casein B variant, with the BB genotype appearing in 63.8% to over 80% of specific populations, compared to less than 20% in standard Holstein herds²⁸. The BB genotype is associated with smaller, more numerous casein micelles, which drastically reduces rennet coagulation time, increases curd firmness, and improves the retention of both fat and protein in the cheese matrix²⁷. Milk containing the BB variant yields up to 13% more cheese per unit of volume than milk containing the AA variant²².

The Rumen Microbiome and Lipid Biohydrogenation

A 100% forage diet induces profound shifts in the fatty acid (FA) profile of the milk fat. Fresh pasture grass is rich in polyunsaturated fatty acids (PUFAs), particularly α -linolenic acid (an Omega-3 fatty acid)³⁴. When these lipids enter the bovine rumen, the microbial consortium engages in biohydrogenation to neutralize the toxicity of unsaturated double bonds, a critical survival mechanism for the microflora.

This process converts linolenic and linoleic acids into various intermediate saturated and monounsaturated configurations, most notably Vaccenic Acid (VA). Upon absorption into the mammary gland, the endogenous enzyme Δ^9 -desaturase converts a substantial portion of this Vaccenic Acid into Conjugated Linoleic Acid (CLA, specifically the *cis*-9, *trans*-11 isomer, or rumenic acid)⁸. Consequently, milk from cows on a 100% forage diet contains up to 147% more Omega-3 fatty acids and up to 500% more CLA than milk from TMR-fed cows consuming massive quantities of corn silage and grain³⁵.

This biochemical shift alters the Omega-6 to Omega-3 ratio from a highly inflammatory 5.7:1 (typical of grain-fed dairies) to a near 1:1 physiological balance³⁵. The daily grazing pattern further influences this profile; night-time grazing has been shown to yield a higher proportion of specific polyunsaturated fatty acids compared to daytime grazing, highlighting the sensitivity of the lipidomic output to environmental variables³⁸. Furthermore, the increased proportion of unsaturated fatty acids physically lowers the melting point of the milk fat, directly altering the rheology and spreadability of manufactured butter³⁹.

Mathematical Modeling of Butter Yield

Butter production is fundamentally a process of phase inversion, transitioning milk from an oil-in-water emulsion (cream) into a water-in-oil emulsion (butter) through rigorous mechanical

agitation⁴¹.

The Mechanics of Phase Inversion

When cream is mechanically churned at controlled temperatures (ideally 8°C to 12°C), the physical impact disrupts the protective Milk Fat Globule Membrane (MFGM)⁴¹. Liquid fat exudes from the fractured globules, acting as a cementing agent that binds the remaining intact solid fat globules together into macroscopic butter granules. As these granules coalesce and grow in density, the aqueous phase (buttermilk) separates completely and is drained off⁴². The granules are then mechanically worked to finely disperse residual moisture droplets throughout the continuous fat matrix, ensuring microbiological stability and textural smoothness.

Yield Calculation and Over-run Dynamics

Butter yield is dictated by the efficiency of fat recovery and the controlled incorporation of non-fat components, primarily moisture, milk solids-not-fat (curd), and salt. This incorporation is quantified in the dairy industry as "over-run," which represents the percentage increase in the weight of the finished butter compared to the absolute weight of the churned fat⁴¹.

The exact yield formula utilized by dairy technologists is defined mathematically as:

$$Y = \frac{F \times (100 + \%OR)}{100}$$

Where:

- Y = Yield of butter (in kg).
- F = Weight of fat inside the churn (in kg).
- $\%OR$ = Percentage of over-run⁴².

By strict legal and commercial standards worldwide, butter must contain a minimum of 80% milk fat by weight. Therefore, 80 kg of pure fat theoretically yields 100 kg of butter (the remaining 20% comprising up to 16% moisture, 1.5% to 2.0% salt, and 1.5% to 2.0% curd)²¹. This rigid constraint sets the absolute theoretical maximum over-run at exactly 25%⁴². However, accounting for unavoidable mechanical fat losses expelled in the buttermilk, practical factory over-run typically operates between 20% and 22.7%⁴¹.

Assuming an optimized commercial churning efficiency where 99.3% of the fat is recovered from the cream matrix¹⁸, the total theoretical butter yield from the heritage cow's entire annual fat production (252 kg) can be calculated to demonstrate maximal capacity:

- Total Milk Fat = 252.0 kg.
- Churned Fat successfully retained (F) = $252.0 \times 0.993 = 250.23$ kg.
- Applying a highly efficient, practical 20% over-run: $Y = \frac{250.23 \times (100 + 20)}{100} = 300.27$ kg of butter.

While dedicating the entire yield to butter is theoretically possible, artisanal creameries seek

multi-product diversification. The butter derived from this 100% forage diet will possess a distinctly lower solid fat content at room temperature, manifesting as superior spreadability, alongside a deep, brilliant yellow hue driven by elevated beta-carotene transferred directly from the pasture sward³⁹.

Thermo-Acid Coagulation: The Biochemistry of Mascarpone

Unlike traditional cheeses that rely entirely on the enzymatic cleavage of casein by chymosin, mascarpone is a high-moisture, extra-fat fresh cheese produced via the complex thermo-acid coagulation of heavy cream⁴³.

Isoelectric Precipitation and Whey Protein Denaturation

The physical chemistry of authentic mascarpone hinges on the simultaneous application of extreme thermal stress and organic acidification. A standardized cream mixture is heated to temperatures between 85°C and 90°C (185°F to 195°F)¹⁷. At these elevated temperatures, the

tertiary structures of the whey proteins (β -lactoglobulin and α -lactalbumin) completely unfold, exposing highly reactive sulfhydryl groups hidden within their native globular conformation⁴⁶.

Following this severe thermal activation, a dilute organic acid—typically tartaric or citric acid—is introduced to rapidly depress the pH of the matrix toward the isoelectric point of casein (pH 4.6)⁴⁵. The aggressive neutralization of the negative surface charges on the casein micelles abolishes their electrostatic repulsion. Concurrently, the unfolded, reactive whey proteins co-precipitate and form dense, complex aggregates with the collapsing casein micelles, trapping massive quantities of milk fat within the resulting coagulum⁴⁸. This extensive co-precipitation of whey proteins is entirely responsible for the dense, velvety, and easily spreadable rheology characteristic of mascarpone¹⁹. It is critical to note that cream subjected to Ultra-High Temperature (UHT) processing prior to this stage cannot be utilized effectively, as premature thermal denaturation destroys the requisite protein scaffolding necessary for proper curd formation upon acidification⁵¹.

Mascarpone Yield Dynamics

To achieve the optimal sensory profile, the initial cream matrix must be strictly standardized to a precise butterfat content of 20% to 25%¹⁹. In an artisanal setting, this is typically achieved by blending separated heavy whipping cream (36-40% fat) with fresh whole milk (3.25-4.20% fat)⁴⁵.

If an operation were to dedicate the cow's entire 252 kg of annual milk fat to mascarpone production, the mass balance requires heavy dilution with the aqueous phase of the milk to reach the 22.5% target. The thermo-acid coagulation process is highly efficient at retaining lipids, boasting a fat recovery coefficient of approximately 80%²⁰.

- Recovered Fat in Curd = $252 \times 0.80 = 201.6$ kg.
- Given that finished, drained mascarpone is classified legally as an extra-fat or

double-cream cheese containing roughly 50% fat by absolute weight⁴³, the final yield calculation is mathematically straightforward:

- Total Mascarpone Yield = $201.6 \text{ kg} / 0.50 = 403.2 \text{ kg}$.

Enzymatic Coagulation and Aged Cheese Yield Prediction

The conversion of fluid milk into hard, aged cheeses (e.g., Cheddar or Parmigiano-Reggiano styles) relies on the precise enzymatic cleavage of κ -casein by rennet, followed by extensive syneresis and whey expulsion⁵³. Predicting the exact volumetric yield of this biological process is an economic imperative for the dairy industry, universally calculated using the Van Slyke and Price equation⁵⁵.

The Modified Van Slyke Equation

Formulated over a century ago, the Van Slyke equation operates as an exact mass-balance algorithm. It quantifies the theoretical transfer of fat and casein from the liquid phase into the solid curd matrix, adjusted for unavoidable process losses and target moisture parameters¹⁶. The contemporary modified exact Van Slyke formula is expressed as:

$$Y = \frac{((F \times RF) + (C \times RC)) \times 1.09}{1 - M}$$

Variable	Definition	Typical Values & Impact
Y	Cheese yield	Expressed as kg of cheese per 100 kg of milk.
F	Milk fat percentage	Primary driver of yield and mouthfeel.
RF	Fat recovery factor	Proportion of fat trapped in the matrix (0.92 to 0.95) ⁵⁵ .
C	Milk casein percentage	The structural protein scaffold. Derived as True Protein $\times 0.78$ ⁵⁶ .
RC	Casein recovery factor	Proportion of casein

		retained (0.95 to 0.98) ⁵⁵ .
1.09	Whey solids retention factor	Accounts for soluble salts, minerals, and lactose remaining in the curd moisture ⁵⁵ .
M	Target cheese moisture	Expressed as a decimal (e.g., 0.37 for Cheddar, 0.30 for Parmesan) ⁵⁵ .

Note: Older iterations of the Van Slyke formula utilized a rigid, constant subtraction of 0.1 for casein fines lost in the whey (e.g., $(C - 0.1)$)⁵³. Contemporary predictive models favor the fractional recovery coefficient (RC), as it dynamically scales with the exceptional casein retention facilitated by the κ -casein BB variant highly prevalent in heritage breeds³¹.

Standardization and the Casein-to-Fat Ratio

Cheese yield and structural integrity are fundamentally governed by the Casein-to-Fat (C/F) ratio of the initial cheese milk in the vat⁵³. If the fat content is too high relative to the casein scaffolding, the protein network lacks the tensile strength to encapsulate the massive fat globules. This results in severe lipid leakage into the whey during pressing, yielding a greasy, structurally compromised cheese that will rapidly spoil during aging⁶⁰.

The raw milk of the heritage cow under analysis contains 4.20% fat and 3.50% protein. Given the 0.78 casein conversion factor, the true casein content is 2.73%. This yields a raw C/F ratio of $0.65 (2.73/4.20)$.

For a full-fat Cheddar-style cheese, the optimal C/F ratio is tightly defined between 0.64 and 0.70⁵³. Therefore, the raw heritage milk is naturally perfect for Cheddar applications and requires no centrifugal standardization. However, hard alpine grating cheeses, such as Parmigiano-Reggiano, require immense structural rigidity to survive extreme, long-term aging without weeping fat or collapsing under their own weight. The strict parameters for these styles dictate that the vat milk must be partially skimmed to achieve a significantly higher C/F ratio, typically targeting between 0.80 and 1.10⁵³.

The Paradox of Aging: Moisture Loss and Proteolysis

Aged cheeses undergo a continuous, slow loss of mass during maturation in the climate-controlled ripening facility. For a hard Parmigiano-style cheese aged over 24 to 36 months, the internal moisture evaporates continuously through the dense rind, dropping from an initial 30% down to approximately 25% by month 50⁶⁵. This profound desiccation routinely causes an absolute weight reduction of 10% to 15% from the fresh yield out of the press. However, this physical shrinkage drives intense biochemical refinement. As moisture decreases, microbial enzymes and residual endogenous plasmin orchestrate profound

proteolysis—the systematic breakdown of the massive para-casein network into highly volatile smaller peptides and free amino acids⁵⁴. As the water matrix evaporates, the concentration of the specific amino acid tyrosine exceeds its solubility threshold, causing it to precipitate directly out of the paste into macroscopic, crunchy white crystals⁶⁶. These tyrosine crystals, alongside calcium lactate, serve as the ultimate biochemical markers of advanced age, signifying the transition from basic structural decay into profound sensory delicacy and umami⁵⁴.

Next-Generation Yield Optimization: Whey Microparticulation

When raw milk is converted into cheese via enzymatic coagulation, the casein matrix traps the fat, but over 95% of the highly nutritious whey proteins (β -lactoglobulin and α -lactalbumin) are flushed out of the vat as liquid whey⁶⁷. Historically, this vast volume of acid or sweet whey represented a catastrophic environmental liability due to its massive Biochemical Oxygen Demand (BOD) and Chemical Oxygen Demand (COD). Discharging raw whey directly into waterways severely depletes aquatic oxygen levels, causing devastating ecological toxicity⁶⁹. Contemporary dairy engineering circumvents this environmental hazard while drastically improving plant economics through Whey Protein Upcycling via Microparticulation. The expelled liquid whey undergoes ultrafiltration to concentrate the soluble proteins, followed by a precision thermal-mechanical treatment. The optimal parameters dictate heating the whey protein concentrate to 93°C for exactly 5.4 minutes under continuous, aggressive high shear⁶⁸. This process intentionally denatures the whey proteins, forcing them to aggregate into microscopic, uniform spherical structures measuring between 1 and 10 micrometers in diameter⁷².

Remarkably, these microparticulated whey proteins (MWP) possess the exact morphological dimensions of native milk fat globules⁷². When reintegrated in liquid form into subsequent batches of cheese milk, these inert protein spheres become physically entrapped within the three-dimensional para-casein network during the coagulation phase⁷². MWP exhibits a massive, sponge-like water-holding capacity. Its inclusion acts as an invisible rheological fat replacer, actively boosting the total solids recovery, driving up cheese yield by several percentage points, and restoring the creamy palatability and structural elasticity often lost in lower-fat, standardized hard cheeses⁷³. By intercepting the whey stream and converting a potent pollutant into highly functional structural yield, microparticulation secures the absolute maximization of the herd's biological output⁷².

The Integrated Artisanal Mass-Balance Model

To demonstrate the full economic and physical potential of the forage-fed heritage cow, we map the 6,000 kg annual yield into a diversified, integrated mass-balance model. Rather than dedicating the entire volume to a single bulk commodity, an optimized artisanal creamery splits the milk mathematically to simultaneously capture the value of aged alpine cheeses, superior grass-fed butter, and rich mascarpone.

Initial Raw Material Input (Annual Yield of One Cow):

- Total Milk Volume: 6,000 kg
- Milk Fat: 4.20% (252.0 kg total fat)
- Milk True Protein: 3.50% (210.0 kg total protein)
- Calculated Casein: 2.73% (163.8 kg total casein)⁵⁷
- Initial C/F Ratio: 0.65

Step 1: Centrifugal Skimming for Standardization

To achieve an ideal Casein-to-Fat ratio for a structurally sound aged alpine cheese (targeting a C/F ratio of 0.81), 1,200 kg of the raw milk is diverted through a centrifugal separator⁵³.

- Separation of 1,200 kg of whole milk yields roughly 126 kg of heavy cream (at 40% fat) and 1,074 kg of skim milk (at a negligible 0.05% fat)⁶⁰.
- Fat partitioned into the heavy cream: $126 \times 0.40 = 50.4$ kg of fat.

Step 2: Formulating the Cheese Vat and Van Slyke Yield

The 1,074 kg of skim milk is blended back into the remaining 4,800 kg of untouched whole raw milk.

- Total Standardized Cheese Milk in Vat: 5,874 kg.
- Total Fat remaining in Vat: $252.0 \text{ kg (initial)} - 50.4 \text{ kg (removed)} = 201.6$ kg of fat.
- Standardized Fat Percentage: $201.6/5874 = 3.43\%$ fat.
- Total Casein in Vat: 163.8 kg (casein remains entirely in the skim phase and is unaffected by centrifugal separation) $\approx 2.79\%$ casein.
- New C/F Ratio: $2.79/3.43 = 0.81$. (An optimal, legal parameter for firm, long-aged alpine cheeses)⁵³.

Yield Calculation for the Alpine Cheese (targeting 32% moisture, using standard recovery factors of $RF = 0.94$, $RC = 0.97$):

$$Y = \frac{((3.43 \times 0.94) + (2.79 \times 0.97)) \times 1.09}{1 - 0.32}$$

$$Y = \frac{(3.2242 + 2.7063) \times 1.09}{0.68} = \frac{5.9305 \times 1.09}{0.68} = 9.50\%$$

- **Total Fresh Cheese Yield:** $5,874 \text{ kg} \times 9.50\% = 558.0 \text{ kg of fresh cheese wheels.}$

- **Aged Cheese Yield (Post-Ripening):** Assuming a standard 12% moisture evaporation loss over a 24-month affinage period, the final harvestable mass is reduced to **491.0 kg of aged cheese**⁶⁵.

Step 3: Utilizing the Extracted Cream for Butter and Mascarpone

The 126 kg of heavy cream (containing 50.4 kg of pure fat) extracted during the standardization phase is now split to manufacture the high-value lipid products.

Product Stream	Cream Allocation	Fat Allocation	Yield Calculation	Final Product Weight
Grass-Fed Butter	76.0 kg (at 40% fat)	30.4 kg fat	$30.4 \text{ kg} \times 0.99$ [cite: 42]	36.2 kg of Butter
Mascarpone	50.0 kg (at 40% fat)	20.0 kg fat	Dilute to 22.5% fat using 38.9 kg whole milk (adds 1.63 kg fat). Total mix: 88.9 kg containing 21.63 kg fat. Yield: (21.63×0.80) [cite: 43, 52]	34.6 kg of Mascarpone

Economic Ramifications and Conclusion

The utilization of a heritage dual-purpose bovine grazing exclusively on perennial forage represents a masterclass in biological efficiency, environmental stewardship, and biochemical complexity. While abandoning energy-dense grain rations intentionally restricts the absolute volumetric yield of the animal to approximately 6,000 kg annually, it fundamentally elevates the lipidomic and genetic value of the milk. Federal Milk Marketing Orders (FMMOs) and modern multiple-component pricing systems globally are increasingly penalizing mere volume while heavily rewarding dense concentrations of protein and fat, perfectly aligning with the heritage breed's physiological output⁷⁶.

The overwhelming prevalence of the κ -casein BB allele in these cattle supercharges the efficiency of the Van Slyke mass-transfer equation, aggressively pulling casein and fat out of the aqueous phase and locking it into the cheese matrix, routinely providing a 10% to 13% yield dividend over Holstein milk²². Concurrently, the ruminal biohydrogenation of lush pasture grasses imbues the milk fat with extraordinary concentrations of conjugated linoleic acid and

Omega-3 fatty acids, shifting the Omega 3:6 ratio to a near 1:1 balance and fundamentally altering the melting rheology and vibrant hue of the resulting butter³⁵.

Through the precise, calculated application of centrifugal standardization, enzymatic and thermo-acid coagulation, and advanced whey microparticulation to mitigate environmental COD/BOD loads, the modern artisanal facility can mathematically trace every gram of human-inedible forage into highly refined, premium dairy solids⁷⁰. As definitively demonstrated in the integrated mass-balance model, a single lactation of 6,000 kg can be elegantly and precisely partitioned to yield nearly 500 kilograms of crystallized, aged alpine cheese, alongside high-value secondary streams of 36.2 kg of golden grass-fed butter and 34.6 kg of rich mascarpone. This paradigm proves that operating within the natural metabolic constraints of the grazing bovine secures both profound sensory quality and robust operational viability.

Works cited

1. Production Costs of Grass-Fed Organic Milk in the Northeastern, <https://www.mdpi.com/2071-1050/17/24/11324>
2. Increasing Grazing in Dairy Cow Milk Production Systems in Europe, <https://www.mdpi.com/2071-1050/12/6/2443>
3. (PDF) Comparative study on production, reproduction and functional, https://www.researchgate.net/publication/315770920_Comparative_study_on_production_reproduction_and_functional_traits_between_Fleckvieh_and_Braunvieh_cattle
4. Comparative Study on Milk Production, Milk Components ... - PubMed, <https://pubmed.ncbi.nlm.nih.gov/40387471/>
5. 44 Bovine Reproduction And Herd Management.pdf
6. Fleckvieh Changes Environmentally friendly cattle breeding Eight, https://www.fleckvieh.at/wp-content/uploads/2025/03/FC_Magazin-1_0121.pdf
7. Pasture-based dairy farm systems increasing milk production, https://www.researchgate.net/publication/229915621_Pasture-based_dairy_farm_systems_increasing_milk_production_through_stocking_rate_or_milk_yield_per_cow_pasture_and_animal_responses
8. Changes in milk production and milk fatty acid composition of cows, <https://www.tandfonline.com/doi/full/10.1080/1828051X.2016.1141330>
9. German Fleckvieh – the universal breed! - Service - GGI-SPERMEX, <https://www.ggi-spermex.de/en/service/german-fleckvieh-%E2%80%93-the-universal-breed!-1729.html>
10. Can all-forage dairy diets be practical and profitable?, <https://www.fwi.co.uk/livestock/dairy/can-all-forage-dairy-diets-be-practical-and-profitable>
11. Low input, full pasture and seasonal calving with great Fleckvieh cows, <https://www.ggi-spermex.de/en/service/insight:-low-input-full-pasture-and-seasonal-calving-with-great-fleckvieh-cows-1850.html>
12. (PDF) Economic Analysis of Forage Mixture Productivity in Pastures, https://www.researchgate.net/publication/43263803_Economic_Analysis_of_Forage_Mixture_Productivity_in_Pastures

- [ge_Mixture_Productivity_in_Pastures_Grazed_by_Dairy_Cattle](#)
13. The “Grass-Fed” Milk Story: Understanding the Impact of Pasture, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6723057/>
 14. German Simmental Fleckvieh – The Best Dual-Purpose Breed in the, <https://alpineagriculture.ch/en/german-simmental-fleckvieh-the-best-dual-purpose-breed-in-the-world/>
 15. About Fleckvieh - GGI-SPERMEX, <https://www.ggi-spermex.de/en/fleckvieh/about-fleckvieh-92.html>
 16. How to Calculate Cheese Yield from Milk | DairyCraftPro, <https://dairycraftpro.com/how-to-calculate-cheese-yield-from-milk-the-complete-guide-for-cheesemakers/>
 17. Homemade Mascarpone Cream Cheese - Kvalifood, <https://kvalifood.com/posts/homemade-mascarpone-cream-cheese/>
 18. Butter Manufacture – Dairy Science and Technology eBook, <https://books.lib.uoguelph.ca/dairyscienceandtechnologyebook/chapter/butter-manufacture/>
 19. Mascarpone Cheese Making Recipe, <https://cheesemaking.com/products/mascarpone-cheese-making-recipe>
 20. What it takes to make a perfect batch of Mascarpone Cheese, <https://scispace.com/pdf/mastering-mascarpone-what-it-takes-to-make-a-perfect-batch-2ju104fnon.pdf>
 21. Over Run - Dairy Technology, <http://dairy-technology.blogspot.com/2014/01/over-run.html>
 22. European Brown Swiss Federation www.brown-swiss.org, <http://www.germanlivestock.de/services/files/BS-MORE-THAN-MILK-Press-release.pdf>
 23. BRS - Fleckvieh - Bundesverband Rind und Schwein, <https://www.rind-schwein.de/brs-cattle/fleckvieh-en.html>
 24. Heritability and correlations of milk traits in the view of kappa-casein, <https://hrcak.srce.hr/file/150939>
 25. (PDF) Polymorphisms in CSN3, CSN2 and LGB Genes and Their, https://www.researchgate.net/publication/331426473_Polymorphisms_in_CSN3_CSN2_and_LGB_Genes_and_Their_Relation_to_Milk_Production_in_Dairy_Cattle_in_the_Czech_Republic
 26. Exploring the Genotype at CSN3 Gene, Milk Composition ... - MDPI, <https://www.mdpi.com/2076-2615/10/11/1995>
 27. (PDF) Kappa casein (CSN3) gene polymorphism and its effect on, https://www.researchgate.net/publication/356185124_Kappa_casein_CSN3_gene_polymorphism_and_its_effect_on_cumulative_milk_yields_of_Holstein_Friesian_dairy_cattle
 28. European Brown Swiss Federation, <https://homepage.braunvieh.ch/wp-content/uploads/2021/08/EBSF-BS-school-leaftlet-A4-E.pdf>
 29. Milk Caseins: Updated Genotype Frequencies by Breed - Lactanet, <https://lactanet.ca/en/milk-caseins-updated-genotype-frequencies-by-breed/>

30. Breeding for Kappa Casein to Increase Cheese Yield | The Bullvine,
<https://www.thebullvine.com/genetics/breeding-for-kappa-casein-to-increase-cheese-yield/>
31. Estimating the Effect of the Kappa Casein Genotype on Milk ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10778097/>
32. Comparative analysis of κ -casein expressions and dairy herd ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11928332/>
33. Genetic variants of milk proteins and their effects on the yield and quality of cheese? tp=eyJjb250ZXh0ljp7lnBhZ2UiOiJzY2llbnRpZmljQ29udHJpYnV0aW9ucyIsbnByZXZpb3VzUGFnZSI6bnVsbCwic3ViUGFnZSI6bnVsbH19
https://www.researchgate.net/publication/240511667_Genetic_variants_of_milk_proteins_and_their_effects_on_the_yield_and_quality_of_cheese?tp=eyJjb250ZXh0ljp7lnBhZ2UiOiJzY2llbnRpZmljQ29udHJpYnV0aW9ucyIsbnByZXZpb3VzUGFnZSI6bnVsbCwic3ViUGFnZSI6bnVsbH19
34. Composition and Profile of Fatty Acids in Milk on Quality and Health,
<https://ejournal.cwcu.ac.id/index.php/IJSTeMLaH/article/download/111/161/864>
35. Milk from Pastured Cows is Better for You - Cornucopia Institute,
<https://www.cornucopia.org/2018/03/milk-pastured-cows-better/>
36. Grain and CLA | Keeping A Family Cow,
<https://familycow.proboards.com/thread/63436/grain-cla>
37. Organic Production Enhances Milk Nutritional Quality by Shifting,
<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0082429>
38. Effect of total mixed ration composition and daily grazing pattern on, Effect of total mixed ration composition and daily grazing pattern on milk production composition and fatty acids profile of dairy cows
https://www.researchgate.net/publication/266188499_Effect_of_total_mixed_ratio_n_composition_and_daily_grazing_pattern_on_milk_production_composition_and_fatty_acids_profile_of_dairy_cows
39. The linear relationship between the proportion of fresh grass in the,
<https://pubmed.ncbi.nlm.nih.gov/16702259/>
40. The Influence of Different Butter Type, Their Fatty Acid Composition,
<https://www.mdpi.com/2304-8158/13/23/3803>
41. Understanding Over-Run in Butter Production - Agriculture Institute,
<https://agriculture.institute/dairy-products-i/understanding-over-run-butter-production/>
42. How to Calculate Butter Yield and Maximize Production,
<https://agriculture.institute/dairy-products-i/calculate-butter-yield-maximize-production/>
43. Mascarpone Cheese Production - Узловский молочный комбинат,
<https://uzmolkom.com/en/blog/mascarpone-cheese-production/>
44. Mascarpone - Description - Tiiips,
<https://www.tiiips.com/m/tiiips/home?action=listReviews&reviewID=38935&oldID=38414>
45. Mascarpone Cheese Making Recipe Instructions - roulette-one,
<https://roulette-one.myshopify.com/pages/mascarpone-cheese-making-recipe-instructions>
46. Heat-Induced Interaction of Milk Proteins: Impact on Yoghurt Structure,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8483934/>
47. An Integrated Approach in Assessing the Food-Related Properties of,

- <https://www.mdpi.com/2304-8158/14/19/3421>
48. Mascarpone, what is it? - GEA,
<https://www.gea.com/en/campaigns/fresh-cheese-uncovered/mascarpone/>
 49. Mascarpone Cheese - Catalina Bakes,
<http://catalinabakes.blogspot.com/2011/08/mascarpone-cheese.html>
 50. Masterarbeit Fettreduzierung bei Mascarpone,
https://digibib.hs-nb.de/file/dbhsnb_thesis_0000002499/dbhsnb_derivate_0000003118/Masterarbeit-Cao-2020.pdf
 51. Mascarpone Cheese - Probably Worth Sharing,
<https://probablyworthsharing.com/recipes/mascarpone-cheese>
 52. Mascarpone Info Cheese Making Recipe,
<https://cheesemaking.com/pages/mascarpone-cheese-making-recipe-info>
 53. Factors that Affect Cheese Yield,
https://adpi.org/wp-content/uploads/2022/08/COE-Article_Factors-That-Affect-Cheese-Yield_2022.pdf
 54. How Cheese Gets Sharper And Crumblier With Age Naturally,
<https://www.youtube.com/watch?v=oucu-rlpFI8>
 55. Cheese Yield Calculator (Van Slyke) - Watson Dairy Consulting,
<https://dairyconsultant.co.uk/cheese-yield-van-slyke.php>
 56. Free Cheese Yield Calculator (Van Slyke) - Vesper Tool,
<https://vespertool.com/calculators/dairy/cheese-yield-estimation/>
 57. Standardization of Milk for Cheese Making,
<https://books.lib.uoguelph.ca/cheesemakingtechnologyebook/chapter/3-3-standardization-of-milk-for-cheese-making/>
 58. Yield Prediction – Cheese Making Technology e-Book,
<https://books.lib.uoguelph.ca/cheesemakingtechnologyebook/chapter/yield-prediction/>
 59. Calculating Component Values from the Modified Van Slyke Cheese,
<https://www.ams.usda.gov/sites/default/files/media/DYCompValMBrownIIIVHearing2000.pdf>
 60. How to Standardize Milk Fat for Cheese | DairyCraftPro,
<https://dairycraftpro.com/how-to-standardize-milk-fat-for-cheese-a-complete-guide-for-cheesemakers/>
 61. Effect of the casein to fat ratio on the functional properties of Prato,
<https://rsdjournal.org/rsd/article/download/27812/24354/324269>
 62. Quantification of Cheese Yield Reduction in Manufacturing ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7074290/>
 63. FOR PARMIGIANO REGGIANO CHEESE,
https://backend.parmigianoreggiano.com/uploads/0325_disciplinare_docx_EN_2eda3456af.pdf
 64. PRODUCT SPECIFICATION OF THE PROTECTED DESIGNATION,
https://backend.parmigianoreggiano.com/uploads/disciplinare_PR_EN_clean_02072025_55f8b10686.pdf
 65. Impact of Extending Hard-Cheese Ripening - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7143483/>

66. Parmesan Tasting: Ages Compared (12–48+ Months),
<https://www.parmesandairytour.com/parmesan-tasting-ages-compared/>
67. Understanding the basics: Techniques to attain a high cheese yield,
<https://www.dairyfoods.com/articles/92100-understanding-the-basics-techniques-to-attain-a-high-cheese-yield>
68. Effect of microparticulate from whey protein concentrates on the,
http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-01352023000200005
69. Reuse of waste acid whey for extraction of high added value,
<https://webgate.ec.europa.eu/life/publicWebsite/project/LIFE16-ENV-SI-000335/reuse-of-waste-acid-whey-for-extraction-of-high-added-value-bioactive-proteins>
70. Whey Utilization: Sustainable Uses and Environmental Approach,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8284110/>
71. Paving the whey - transforming dairy waste - Teagasc,
<https://teagasc.ie/news--events/daily/paving-the-whey-transforming-dairy-waste/>
72. Whey Protein Upcycling & Microparticulation | Perinox,
<https://perinox.com/en/whey-protein-upcycling-microparticulation/>
73. Effect of microparticulated whey protein concentration and protein-to-fat ratio on Caciotta cheese yield and composition
https://www.researchgate.net/publication/273789867_Effect_of_microparticulated_whey_protein_concentration_and_protein-to-fat_ratio_on_Caciotta_cheese_yield_and_composition
74. (PDF) Low-fat Cheddar cheese made using microparticulated whey,
https://www.researchgate.net/publication/316311973_Low-fat_Ceddar_cheese_made_using_microparticulated_whey_proteins_Effect_on_yield_and_cheese_quality
75. Turning Whey Into Value with microparticulation - SPX Flow,
<https://www.spxflow.com/blog/turning-whey-into-value-with-microparticulation/>
76. Economic Analysis Class III and IV Pricing Formulas,
<https://www.ams.usda.gov/sites/default/files/media/Hearings%20Class%20III%20and%20IV%20Economic%20Analysis.pdf>
77. Changes to Milk Pricing Formulas Under Federal Milk Marketing,
<https://www.choicesmagazine.org/choices-magazine/theme-articles/dairy-theme/changes-to-milk-pricing-formulas-under-federal-milk-marketing-orders-will-affect-farm-milk-prices>
78. End of Project Report A New Direction for the Payment of Milk - T-Stor,
<https://t-stor.teagasc.ie/bitstreams/16952aa1-af50-427a-a2dd-e131c21b897a/download>