

Sylvatic Bio-Resource Management and Haute Chasse Cuisine in Post-Industrial Brandenburg: Wildlife Ecology, Non-Ballistic Harvesting, Post-Mortem Biochemistry, and Gastronomic Valorisation

Regional Agro-Ecological Baseline and Ungulate Population Dynamics

The Pleistocene glacial retreats formed the physical landscape of the North German Plain into a distinct succession of terminal moraines, outwash sandur plains, and broad urstromtal valleys. In the federal state of Brandenburg, which encircles the Berlin metropolitan basin, the pedological profile is defined by nutrient-deprived, highly permeable Arenosols, colloquially known as glacial blow sands and outwash sands, which register modest agricultural soil values (*Bodenwertzahlen*) between 25 and 40¹. Operating within USDA plant hardiness zone 7b, the territory is subject to a semi-continental climate characterized by an average annual

precipitation of 500 to 600 mm, an average annual temperature of roughly 9.5°C, and a restricted frost-free vegetative window of approximately 137 days¹.

While these sandy, acidic soils severely limit high-yield arable agriculture without industrial synthetic inputs, the regional forest canopy encompasses approximately 1.1 million hectares. Historically planted with Scots pine (*Pinus sylvestris*) monocultures, these woodlands are increasingly diversified with pedunculate oak (*Quercus robur*), sessile oak (*Quercus petraea*), and European beech (*Fagus sylvatica*). This extensive sylvatic environment provides exceptional forage and cover for wild ungulates, allowing Brandenburg to maintain some of the highest recorded biomass densities of wild boar (*Sus scrofa*) and roe deer (*Capreolus capreolus*) in Central Europe².

The wild boar thrives across the sandy outwash landscapes of Brandenburg due to recurrent mast years (*Mastjahre*) produced by mature oak and beech stands, mild winter conditions that reduce piglet mortality, and agricultural edge ecotones. Roe deer occupy the brushy forest fringes and regenerating understories as concentrate selectors (*Konzentratselektierer*), browsing selectively on dicotyledonous herbs, terminal buds, and agricultural greens. Red deer (*Cervus elaphus*) and fallow deer (*Dama dama*) maintain robust populations across extensive, contiguous sylvatic tracts such as the Schorfheide, the Fläming, and the Kyritz-Ruppiner Heide².

In contrast, the European brown hare (*Lepus europaeus*) inhabits the drier agricultural fields, pastures, and open clearings, though its populations remain constrained by historical mechanization and ground predation². Waterfowl, predominantly the mallard (*Anas platyrhynchos*) and greylag goose (*Anser anser*), congregate in massive seasonal flights along the Havel, Spree, and Oder river networks and their associated post-glacial wetland complexes³.

The trophic structure of Brandenburg includes the established presence of an apex carnivore: the grey wolf (*Canis lupus*). Recent systematic surveys confirm between 54 and 58 reproductive packs (*Rudel*) and 6 to 8 resident scent-marking pairs within state borders, representing a dynamic wild population of approximately 600 to 700 wolves⁹.

Dietary biomass analyses of wolf scats across Brandenburg establish that wild ungulates constitute over 95% of their prey intake, partitioned primarily into roe deer (40–50%), red deer (20–25%), and wild boar (15–20%), alongside localized predation on fallow deer¹². While this carnivore pressure curbs runaway ungulate recruitment and alters herd grazing behavior, the net annual biomass production across Brandenburg remains capable of supporting human nutritional extraction, provided that harvest rates respect demographic replacement boundaries².

Species	Baseline Spring Density (N/km2)	Annual Reproductive Rate (λ -1)	Brandenburg Annual Cull (Jagdstrecke 2024/25)	Trophic Role and Diet Classification
Wild Boar (<i>Sus scrofa</i>)	4.0 – 12.0 (locally >15.0)	150% – 250% (of spring adult stock)	60,208 ⁴	Opportunistic omnivore; roots, tubers, acorns, carrion, crops ¹
Roe Deer (<i>Capreolus capreolus</i>)	6.0 – 14.0	30% – 45% (80–100% of adult females) ¹⁵	45,351 ⁴	Concentrate selector; high-tannin browse, herbs, buds, soft mast
Fallow Deer (<i>Dama dama</i>)	2.0 – 8.0 (patchy distribution) ¹⁶	25% – 35% (75% of adult females) ¹⁵	7,440 ⁴	Intermediate feeder; grasses, heather, tree

				bark, mast
Red Deer (<i>Cervus elaphus</i>)	1.0 – 3.5 (large contiguous forests) ²	20% – 30% (75% of adult females) ¹⁵	6,205 ⁴	Intermediate/roughage grazer; grasses, sedges, shoot tips, bark
European Hare (<i>Lepus europaeus</i>)	2.0 – 6.0 (open agricultural land)	100% – 200% (high juvenile mortality)	1,183 ⁴	Herbivorous specialist; wild graminoids, herbaceous perennials
Waterfowl (<i>Anas/Anser</i> spp.)	Seasonal migrants / riparian	150% – 300% (variable hatch rate)	5,590 (3,115 ducks; 2,475 geese) ⁸	Graminoid grazers, aquatic macrophyte and seed foragers

Sustainable Harvest and Land-Budget Model for a 300-Person Agrarian Community

In an agrarian economy lacking mechanized equipment, synthetic petrochemical fertilizers, and imported goods, the harvest of wild ungulates provides dense animal protein, essential lipids, and bioavailable micronutrients without diverting arable land from subsistence crop production. Wild ruminants and suids convert indigestible sylvatic cellulose, browse, mast, and aquatic vegetation into complete proteins and fats.

For a community expanding from an initial domestic nucleus to a structured village of 300 individuals, setting wild game consumption at a target of 25 to 35 kg of dressed, boneless culinary meat per capita annually provides 68.5 to 95.8 grams of animal muscle daily per person. This intake provides 15 to 21 grams of biological protein per person each day, fulfilling approximately 30% to 40% of the physiological protein baseline, with the remainder supplied by pulses, cultivated grains, and domestic livestock.

The aggregate annual community demand is calculated as:

$$\text{Total Boneless Meat Demand} = 300 \text{ persons} \times 30.0 \text{ kg/yr} = 9,000.0 \text{ kg/yr}$$

To convert this net boneless target into harvest quotas, the physiological yields of wild carcasses must be evaluated across species. Wild cervids yield roughly 50% of their live weight as cold dressed hanging carcass, and approximately 68% of that hanging carcass is recoverable as deboned culinary muscle¹⁷. This establishes an overall boneless yield factor of 34% relative to live weight¹⁷.

Wild boar yields roughly 60% of live weight as dressed carcass (head-on, skin-on)¹⁹, and approximately 65% of that carcass mass as culinary cuts, processing trim, and solid fatback¹³, resulting in a net recovery factor of 39% relative to live weight¹³.

Target Harvest Contribution	Percent of Meat Demand	Net Boneless Meat (kg)	Average Live Weight per Specimen (kg)	Culinary Yield per Specimen (kg)	Total Head Required Annually
Wild Boar (<i>Sus scrofa</i>)	45%	4,050	60.0 (mixed cohorts)	23.4	173 animals
Roe Deer (<i>Capreolus capreolus</i>)	25%	2,250	18.0 (adult/subadult)	6.1	369 animals
Fallow Deer (<i>Dama dama</i>)	15%	1,350	55.0 (calves/adults)	18.7	72 animals
Red Deer (<i>Cervus elaphus</i>)	10%	900	110.0 (calves/hinds)	37.4	24 animals
Hare & Waterfowl	5%	450	3.5 (hares) / 1.2 (ducks)	1.8 / 0.6	~400 head combined
Total	100%	9,000 kg	—	—	1,038 head

To guarantee ecological equilibrium, total mortality from hunting extraction (H), natural predation (P , including wolves), and environmental attrition (M) must not exceed the demographic recruitment rate (R) of the target populations:

$$H + P + M \leq R = N_0 \cdot (\lambda - 1)$$

Given the landscape composition of Brandenburg, featuring approximately 40% mixed pine-hardwood canopy, 50% arable fields and fallow clearings, and 10% lakes and peat bogs, sustainable annual harvest rates per 100 hectares (1 km^2) under non-mechanized conditions are established as:

- Wild boar: 4.5 head harvested per km^2
- Roe deer: 8.0 head harvested per km^2
- Fallow deer: 1.5 head harvested per km^2
- Red deer: 0.5 head harvested per km^2

Dividing the required cull by the spatial harvest rates establishes the territory required to support the village:

$$A_{\text{boar}} = \frac{173}{4.5/\text{km}^2} = 38.4 \text{ km}^2 = 3,840 \text{ ha}$$

$$A_{\text{roe}} = \frac{369}{8.0/\text{km}^2} = 46.1 \text{ km}^2 = 4,610 \text{ ha}$$

$$A_{\text{red}} = \frac{24}{0.5/\text{km}^2} = 48.0 \text{ km}^2 = 4,800 \text{ ha}$$

A contiguous communal hunting territory (*Jagdallmende*) encompassing 3,500 to 4,500 hectares ($35 - 45 \text{ km}^2$) provides the necessary resource base. This represents an operational radius of roughly 3.3 to 3.8 km radiating outward from a central settlement, a

distance accessible to human foot travel and draft-animal transport.

To maintain demographic resilience, harvesting must strictly track traditional German

age-class quotas (*Altersklassen*)¹⁵. For wild boar, at least 80% of the annual take must consist of juveniles under one year of age (*Frischlinge*, 15–35 kg) and yearlings (*Überläufer*, 35–55 kg)¹⁵.

Adult reproductive females (*Bachen*, ≥ 2 years) are fully protected; removing lead sows disrupts sounder structure, causes social dispersal, and triggers compensatory breeding cycles. Mature solitary boars (*Keiler*) comprise no more than 5% of the bag¹⁵.

For roe deer, a strict 1 : 1 sex ratio is maintained, with juveniles (*Kitze*) and yearlings (*Schmalrehe/Jährlinge*) comprising 60% of the extraction, and adult does and bucks making up the remaining 40%¹⁵. For red and fallow deer, calves and yearlings must comprise 60% to 70% of the harvest, restricting the culling of mature antlered stags to post-rut individuals (*Altersklasse IV*, ≥ 10 years)¹⁵.

Non-Ballistic Harvesting Methodologies and Animal Welfare

The permanent absence of manufactured metallic cartridges, percussion primers, and propellant powders requires a return to mechanical projectile weapons, static traps, and collective driven hunts. Each technique involves distinct biomechanical parameters and animal welfare implications that directly determine meat quality via post-mortem muscle physiology.

Traditional Archery and Heavy Crossbows

The mechanical bow and crossbow function on principles distinct from firearm terminal ballistics. While expanding bullets impart substantial kinetic energy (1,500 – 4,000 J), causing hydrostatic shock waves, cavitation, and concussive organ failure¹⁷, arrows and bolts operate as low-velocity, high-momentum cutting instruments.

The kinetic energy (KE) and linear momentum (p) of an archery projectile are defined by:

$$KE = \frac{1}{2}mv^2$$

$$p = mv$$

A forged steel leaf-blade broadhead with an acute cutting edge (mass 125 to 175 grains),

affixed to a dense Scots pine or ash shaft (total projectile mass $m = 32 - 45 \text{ g}$ / 500–700 grains), discharged from a 70-pound wooden longbow or a 150-to-220-pound recurve steel-limbed crossbow, produces an initial launch velocity of 55 to 75 m/s. This yields:

$$KE = \frac{1}{2}(0.040 \text{ kg})(65 \text{ m/s})^2 \approx 84.5 \text{ Joules}$$

$$p = (0.040 \text{ kg})(65 \text{ m/s}) = 2.6 \text{ kg} \cdot \text{m/s}$$

While 85 Joules cannot penetrate the dense, pneumatic frontal sinus bone of an adult boar or stag¹⁷, it possesses ample momentum to pass completely through the intercostal ribs of the thorax. Broadheads cause death through vascular transection rather than concussive trauma. A sharp broadhead severs the microvasculature of both pulmonary lobes (inducing bilateral tension pneumothorax) and transects the cardiac muscle, aortic arch, or pulmonary trunk¹⁷. This precipitates rapid hypovolemic shock, collapsing mean arterial pressure, terminating oxygen delivery to the brain, and causing unconsciousness and clinical death within 15 to 45 seconds¹⁷.

When executed from elevated platform tree stands (*Hochsitze*) or concealed blinds, broadside thoracic archery shots rarely provoke acute panic. The quarry often runs only a short distance before collapsing. As a result, catastrophic ante-mortem glycogen depletion is avoided, leaving glycogen reserves intact for post-mortem glycolysis and optimal muscle acidification¹⁹.

Visceral strikes, however, cause prolonged agony, violate humane standards, and trigger extreme catecholamine release, which impairs meat quality.

Collective Driven Hunts: The Lineage of Zeugjagd, Lappjagd, and Cordage Nets

The most labor-efficient methodology for harvesting game without ammunition is the collective driven hunt, adapted from the historical *Zeugjagd* and *Einstelljagd* practiced across the Electoral March of Brandenburg from the sixteenth through the eighteenth centuries²¹.

This operational sequence relies on non-metallic physical barriers:

1. **Lappjagd (Flap Drives):** Hemp ropes are suspended at human chest height (1.2 to 1.5 meters) along designated forest corridors, strung with hanging rectangles of linen or canvas (*Jagdplatten*, $30 \times 50 \text{ cm}$) spaced 2 to 3 meters apart. Cervids avoid these optical barriers, which are treated with wood ash, tallow, or human scent, allowing beaters to direct moving herds along pre-cleared forest rides (*Flügel*).
2. **Nets and Corrals (*Tümmletetze*):** Heavy cordage nets braided from four-ply hemp or flax twine, with a mesh width of 12 to 15 cm and a height of 2.0 to 2.5 meters, are suspended in dense undergrowth along natural bottlenecks. The bottom line is staked to the soil, while the top line is suspended from branched poles. As fleeing game contacts

the barrier, the top line releases, entangling the quarry in the falling mesh.

3. **Dispatch Protocol (*Abfangen*):** Handlers stationed behind blinds approach the entangled animals. Boars are pinned using heavy, forked ash poles and dispatched with a wild boar spear (*Saufeder*). The forged leaf blade, fitted with a cross-guard (*Knebel*) to arrest over-penetration, is thrust into the thoracic inlet behind the left shoulder to sever the great vessels at the base of the heart¹⁷. Cervids are dispatched by pithing (*Abnicken*)—driving a short blade (*Knicker*) directly between the occipital condyles and the atlas vertebra (C1), severing the brainstem and arresting autonomic functions¹⁷.

Driven hunts subject animals to sustained physical exertion and stress. This exertion depletes cellular adenosine triphosphate (ATP) and muscle glycogen, leading to premature lactic acid

accumulation while the core body temperature is elevated ($> 39^{\circ}\text{C}$). This dynamic causes pale, soft, and exudative (PSE) defects in boar meat, or dark, firm, and dry (DFD) meat in exhausted cervids¹⁹. To preserve meat quality, driven operations must be limited to short drives

under 45 minutes during cool weather below 5°C .

Trapping Mechanisms: Box Traps, Corrals, and Snares

Passive mechanical traps enable continuous animal harvesting while human labor is directed toward agriculture and forestry:

- **Saufang (Wild Boar Corral Trap):** Circular enclosures built from vertical pine timbers (10 to 15 cm diameter) driven 80 cm into the ground, bound with split-pine rails to a height of 1.8 meters. The entrance features a weighted drop-gate (*Falltor*) suspended by hemp rope connected to a mechanical root-treadle. Baited with acorns, cracked maize, or fermented rye, entire sounders (*Rotten*) enter the pen. Once trapped, animals are selectively dispatched from an elevated exterior walkway using spears or crossbows. This method permits the sparing of reproductive sows while capturing subadult biomass.
- **Heavy Box Traps (*Kastenfallen*):** Constructed from heavy pine or oak boards (5 cm thickness), measuring $2.0 \times 0.8 \times 1.0$ meters. Configured with vertical falling gates and a center treadle, these traps are set along game paths for roe deer or subadult boars. The darkened interior keeps the trapped animal calm, reducing impact trauma caused by escape attempts.
- **Pitfall Traps (*Hirschgruben* / *Wolfsgruben*):** Deep excavations (2.5 to 3.5 meters deep) with sheer walls, covered by light lattices of twigs, pine needles, and turf. Pitfalls compromise animal welfare; falling cervids frequently suffer bone fractures, internal hemorrhaging, and dislocated joints. The resulting distress causes deep muscular hematomas and systemic blood splash (*Ecchymosis*), ruining prime culinary cuts¹⁹. Pitfalls are therefore restricted to predator defense perimeter ditches around livestock pens.
- **Snares (*Schlingen*):** Twisted steel wire or braided cordage set along trails. Under ethical hunting standards, snares are prohibited due to prolonged strangulation and trauma. From a culinary perspective, meat from snared animals is severely compromised: violent agonal struggling causes venous pooling, prevents proper blood drainage, and leaves high residual blood in muscle capillary beds¹⁷.

Legal Frameworks, Governance Transitions, and Commons Management

The Statutory Baseline: German Hunting Law

The contemporary regulatory framework for hunting across the region is established by the Federal Hunting Act (*Bundesjagdgesetz*, BJagdG) and the Brandenburg State Hunting Act (*Brandenburgisches Jagdgesetz*, BbgJagdG)⁷. This legal corpus is built upon the *Reviersystem* (territorial hunting district system), which links the right to hunt (*Jagdausübungsrecht*) to land ownership while preventing fragmented exploitation through mandatory surface area thresholds:

- *Eigenjagdbezirk* (Private hunting district): Contiguous land area of at least 75 hectares owned by a single individual or entity.
- *Gemeinschaftlicher Jagdbezirk* (Community hunting district): The consolidated aggregate of all contiguous parcels within a municipality measuring at least 150 hectares, managed collectively by landowners organized into a statutory public corporation (*Jagdgenossenschaft*).

Central to this legal framework is the principle of game stewardship (*Hegepflicht*), codified in § 1 BJagdG. This principle mandates that hunters maintain balanced, species-rich wildlife populations adapted to regional ecosystems, actively preventing severe agricultural browsing (*Verbiss*) and timber peeling (*Schälen*)¹³.

Harvest yields are legally controlled through statutory shooting plans (*Abschusspläne*), which are formulated by local wildlife associations (*Hegegemeinschaften*), certified by the lower hunting authority (*Untere Jagdbehörde*), and legally binding for all cloven-hoofed game (*Schalenwild*)¹⁵.

Statutory hunting seasons (*Jagdzeiten*) reflect wildlife reproductive cycles: roe bucks and yearlings are hunted from April 16 to January 15; adult does from August 1 to January 15; red and fallow deer from August 1 to January 15; while wild boar may be taken year-round to control disease and agricultural losses, subject to § 22 Abs. 4 BJagdG, which protects mothers nursing dependent offspring⁷.

The Post-Collapse Paradigm: Managing the Sylvatic Commons

Following the collapse of central state authority and regulatory oversight, the legal framework is vulnerable to rapid degradation. Uncontrolled hunting by a stressed population can quickly deplete wild stocks, replicating the wildlife extirpations observed in Central Europe during the Thirty Years' War (1618–1648) and the social breakdowns of 1945–1946.

To protect the wildlife resource base, the community must transition from statutory governance to customary sylvatic law, adapting historical *Forst- und Jagdverwaltung* institutions:

1. **The Village Hunting Cooperative (*Jagdgenossenschaft*)**: All hunting rights across the 4,000-hectare communal territory are placed under communal governance. Individual unauthorized hunting is treated as resource destruction and biosecurity endangerment.
2. **The Office of the Head Gamekeeper (*Forstmeister / Revierförster*)**: Qualified

gamekeepers are appointed by the village council. These wardens manage collective drives, inspect fixed traps, monitor sounder dynamics, conduct population censuses, and enforce species quotas.

3. **Rotational Wildlife Sanctuaries (*Wildschonbezirke*):** A minimum of 30% of the woodland canopy (roughly 1,200 hectares) is designated as a permanent, undisturbed wildlife sanctuary. Human entry into these sectors is prohibited, providing undisturbed breeding and bedding cover to preserve herd recruitment despite harvest pressure on periphery lands.
4. **Integration of *Waidgerechtigkeit* into Customary Law:** The historical unwritten code of hunting ethics—*Waidgerechtigkeit*—is incorporated into communal law. Strict prohibitions against killing lead sows (*Leitbachen*) or leading hinds (*Leittiere*) are enforced. Lead sows guide group movement and maintain social order; their removal causes sounders to scatter into agricultural crops, increasing food losses.

Post-Mortem Biochemistry, Field Dressing, Hanging, and Maturation

Field Dressing (*Aufbrechen*) and Sanitary Evisceration

Transforming a harvested wild ungulate into high-quality meat without access to commercial slaughterhouses requires precise field dressing to prevent contamination of sterile muscle by gastrointestinal bacteria such as *Escherichia coli*, *Salmonella enterica*, and *Clostridium perfringens*¹⁷. Complete evisceration must occur within 60 minutes of death.

The primary dressing sequence begins with exsanguination:


- **Exsanguination:** If the animal was taken by trap or broadhead, an immediate thoracic stick is performed. The point of a breaking knife is inserted cranial to the manubrium sterni, directed into the chest cavity at a 45-degree angle to sever the brachiocephalic trunk and bicarotid arteries at the aortic arch¹⁷. Evacuating 40% to 60% of total circulating blood volume extracts latent body heat and removes heme iron, which acts as a primary catalyst for lipid oxidation and flavor degradation¹⁷.
- **Two-Knife Protocol:** The external hide carries soil, hair, and bacterial spores. The field dresser uses two dedicated knives: Knife A is used exclusively to cut through the skin, while Knife B, kept sanitized in scalding water, is reserved for sterile cavity and visceral work¹⁷.
- **Circum-Anal Coring (*Ringeln*):** Rather than cleaving the pelvic aitch bone (*Schloss*) with an axe—which creates sharp bone shards and risks rupturing the colon or urinary bladder—the bunging technique (*Ringeln*) is applied¹⁷. The operator makes a circular incision through the perianal skin around the anus and vulva, releasing connective tissues in the pelvic canal without puncturing the rectal wall¹⁷. The free terminal rectum is drawn outward, ligated with cord, and eased forward through the pelvic channel into the abdominal cavity¹⁷.
- **Ventral Opening:** The animal is positioned on an incline or hung by its hind legs. An incision is made along the linea alba from the xiphoid process to the pelvis. The operator

inserts two fingers of the non-dominant hand inside the incision to lift the abdominal wall away from the viscera, guiding the outward-facing knife blade between the fingers to shield the stomachs and intestines from punctures¹⁷.

- **Visceral Evacuation:** The esophagus (*Weasand*) is freed from its tracheal attachments in the neck, knotted, and stripped downward into the thorax¹⁷. The diaphragm is incised along its costal margin¹⁹. By pulling down on the ligated esophagus and rectum, the rumen/stomach mass, intestines, and liver roll out intact onto a clean linen sheet without spillage¹⁹. The pluck (heart, lungs, trachea) is removed immediately afterward¹⁹. The cavity is wiped dry with clean, sterile linen; washing with untreated cold surface water is prohibited, as moisture encourages the colonization and movement of psychrotrophic *Pseudomonas* species across meat surfaces¹⁷.

Post-Mortem Muscle Biochemistry: Glycolysis, Rigor Mortis, and Ageing

The conversion of warm muscle into tender, palatable game meat is governed by post-mortem glycolysis and intracellular energy depletion¹⁹. In the living animal, resting skeletal muscle maintains a neutral pH near 7.2. Following circulatory arrest, tissue oxygen drops to zero, shifting cellular metabolism to anaerobic glycolysis¹⁹. Sarcoplasmic glycogen stores are metabolized into lactic acid ($C_3H_6O_3$) via the Embden-Meyerhof-Parnas pathway²⁴:



This generation of protons drops muscle pH. In an animal dispatched with rested glycogen reserves, the ultimate pH (pH_u) stabilizes between 5.4 and 5.7 within 12 to 24 hours¹⁹. This acidification serves two essential preservation functions:

1. Approaching the isoelectric point of myosin ($pI = 5.3$) reduces the net electrical charge on the filaments, causing the protein matrix to contract and expel surface water. This creates a dry surface film (*Trockenrandschicht*) that impedes bacterial growth²⁴.
2. The acidic environment inhibits deep putrefactive pathogens, particularly *Clostridium putrefaciens*²⁴.

As glycogen is consumed, ATP regeneration ends. When cellular ATP concentrations decline below $1.0 \mu\text{mol/g}$ of muscle, the sarcoplasmic reticulum can no longer pump calcium ions (Ca^{2+}) out of the sarcoplasm¹⁹.

This calcium influx exposes binding sites on the actin chains, allowing myosin heads to lock into place and form permanent actomyosin cross-bridges²⁴. The carcass enters *rigor mortis*, becoming rigid and inextensible¹⁹. Cooking meat during active rigor causes severe toughening, as thermal shrinkage compacts sarcomeres and expels moisture²⁴.

Chilling Thermodynamics and Structural Defects

Controlling carcass temperatures during the onset of rigor mortis is critical to prevent two irreversible processing defects:

- **Cold Shortening (*Kälteverkürzung*):** If a carcass is chilled below 10°C while intracellular ATP levels remain elevated and the pH is above 6.0, cold shock destabilizes the sarcoplasmic reticulum, releasing calcium into the myofibrillar space¹⁹. In the presence of unspent ATP, the muscle contracts forcefully, shortening sarcomeres by up to 50% of their resting length²⁴. This produces severe, irreversible toughness that cannot be resolved by subsequent ageing²⁴.
- **Heat Shortening (*Wärmeverkürzung*):** If the carcass is held at warm ambient temperatures ($> 35^{\circ}\text{C}$) during early post-mortem glycolysis, rapid lactic acid production denatures sarcoplasmic and myofibrillar proteins¹⁹. This denaturation destroys water-holding capacity, resulting in pale, watery, and structurally compromised meat that loses excessive moisture during cooking¹⁹.

To avoid both issues, the carcass must follow a structured chilling profile: internal core temperatures must remain above 10°C for the first 10 to 12 hours post-mortem, but drop below 7°C within 24 hours²⁷.

Endogenous Proteolysis and Deep-Cellar Maturation

Following the resolution of rigor mortis (24 to 48 hours post-mortem), the tenderization and flavor development phase—maturation (*Abhängen/Reifung*)—begins. Tenderization does not reverse actomyosin bonds; rather, it is driven by endogenous proteases that dismantle the structural cytoskeleton of the muscle fibers²⁴:

- **Calpains (μ -calpain and m-calpain):** Calcium-activated neutral cysteine proteases. Activated by the calcium ions released during rigor, calpains selectively cleave the structural proteins that support the myofibril, degrading titin, nebulin, and desmin, and liberating troponin-T²⁴. This breaks the lateral register of the muscle fibers, tenderizing the meat without hydrolyzing tough connective tissue²⁴.
- **Cathepsins (B, D, H, and L):** Lysosomal acid proteases released as cell membranes break down²⁹. Operating optimally at low pH ($\text{pH} \leq 5.5$), cathepsins cleave both myofibrillar proteins and the terminal telopeptides of intramuscular collagen, facilitating tenderization over extended hanging periods²⁴.

In a low-technology environment, controlled cooling relies on two passive structures: deep earth cellars (*Erdkeller*) and winter ice-houses (*Eiskeller*). Earth cellars excavated 4 to 6 meters into subsoil sands maintain stable year-round temperatures of $8^{\circ}\text{C} - 10^{\circ}\text{C}$.

For colder storage, an ice-house filled with winter lake ice packed beneath a 1-meter layer of dry Scots pine sawdust maintains internal temperatures between 1°C and 4°C through late

summer. Carcasses should be hung by the pelvic obturator foramen (the "tenderstretch" method) rather than the Achilles tendon. Pelvic suspension stretches the lumbar spine, preventing the *Longissimus dorsi* and haunch muscles from contracting, which reduces Warner-Bratzler shear force values²⁴.

Species	Carcass Weight (Dressed, kg)	Minimum Ageing Period (Days)	Optimal Maturation Window (Days)	Maximum Safe Hanging Limit (Days)	Primary Proteolytic Profile
Roe Deer (<i>Capreolus capreolus</i>)	12.0 – 18.0	3	5 – 7	10	Rapid μ -calpain activation; low connective tissue density
Wild Boar (<i>Sus scrofa</i>)	25.0 – 65.0	4	7 – 10	14	Rapid surface drying; fat cap protects muscle from desiccation ¹⁹
Fallow Deer (<i>Dama dama</i>)	30.0 – 55.0	5	8 – 12	16	Intermediate fiber diameter; stable catheptic breakdown
Red Deer (<i>Cervus elaphus</i>)	60.0 – 140.0	7	12 – 18	21	High collagen cross-linking; requires

					extended structural relaxation ²⁴
European Hare (<i>Lepus europaeus</i>)	2.5 – 4.0	2	4 – 5	7	Skin left intact during hanging to prevent severe surface desiccation

Sylvatic Pathogens, Parasitology, and Radiometric Risk Mitigation

Wild game meat from post-industrial landscapes carries risks from endemic zoonoses, epizootic viruses, and radiological fallout residues. Safeguarding public health requires structured diagnostics, biosecurity protocols, and culinary-chemical remediation.

***Trichinella spiralis* and *Trichinella britovi* in Wild Boar**

Trichinella nematodes cause human trichinellosis, a parasitic disease characterized by acute gastroenteritis, fever, periorbital eyelid edema, severe muscle inflammation, and potentially fatal myocarditis or encephalitis¹⁷. In Brandenburg, *Trichinella spiralis* and the freeze-tolerant *Trichinella britovi* circulate in wild populations via the scavenging of carcasses by wild boars, red foxes (*Vulpes vulpes*), and raccoon dogs (*Nyctereutes procyonoides*)².

Infective larvae ingested through raw or undercooked meat excyst in the stomach, develop into adult worms in the small intestine, and produce newborn larvae that migrate via the circulatory system into striated muscle, where they encapsulate within nurse cells¹⁷.

To detect larvae without commercial automated equipment, the community can use an adapted low-technology artificial digestion protocol based on ISO 18743:2015¹⁹:

- Reagents: 1,000 mL of water heated to $46^{\circ}\text{C} - 48^{\circ}\text{C}$, 16.0 g of high-activity pepsin (1:10,000 NF), and 10.0 mL of concentrated hydrochloric acid (HCl , approximately 37%, salvaged or distilled from salt and vitriol)¹⁹.
- Sampling: Excise a 5.0 g sample of clean muscle tissue from the muscular pillar of the diaphragm (*Zwerchfellpfeiler*) at the transition to the central tendon¹⁷. If unavailable, take samples from the tongue or internal masseter muscles¹⁷.
- Digestion: Finely chop the muscle into pieces smaller than 2 mm. Mix the sample into the acidified pepsin solution ($\text{pH} \approx 1.5 - 2.0$) within a glass vessel¹⁹. Maintain the

mixture in an insulated water bath at $44^{\circ}\text{C} - 46^{\circ}\text{C}$ for 30 to 45 minutes, agitating continuously with a glass rod or mechanical paddle¹⁹. The pepsin-acid solution hydrolyzes the myofibrillar and connective proteins, releasing any encysted larvae¹⁷.

- Sedimentation and Examination: Pour the digested fluid through an $180\text{-}\mu\text{m}$ wire sieve into a glass sedimentation funnel. Allow the liquid to rest for 30 minutes. Draw off the bottom 40 mL of sediment into a flat watch glass or Petri dish. Examine the sample using a salvaged stereomicroscope or botanical field loupe at $20\times$ to $40\times$ magnification. *Trichinella* larvae appear as coiled, comma-shaped or spiral worms approximately 1.0 mm long and $30\text{ }\mu\text{m}$ wide¹⁷.

If laboratory chemicals are unavailable, the historical compression squash method (*Kompressorium*) must be employed¹⁷. Using fine scissors, cut 28 oat-grain-sized longitudinal tissue sections ($2 \times 10\text{ mm}$) along the muscle fibers of the diaphragm pillars¹⁷. Compress the samples between two thick glass plates tightened with thumbscrews until translucent. Inspect each section under $30\times$ to $50\times$ magnification for encapsulated larvae¹⁷. Any positive sample mandates the condemnation and incineration of the entire carcass.

Freezing is not a dependable control for *Trichinella britovi*, which resists sub-zero temperatures. All suid meat must reach an internal core temperature of at least 71°C for 60 seconds or 65°C for 15 minutes to guarantee thermal destruction¹⁹.

African Swine Fever (ASFV / Afrikanische Schweinepest)

African Swine Fever is caused by a large, enveloped DNA virus of the family *Asfarviridae*. The pathogen is not zoonotic and cannot infect humans; consuming meat from an infected wild boar poses zero risk of human illness.

However, ASFV is a highly contagious, fatal hemorrhagic disease of suids, causing mortality rates approaching 100%⁸. Brandenburg is an active front line for ASFV in Western Europe, with thousands of cases confirmed across eastern districts such as Oder-Spree, Spree-Neiße, and Märkisch-Oderland⁸.

The critical risk is biosecurity: feeding, handling, or improperly disposing of infected wild boar tissues can transmit the virus into the community's domestic pig herds, destroying domestic pork production.

The ASFV virion exhibits extreme environmental stability:

- Persists in chilled fresh pork for up to 150 days.
- Survives in dry-cured hams, unsmoked bacon, and dried sausages for 180 to 400 days.
- Remains viable in frozen meat for years and in contaminated mud or feces for months.
- Inactivated reliably only by heating to 60°C for 30 minutes, or boiling (100°C) for 5 minutes.

Hunters must perform field post-mortem examinations for indicative lesions: severe splenomegaly (a dark purple, swollen, friable spleen), petechial hemorrhages on the cortical surface of the kidneys ("turkey-egg kidney"), and swollen, marbled-patterned gastrohepatic and mesenteric lymph nodes.

Any animal displaying these signs must be quarantined, and the carcass, blood, and offal destroyed by high-temperature wood pyrolysis followed by burial under quicklime. Personnel processing wild boars must pass through wood-ash lye footbaths and change garments before entering domestic livestock enclosures. Under no circumstances may raw wild boar scraps enter domestic pig feeding troughs.

Cestode Zoonoses: *Echinococcus multilocularis* and *Echinococcus granulosus*

Echinococcosis presents significant health risks across the region:

- *Echinococcus multilocularis* (Fox tapeworm / *Fuchsbandwurm*): Adult tapeworms reside in the intestines of red foxes, raccoon dogs, and domestic dogs. Microscopic oncosphere eggs shed in feces contaminate ground vegetation, fallen berries, and water sources. Ingestion by humans leads to alveolar echinococcosis, forming destructive, tumor-like parasitic growths within the liver parenchyma that are fatal without pharmaceutical or surgical intervention.
- *Echinococcus granulosus* (Cystic echinococcosis): Cycles between wild canids (wolves, feral dogs) and wild ungulates. Ingestion of eggs causes large, fluid-filled hydatid cysts within pulmonary and hepatic tissues.

Field dressers must wear protective leather or waxed-linen gloves during evisceration. The liver and lungs of all harvested game must be palpated and sliced; organs containing hydatid cysts must be excised and cremated²⁷. Raw offal must never be fed to hunting hounds.

Community dogs must receive regular anthelmintic treatments using decoctions of wormwood (*Artemisia absinthium*), male fern rhizome (*Dryopteris filix-mas*), or salvaged veterinary praziquantel. Forest herbs, roots, and mast intended for consumption must be thoroughly cooked; cold-water washing does not reliably remove adhesive cestode eggs.

Radiocaesium (^{137}Cs) Persistence in Brandenburg Wild Boar

Following the 1986 Chernobyl disaster, atmospheric fallout deposited significant quantities of

Caesium-137 (^{137}Cs , half-life $t_{1/2} = 30.17$ years) across Central Europe¹. While cervids show low contamination levels due to their consumption of above-ground foliage, wild boars in Brandenburg frequently exhibit activity concentrations exceeding the European safety threshold of 600 Bq/kg fresh mass¹.

This persistent contamination—termed the "wild boar paradox"—is driven by the animal's winter foraging ecology and soil chemistry¹. Over decades, caesium ions percolated through Brandenburg's sandy Arenosols, settling at the interface between the organic humus layer and the mineral subsoil (5 – 20 cm depth)¹.

This horizon is colonized by the subterranean mycorrhizal deer truffle (*Elaphomyces*

granulatus)¹. The mycelial network of *Elaphomyces* hyperaccumulates radiocaesium, producing fruiting bodies that register radioactivity levels between 2,000 and 10,000 Bq/kg¹. During winter, when surface acorns are exhausted, boars root deeply, consuming deer truffles as a primary calorie source¹. The ingested caesium is absorbed into the bloodstream and deposited throughout skeletal muscle tissue as a chemical analogue to potassium (K^+)¹. To reduce internal exposure without laboratory spectrometers, the community can employ culinary-chemical leaching techniques¹:

- **Aqueous Leaching Kinetics:** Caesium exists within muscle cells as an unbound, water-soluble monovalent cation ($^{137}Cs^+$)¹⁴. When meat is soaked or simmered in water, caesium leaches into the surrounding fluid along osmotic gradients¹⁴.

- **Processing Retention Factors:**

The food processing retention factor (F_r) is expressed as:

$$F_r = \frac{C_{\text{cooked}} \cdot m_{\text{cooked}}}{C_{\text{raw}} \cdot m_{\text{raw}}}$$

where C represents activity concentration and m is tissue mass¹⁴. Grilling yields an $F_r \approx 0.90$ (retaining 90% of total radioactivity while concentrating it through moisture evaporation)¹⁴. Steaming achieves an $F_r \approx 0.50$ (50% removal)¹⁴. Boiling produces an $F_r \approx 0.40 - 0.60$ (40% to 60% reduction)¹⁴.

- **Culinary Decontamination Protocol:** Boar meat from winter hunts is diced into small cubes ($2 \times 2 \text{ cm}$) to maximize the surface area for diffusion¹⁴. The meat is submerged in a 3% aqueous salt solution ($NaCl$) at a 1 : 5 meat-to-water ratio for 24 to 48 hours at $4^\circ C$, changing the brine bath twice¹⁴. This process extracts up to half of the cellular caesium into the waste brine¹⁴, which is discarded away from crops. Alternatively, the meat is placed in cold water, brought to a boil for 10 minutes, and the initial water discarded¹⁴. Subsequent slow braising in fresh liquid produces a safe, high-quality dish, removing up to 70% of the initial radiocaesium load¹⁴. If salvaged supplies of Prussian Blue (ferric ferrocyanide, $Fe_7(CN)_{18}$) are available, adding 1 to 2 grams per kilogram of cooked meat irreversibly binds $^{137}Cs^+$ ions into an insoluble complex, preventing intestinal uptake during digestion.

Manual Seam Butchery, Carcass Breakdown, and Byproduct Valorisation

Artisanal manual butchery differs fundamentally from mechanized band-saw fabrication. Rather than sawing straight, cross-sectional steaks across muscle groups—which splinters bone and mixes disparate muscle fiber types—manual seam butchery (*Präzisions-Nahtschnur-Zerlegung*) follows natural connective tissue planes²⁴. This technique maximizes yield, reduces kerf loss, and isolates individual muscles according to their connective-tissue content and culinary use²⁴.

Tool Metallurgy and Manual Ergonomics

Executing precise manual butchery without power tools requires specific edged tools²⁴:

- **Boning and Seam Knives:** Forged from high-carbon spring or tool steel (such as recycled leaf-spring 5160 or 1095 steel), hardened to 58–60 HRC, and sharpened to an acute $15^{\circ} - 20^{\circ}$ V-bevel²⁴. The thin, semi-flexible blade traces fascial sheets without cutting into adjacent muscle tissue²⁸.
- **Heavy Cleaver:** Ground to an obtuse $50^{\circ} - 60^{\circ}$ inclusive convex profile²⁴. The mass of the heavy spine (3 to 4 mm thick) absorbs impact shock when splitting the pubic symphysis or severing vertebrae without chipping or rolling the edge²⁴.
- **Butcher's Bow Saw:** Fitted with high-tension carbon steel blades; 4 teeth per inch (TPI) for chilled, firm carcasses, and 3 TPI for heavy bone sectioning²⁴. The thin blade minimizes kerf loss and limits bone dust²⁴. Bone dust contains fats and cellular debris that accelerate bacterial growth (*Pseudomonas* spp.), causing deep putrefaction known as "bone sour"²⁴. Cut bone faces must be scraped clean with a flat steel scraper²⁸.

Anatomical Breakdown of the Cervid Carcass

Following chilling and the resolution of rigor mortis, the skinned cervid carcass is quartered between the 12th and 13th ribs, leaving the 13th rib on the hindquarter to support the loin primal²⁴.

The forequarter is deconstructed into functional culinary cuts:

- **Shoulder (*Blatt / Schulter*):** The forelimb is lifted outward and separated from the thorax by following the connective tissue plane beneath the *Serratus ventralis*; no bone sawing is required, as ungulates lack a clavicle²⁴. The blade is deboned to isolate the *Supraspinatus* (chuck tender / *Falsches Filet*) and the *Infraspinatus* (top blade / *Schulterschmelze*)²⁴. The *Infraspinatus* features fine, tender fibers and can be grilled or seared once its central internal connective tissue strip is excised²⁴. The remaining shoulder trimmings are reserved for braising, terrines, or sausage emulsions²⁴.
- **Neck (*Träger / Hals*):** Severed at the C7-T1 intervertebral junction with a bow saw²⁴. Rich in cervical vertebrae, intermuscular fat, and ligamentous tissue (*Ligamentum nuchae*), the neck is braised whole on the bone or broken down for hearty stews (*Pfeffer*)²⁸.
- **Rib and Brisket (*Brust / Querrippe*):** The sternum and costal cartilage plates (*Pectoralis*

complex) are separated from the loin. High in structural collagen, these cuts are used for enriched game stocks, glazes, or fine mince²⁴.

The hindquarter and saddle yield the prime roasting cuts:

- **The Saddle / Loin (Rücken):** Extends from T6 through the lumbar spine to the sacrum²⁴. The tenderloin (*Psoas major* / *Filet*) rests along the ventral side of the lumbar transverse processes; non-locomotive and minimally worked, it is the most tender muscle in the carcass²⁴. The dorsal muscle (*Longissimus dorsi* / *Lachs*) is freed from the spinous and transverse processes using a boning knife²⁴. It can be portioned into medallions or prepared whole as a bone-in *Rehrücken* roast²⁴.
- **The Haunch / Leg (Keule):** Separated from the pelvis through the sacroiliac joint and the acetabular hip socket²⁴. Seam butchery divides the leg into four primary muscles²⁴:
 - **Top Round (Oberschale):** Comprising the *Semimembranosus* and *Adductor*. Large, fine-grained, and lean; suitable for fine roasts, escalopes, or cured charcuterie²⁴.
 - **Bottom Round (Unterschale):** Formed by the *Biceps femoris*. Coarser grain with a heavy connective outer sheet; requires slow roasting or braising²⁴.
 - **Knuckle (Nuss / Kugel):** Formed by the *Quadriceps femoris*. Dense and structural; sliced into uniform steaks or cured²⁴.
 - **Rump (Hüfte):** Formed by the *Gluteus medius*. Highly tender; roasted whole or pan-seared²⁴.
 - **Shank (Haxe):** Formed by the *Flexor* and *Extensor digitorum* wrapped around the tibia²⁴. Dense with Type I collagen, requiring extended braising to melt into gelatin²⁴.

Primal / Subprimal Cut	Anatomical Muscle Group	Yield Mass (kg)	% of Live Weight	Connective Tissue Level	Optimal Culinary Method
Tenderloin (<i>Filet</i>)	<i>Psoas major</i> [cite: 24]	0.40	2.0%	Very Low (WBSF <2.5 kg) ²⁴	Quick high-heat sear; rare (52°C) ²⁴
Saddle / Loin (<i>Rücken</i>)	<i>Longissimus dorsi</i> [cite: 24]	1.80	9.0%	Low (WBSF 3.0–3.8 kg) ²⁴	Dry-heat roast; medium-rare (54°C) ²⁴
Top Round (<i>Oberschale</i>)	<i>Semimembranosus/Adductor</i> [cite: 24]	1.60	8.0%	Low-Moderate	Dry roast, schnitzel, cold

					curing ²⁴
Bottom Round (Unterschale)	<i>Biceps femoris</i> [cite: 24]	1.50	7.5%	Moderate (dense fascia) ²⁴	Slow braise or fine comminution ²⁴
Knuckle & Rump (Nuss & Hüfte)	<i>Quadriceps /Gluteus</i> [cite: 24]	1.70	8.5%	Low-Moderate	Pan roasting, medallion steaks ²⁴
Shoulder (Blätter)	<i>Triceps, Supraspinatus</i> [cite: 24]	1.80	9.0%	High	Braised ragoût, terrines, pasties ²⁴
Neck & Flank (Hals & Dünnung)	<i>Cervical group / Abdominal</i> [cite: 24]	1.60	8.0%	High	Stewing (Hasenpfeffer style), stock ²⁴
Shanks (Haxen)	<i>Flexor/Extensor digitorum</i> [cite: 24]	0.80	4.0%	Maximum (heavy collagen) ²⁴	Slow-simmered gelatin broths ²⁴
Bones & Skeletal Frame	Complete skeleton ²⁸	4.20	21.0%	Non-edible structural	Pyrolysis, brown stock, bone tools ²⁸
Viscera & Offal	Liver, heart, kidneys, blood ²⁷	1.60	8.0%	Varied (high-nutrient)	Pan-fried liver, blood pudding, liver pâté
Hide, Skin, Hooves	Dermis and distal legs ²⁷	3.00	15.0%	Industrial byproduct	Tanned buckskin, marrow glue ²⁷

Complete Byproduct Valorisation

In a self-sustaining agrarian economy, all parts of the harvested animal are utilized:

- **Blood (*Schweiß*):** Captured during exsanguination into glazed earthenware containing salt, stirred with birch twigs to prevent uneven clotting²⁷. Blended with diced wild boar backfat, cooked pearl barley or spelt, onions, and marjoram, it is stuffed into cleaned intestines to produce traditional black pudding (*Rotwurst / Blutwurst*).
- **Suet and Visceral Fats (*Unschlitt / Talg*):** Cervid visceral and perirenal fat contains high levels of saturated stearic acid ($C_{18:0}$), giving it a firm texture and high melting point ($45^{\circ}\text{C} - 50^{\circ}\text{C}$)³³. It is rendered into tallow for candles, leather water-repellents, and bowstring waxes. In contrast, boar caul fat (*Netzfett*) and backfat are salted for culinary cooking fats²⁹.
- **Sinews and Fascia:** The Achilles tendons (*Tendo calcaneus*) and broad back fascia of stags are dried, pounded with wooden mallets to separate the collagen fibrils, and spun into cordage, bowstrings, or applied with hot hide glue as composite backings for wooden hunting bows.
- **Hide Tanning:** Cervid hides are soaked in slaked wood-ash solutions to remove hair, scraped over a wooden beam, and brain-tanned (using the animal's emulsified brain lipids) or bark-tanned with oak and pine bark decoctions, producing supple buckskin for garments and durable sole leather.
- **Bones and Antlers:** Marrow bones (femur, humerus) are split to harvest nutrient-dense marrow²⁸. Bone scraps are simmered for stock, pyrolyzed into bone char (*Knochenkohle*) for water filtration, or carved into needles, buttons, and trap triggers. Shed antlers are worked into tool handles and crossbow nuts.

Culinary Treatment of Game for the Civilised Table

Elevating wild game to fine cuisine requires matching muscle structure to cooking physics.

Cervid meat is exceptionally lean, rarely containing more than 2% intramuscular fat, and locomotive cuts carry dense connective collagen²⁴. Without proper technique, dry cooking drives off internal moisture, leaving the meat tough and dry²⁴.

Marinades and Equilibrium Brining

In classical cuisine, marinades often masked unpleasant odors from poorly aged meat. In this kitchen, marination serves to tenderize tissues and balance flavors:

- **Acidic Marinades (*Beizen*):** Formulated from dry red wine, cider, or lacto-fermented whey, infused with crushed juniper berries (*Juniperus communis*), allspice, cracked peppercorns, wild thyme, and sliced root vegetables. The acidic medium ($\text{pH} \approx 3.5 - 4.2$) swells the myofibrillar protein lattice, assisting the enzymatic cleavage of connective telopeptides³³. Immersion should not exceed 48 hours; prolonged soaking denatures surface proteins, producing a mushy, discolored exterior²⁴.
- **Equilibrium Wet-Brining:** For roasting whole roe shoulders or waterfowl (mallards,

geese), equilibrium brining (2.0% – 2.5% NaCl and 1.0% honey/sugar based on total meat and water mass) is applied for 24 hours at 3°C³³. Chloride ions bind to myofibrils, increasing negative electrostatic charges and swelling the protein structure, which allows the meat to hold water during roasting³³.

Mechanical Lipid Integration: Larding (*Spicken*) and Barding (*Umwickeln*)

Because cervid loins and haunches lack intramuscular fat, external fats must be added manually³⁵:

- **Larding (*Spicken*)**: Solid pork backfat harvested from the inner layer of the pig's back—which possesses the highest concentration of saturated stearic and palmitic acids and the firmest crystal structure—is cut into thin matchsticks (*Lardons*, 5 × 5 × 80 mm)²⁹. The lardons are seasoned with salt, white pepper, and mace, chilled until firm, and threaded into a split-tip steel larding needle (*Spicknadel*). The needle is pushed through the *Longissimus dorsi* along the muscle grain at 2 cm intervals³⁵. As the roast cooks, the interior fat melts slowly, basting the lean fibers and preventing dryness²⁴.
- **Barding (*Umwickeln*)**: Wrapping the exterior of a lean loin, saddle, or game bird breast with thin sheets of salted fatback³³. The fat shields the surface from direct dry heat, reduces moisture evaporation, and conducts heat gently until the interior reaches doneness²⁴.

Thermodynamics of Slow Braising: Collagen-to-Gelatin Hydrolysis

Tough forequarter primals—the shanks (*Haxen*), neck (*Hals*), and brisket (*Brust*)—contain substantial intramuscular connective tissue composed of mature Type I collagen²⁴. These fibers are stabilized by heat-resistant trivalent pyridinoline (PYR) cross-links that do not break down during brief dry roasting²⁴.

The tenderization of these cuts relies on moist-heat thermodynamics:

- Between 60°C and 65°C, the hydrogen bonds of the collagen triple helix break, causing the fibers to shrink longitudinally and force out cellular water¹⁹.
- When held between 80°C and 90°C in an aqueous liquid, the covalent peptide bonds slowly hydrolyze. The rigid triple helix dissolves into soluble gelatin²⁴. Gelatin binds up to ten times its weight in water, coating individual denatured muscle bundles with an unctuous fluid that mimics juiciness and creates a velvety sauce²⁴.
- **The Pectin-Gelatin Synergy (*Saftgulasch* Paradigm)**: In regional braises like *Wiener Saftgulasch* or Brandenburg *Hasenpfeffer*, a 1 : 1 weight ratio of sliced yellow onions to game meat is maintained²⁴. During an extended 4-hour braise at 85°C, the plant cellular architecture of the onions breaks down, releasing structural pectin²⁴. The

released pectin binds with the dissolved gelatin and rendered suet to create a glossy, thick reduction that requires no starch or flour thickeners²⁴.

Cut Classification	Representative Anatomical Primals	Primary Protein Challenge	Core Thermal Target (°C)	Optimal Cooking Technique
Tender Muscle	Roe saddle, stag tenderloin, duck breast ²⁴	Heat-labile actin contracts above 66°C, expelling water ²⁴	52°C — (Rare/Medium-rare) ²⁴	High-heat pan sear or spit roast with butter basting ²⁴
Dense Roasting Cut	Top round (<i>Oberschale</i>), fallow knuckle ²⁴	Moderate cross-linking; easily dried out by prolonged roasting ²⁴	56°C — (Medium)	Larded with fatback; gentle roasting in an earthen bread oven ³⁵
Collagenous Cut	Shanks (<i>Haxen</i>), neck, shoulder, hare legs ²⁴	Mature pyridinoline cross-links; high shear resistance ²⁴	> sustained for 3–5 hours ²⁴	Slow moist-heat braising in an earthenware pot (<i>Römertopf</i>) ²⁴
Comminuted Matrix	Wild boar trim, neck flaps, venison flank ²⁴	Highly variable fat-to-protein ratio; high microbial exposure ¹⁹	Raw fermented ($a_w < \quad$) or cooked ($> \quad$) ¹⁹	Lactic-fermented dry salami, terrines, or emulsified <i>Blutwurst</i> [cite: 29]

Cured Game Charcuterie: Venison Salami (*Hirschsalami* / *Rehwurst*)

Preserving trimmings from stags and boars into shelf-stable, sliceable raw fermented sausages (*schnittfeste Rohwurst*) employs hurdle technology²⁹:

- Formulation: 65% lean venison (trimmed of all connective fascia and silver-skin) and 35% firm wild boar backfat or domestic pork fatback²⁹. The fat must be chilled to −2°C before mincing to prevent fat smearing, which creates a hydrophobic coating that impedes salt penetration and moisture loss²⁹.

- Curing Ingredients (per kg of meat batter): 28.0 g common sea salt (NaCl)²⁹, 0.4 g potassium nitrate (saltpeter, KNO_3 , scraped from dry masonry cellars)²⁹, 5.0 g raw honey or dextrose²⁹, 3.0 g coarsely cracked black pepper²⁹, 2.0 g crushed juniper berries, 1.0 g minced garlic³³, and 10.0 mL dry red wine²⁹.
- Inoculation and Fermentation: Back-slopping with active brine from a previously successful raw sausage introduces wild strains of *Latilactobacillus sakei*, *Latilactobacillus curvatus*, and *Staphylococcus xylosus*²⁹. Over the first 48 to 72 hours at $18^\circ\text{C} - 20^\circ\text{C}$ and 85% relative humidity, the lactic acid bacteria ferment the sugars into lactic acid, dropping the pH from 5.8 down to 5.2²⁹. This drop reaches the isoelectric point of myosin ($pI = 5.3$), coagulating the muscle proteins into an irreversible cold-set gel that provides structural firmness²⁹.
- Nitrate Reduction and Color Development: Concurrently, *Staphylococcus xylosus* produces nitrate reductase, converting NO_3^- into nitrite (NO_2^-), which decomposes in the acidic batter into nitric oxide (NO) gas²⁹. The nitric oxide binds to the iron atom (Fe^{2+}) within muscle myoglobin, synthesizing nitrosylmyoglobin²⁹. This stabilizes a cured ruby-red color while inhibiting the outgrowth of *Clostridium botulinum* spores²⁹.
- Cold Smoking and Maturation: The sausages are hung in a cold smokehouse over smoldering embers of seasoned beech (*Fagus sylvatica*) and dried juniper branches at temperatures below 18°C for 3 to 5 days, depositing antimicrobial phenolic compounds on the casing²⁹. The sausages then dry in a curing cellar at $12^\circ\text{C} - 14^\circ\text{C}$ and 75% relative humidity for 4 to 8 weeks until they lose 35% to 40% of their initial weight, dropping the water activity (a_w) below 0.88, which guarantees shelf stability at room temperature²⁹.

Historical Heritage and Haute Chasse Traditions in Brandenburg-Prussia

The preparation of game for a civilised table draws upon centuries of culinary tradition in the Margraviate and Kingdom of Prussia. Far from utilitarian sustenance, the hunt (*die Jagd*) and the feast (*die Jagdtafel*) were expressions of territorial stewardship, governance, and culinary craftsmanship.

The Medieval Venison Pasty: The Architecture of the Pastry "Coffyn"

In the medieval courts of the Margraviate of Brandenburg under the Ascanian and early Hohenzollern dynasties, hunting large game was an exclusive right of the ruler (*Hohe Jagd*)⁵.

The centerpiece of medieval noble banquets was the venison pasty, designed to tenderize and preserve tough venison without mechanical refrigeration³⁷.

The outer crust, known as the "coffyn," was formulated not as an edible pastry, but as a durable cooking container³⁸. Made from a stiff dough of rye or whole wheat flour, boiling water, and suet, the pasty was hand-raised with thick walls (1.5 to 2.0 cm) and baked firm³⁸.

Tough venison shoulders and shanks were heavily seasoned with black pepper, cloves, mace, cubebs, and cinnamon, packed into the pastry box, and sealed under a generous layer of clarified venison or mutton suet³⁷.

As the pasty baked in a wood-fired masonry oven for 4 to 6 hours, the fat conducted heat evenly, melting the collagen cross-links into rich gelatin²⁴. Upon cooling, the solidified tallow created an airtight barrier, preventing oxygen exposure and bacterial spoilage, allowing the enclosed venison to keep safely in a cold pantry for months³⁸.

Renaissance Courtly Splendor: Joachim II and Marx Rumpolt

On March 7, 1542, Prince-Elector (*Kurfürst*) Joachim II Hector laid the foundation stone for Jagdschloss Grunewald (originally named *Haus zum grünen Wald*) on the shores of the Grunewaldsee, designed by master builder Caspar Theiss²¹. Grunewald served as the primary hunting lodge for courtly hunts across the pine and mixed oak woodlands southwest of Berlin²¹. The culinary treatment of game taken at Grunewald is documented in *Ein new Kochbuch* (published in Frankfurt am Main in 1581) by Marx Rumpolt, master cook (*Mundkoch*) to the Elector of Mainz⁴⁵. Rumpolt detailed elaborate techniques for wild game:

- Stags were partitioned into cuts, marinated in sour wine broths, larded with salt pork, and roasted on turning iron spits over beech wood while being basted with spiced vinegar, butter, and crushed gingerbread⁴⁵.
- Wild boar heads (*Sauköpfe*) were scalded, scraped clean, braised in spiced wine with rosemary, and served whole at banquets, coated in sweet-sour glazes made from wine must, honey, raisins, and pine nuts. Rumpolt's recipes bridged late medieval spicing and early modern balances of acidity, fat, and wild flavors⁴⁵.

The Baroque Pragmatism of the "Soldatenkönig" at Königs Wusterhausen

Under King Friedrich Wilhelm I of Prussia—the "Soldier King" (*Soldatenkönig*, reigning 1713–1740)—the royal court replaced elaborate foreign styles with a disciplined hunting tradition⁴⁸. The King selected Schloss Königs Wusterhausen, southeast of Berlin, as his autumn residence, living there from August through November to direct seasonal hunts⁴⁸.

Friedrich Wilhelm I maintained rigorous records of all game harvested across Brandenburg's state forests⁴⁸. The hunts at Königs Wusterhausen were demanding pursuits on horseback (*Parforcejagd*), for which the King laid out the radial hunting avenues of the *Parforceheide* centered on Jagdschloss Stern in Potsdam⁴⁸.

The harvested game was served at the daily meetings of the *Tabakskollegium* (the King's informal governing circle)⁴⁸. The evening meals featured roasted wild boar hams (*Wildschweinkeulen*), braised stag shoulders served with boiled cabbage and local turnips

(*Teltower Rübchen*), and cold smoked game sausages served with local beer.

Surplus meat was preserved and sent to military regiments in Berlin and Potsdam or sold through regulated municipal markets, establishing game as a practical nutritional resource in the Prussian economy⁴⁸.

Bourgeois Maturation: Henriette Davidis and Imperial Court Banquets

By the 19th and early 20th centuries, game cookery had become part of middle-class households across Berlin and the Mark Brandenburg, guided by cookbooks such as Henriette Davidis' *Praktisches Kochbuch für die gewöhnliche und feinere Küche* (first published in 1845)³⁵. Davidis simplified game preparation, moving away from heavy sour marinades toward precision roasting³⁵:

- Young roe saddles were trimmed of tough silver-skin, larded with fresh fatback, seasoned with salt, and roasted in butter, with the pan juices deglazed with sour cream (*Schmand*) and red currant jelly (*Johannisbeergelee*)³⁵.
- Under Kaiser Wilhelm II, the final imperial court hunts (*Hofjagden*) held at Königs Wusterhausen (ending in 1913) and the Schorfheide concluded with elaborate formal dinners⁴⁹.

These court banquets featured roasted saddles of fallow deer garnished with truffles, wild boar cutlets in port wine reductions, and terrines of hare served in porcelain, demonstrating the enduring role of Brandenburg game in refined regional cuisine⁴⁹.

Synthesis and Systemic Failure Analysis

Managing a sylvatic bio-resource system for a self-contained community requires maintaining a balance between ecological carrying capacity, processing hygiene, and culinary physics. Operational breakdowns in any of these areas can compromise the community's nutrition, health, or food security.

Primary Failure Modes and Field Detection

The three most critical failure modes involve post-mortem temperature management, wildlife population oversight, and zoonotic parasite control:

- **Deep-Tissue Bone Sour (*Knochenfäule*):** Occurs when a large carcass fails to chill below 10°C in the deep muscle core within 18 hours of slaughter²⁷. Body heat remains trapped around the femoral-pelvic joint, and bone dust from sawing introduces *Clostridium* and *Pseudomonas* into deep tissue²⁸. This is detected by a foul, sulfurous odor when cutting into the hip joint, along with a dull grayish discoloration of the *Adductor* and deep marrow²⁸. To correct this, the pelvic aitch bone (*Schloss*) must be removed during warm dressing, all bone dust scraped clean, and heavy haunches split along the seam to accelerate heat loss in the cooling cellar²⁷.
- **Ecological Commons Collapse:** Caused by uncoordinated, excessive hunting, harvesting lead sows (*Bachen*) and matriarchs, or failing to maintain wildlife refuges (*Wildschonbezirke*)¹³. This is detected by shrinking sounder sizes, a lack of juvenile cohorts, low body weights in yearlings, and the disappearance of active signs (wallows, rubs) across the territory. To correct this, driven hunts must be suspended immediately, a

complete 12-month hunting moratorium placed on half the territory, and hunting restricted solely to solitary males (*Keiler*).

- **Zoonotic Parasitic Infection:** Results from consuming undercooked wild boar infected with *Trichinella* larvae¹⁷, handling viscera contaminated with *Echinococcus* eggs, or feeding raw scrap meat to domestic pigs¹⁹. Infected individuals present with gastroenteritis, swollen eyelids, severe muscle pain, and persistent fever within 1 to 3 weeks of eating game¹⁷. To prevent this, 100% artificial digestion testing or compression squash microscopy must be enforced on all suids¹⁷, all wild boar meat cooked to an internal core temperature of at least 71°C ¹⁹, and all suspect viscera buried or incinerated.

Failure Mode	Root Mechanisms	Field Diagnostic Indicators	Operational Corrective Action
1. Deep-Tissue "Bone Sour" (<i>Knochenfäule</i>)	Failure to reduce deep core carcass temperature below 10°C within 18 hours ²⁷ ; heat trapping around the femoral-pelvic joint; saw dust seeding marrow with <i>Clostridium</i> and <i>Pseudomonas</i> ²⁸ .	Unpleasant, sulfurous putrefactive odor upon slicing the deep hip joint; grayish discoloration of the <i>Adductor</i> and bone marrow ²⁸ ; high surface slime.	Debone the aitch bone (<i>Schloss</i>) immediately upon warm dressing; scrape away all bone dust with a clean knife; split the heavy round primals to accelerate heat dissipation in the cooling cellar ²⁷ .
2. Ecological Commons Collapse	Uncoordinated over-harvesting; taking reproductive sows (<i>Bachen</i>) and alpha matriarchs; failure to enforce rotational sanctuaries (<i>Wildschonbezirke</i>) ¹³ .	Sharp reduction in sounder group sizes; rising proportion of yearlings showing low body weight; absence of mature sign (rubbing posts, wallows) across the territory.	Suspend all driven hunts immediately; enforce a complete 12-month hunting ban across 50% of the territory; limit harvesting to solitary males (<i>Keiler</i>) and strictly protect social sounders.
3. Zoonotic /	Ingestion of	Patients present	Require 100%

Parasitic Intoxication	undercooked wild boar infected with <i>Trichinella</i> larvae ¹⁷ ; handling offal carrying <i>Echinococcus</i> ova; feeding raw scraps to domestic pigs ¹⁹ .	with acute gastroenteritis, periorbital eyelid edema, severe muscle pain, and high fever 1 to 3 weeks post-consumption ¹⁷ .	artificial digestion testing or compression squash microscopy on all suids ¹⁷ ; mandate that all boar and bear meats reach an internal core temperature of > 19 ; bury all infected viscera.
-------------------------------	--	--	---

Operational Safety Hazards

Personnel engaged in wild game harvesting, butchery, and preservation face distinct occupational risks:

- Mechanical and Edge Trauma:** Using sharp carbon-steel cleavers, boning knives, and saws in cold, wet work areas presents laceration hazards²⁸. Handlers should wear protective split-leather or wire-mesh gloves on the non-dominant guide hand²⁸. Approaching entangled or wounded boars and stags in nets or traps carries severe risks of puncture wounds or goring; dispatch thrusts must be made from secure angles using long boar spears equipped with cross-guards.
- Biological and Zoonotic Exposure:** Direct contact with infected viscera presents transmission risks for sylvatic pathogens including tularemia (*Francisella tularensis*, contracted while skinning hares), brucellosis, and leptospirosis. Any cuts or punctures sustained during field dressing must be cleaned with antiseptic solutions, cauterized, or dressed with iodine.
- Asphyxiation and Toxic Fumes:** Operating wood-fired smokehouses and deep subterranean cellars presents gas inhalation risks. Enclosed smoking chambers concentrate lethal carbon monoxide (CO) and irritant combustion gases; chambers must be aired out before entry. Similarly, deep earth cellars used for hanging meat can accumulate heavy pools of carbon dioxide (CO_2) generated by microbial respiration and carcass degassing. Deep cellars must be checked using an open-flame candle test before personnel descend into the space.

Works cited

- 137Cs monitoring in the meat of wild boar population in Slovakia, https://www.researchgate.net/publication/303947363_137Cs_monitoring_in_the_meat_of_wild_boar_population_in_Slovakia

2. Long-term trends of hunting bags and wildlife populations in Central,
https://www.researchgate.net/publication/312072090_Long-term_trends_of_hunting_bags_and_wildlife_populations_in_Central_Europe
3. Jagdstrecke des Jagdjahres 2018/2019 im Land Brandenburg,
<https://ljv-brandenburg.de/jagdstrecke-des-jagdjahres-20182019-im-land-brandenburg/>
4. Jagdstrecken - MLEUV - Landesregierung Brandenburg,
<https://mleuv.brandenburg.de/mleuv/de/umwelt/jagd/jagdstatistik/jagdstrecken/>
5. Jagdreisen Brandenburg ▷ Deutschlands günstigste Jagdreise,
<https://jagdreisen-brandenburg.com/>
6. Jagdstrecke 2024/25 - MLEUV - Landesregierung Brandenburg,
<https://mleuv.brandenburg.de/mleuv/de/aktuelles/presseinformationen/detail/~04-12-2025-jagdstrecke-202425>
7. Jagdzeiten Brandenburg - Das Jagdsript - Jagdschein Vorbereitung,
<https://jagdschein-vorbereitung.de/pruefungswissen-online/tabellen/jagdzeiten/jagdzeiten-brandenburg/>
8. Jagdstatistik | MLEUV - Land Brandenburg,
<https://mleuv.brandenburg.de/mleuv/de/umwelt/jagd/jagdstatistik/>
9. Brandenburg zählt Rückgang bei Wolfsfamilien und Territorien - WELT,
<https://www.welt.de/regionales/berlin/article6914c7c635739e197250909a/brandenburg-zaehlt-rueckgang-bei-wolfsfamilien-und-territorien.html>
10. Wölfe in Brandenburg: Konkreter Plan fordert Abschuss von über,
<https://www.jaegermagazin.de/jagd-aktuell/woelfe-in-brandenburg-konkreter-plan-fordert-abschuss-von-ueber-300-tieren/>
11. Wölfe - beim NABU Brandenburg,
<https://brandenburg.nabu.de/tiere-und-pflanzen/saeugetiere/wolf/index.html>
12. Masterthesis „Wölfe in Brandenburg“,
https://digibib.hs-nb.de/file/dbhsnb_thesis_0000002843/dbhsnb_derivate_0000003445/Masterarbeit-Ahrens-2022.pdf
13. Wildtierbestände im Sinkflug – Warnungen der Jägerschaft werden,
<https://ljv-brandenburg.de/wildtierbestaende-im-sinkflug-warnungen-der-jaegerschaft-werden-traurige-realitaet/>
14. Reduction in the Radiocesium in Meats of the Sika Deer and Wild,
https://www.researchgate.net/publication/339315110_Reduction_in_the_Radiocesium_in_Meats_of_the_Sika_Deer_and_Wild_Boar_by_Cooking
15. Gemeinsame Richtlinie für die Hege und Bejagung des ... - BRAVORS,
<https://bravors.brandenburg.de/de/verwaltungsvorschriften-216896>
16. (PDF) Estimating wild boar (Sus scrofa) density using camera traps,
https://www.researchgate.net/publication/266333898_Estimating_wild_boar_Sus_scrofa_density_using_camera_traps_and_distance_sampling
17. Entnahme von Proben zur Untersuchung auf Trichinen,
<https://fortbildung-lazbw.lgl-bw.de/lazbw/webbasys/download/Shop/Entnahme%20von%20Proben%20beim%20Wildschwein.pdf>
18. Entsetzen über Jagdstatistik: „Brandenburg verliert seine großen,
<https://www.tagesspiegel.de/potsdam/brandenburg/entsetzen-uber-jagdstatistik>

[-brandenburg-verliert-seine-grossen-wildarten-und-die-politik-schaut-zu-15023234.html](#)

19. 52 Swine Slaughter and Processing Steps.pdf
20. 45 Bovine Humane Dispatch Methods.pdf
21. Geheimnisse des Jagdschlusses Grunewald - Blog | SPSG,
<https://www.spsg.de/blog/article/2022/03/04/geheimnisse-des-jagdschlusses-grunewald>
22. JAGDSCHLOSS GRUNEWALD,
https://www.spsg.de/fileadmin/user_upload/pdf/Schloesser/SPSG_Flyer_Jagdschlöss-Grunewald.pdf
23. Jagdschloß Grunewald - Denkmaldatenbank - Berlin.de,
https://denkmaldatenbank.berlin.de/daobj.php?obj_dok_nr=09075448
24. 48 Beef Cuts And Cooking Methods.pdf
25. Grundwissen Jägerprüfung - Das Standardwerk zum Jagdschein,
<https://dokumen.pub/grundwissen-jagerprufung-das-standardwerk-zum-jagdschein.html>
26. Jagdzeiten - Landesjagdverband Brandenburg e.V.,
<https://ljv-brandenburg.de/jagdzeiten/>
27. 46 Bovine Carcass Dressing Guide.pdf
28. 47 Bovine Manual Butchery Dissection Guide.pdf
29. 111 Traditional Salami Formulation And Curing.pdf
30. Merkblatt zur Probenahme für die Trichinenuntersuchung bei Wild,
https://www.landkreis-prignitz.de/globalcontent/documents/veterinaer_ordnung/veterinaer/Merkblatt_Trichinenprobenahme_Wild_3_2020.pdf
31. Radiocesium concentrations in wild boars captured within 20 km of,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7283223/>
32. Selected aspects of caesium-137 contamination in wild boars (Sus,
<https://www.scilit.com/publications/d3acacfc657ee9ec9f45d89671c3d392>
33. 53 Pork Primal Cuts and Curing.pdf
34. Potassium Bioaccessibility in Uncooked and Cooked Plant Foods,
<https://www.mdpi.com/2072-6651/14/10/668>
35. Praktisches Kochbuch für die gewöhnliche und feinere Küche,
<https://de.everand.com/book/292446128/Praktisches-Kochbuch-fur-die-gewohnliche-und-feinere-Kuche-Mit-besonderer-Berucksichtigung-der-Anfangerinnen-und-angehenden-Ein-Klassiker-der-deuts>
36. Rheingauer Hühner - Ein Rezept aus dem Mittelalter - Elmar Lorey,
<https://www.elmar-lorey.de/rheingau/rezhuhn.htm>
37. Kraut und Hexe, Magier - Tempora Historica Mittelalter und Gothic,
<https://www.temporahistorica.de/online-bestellen-webshop/literatur-buch-und-musik/karfunkel-k%C3%BCche/>
38. The Cornish Pasty presents English pasties,
<https://www.cornishpasties.org.uk/otherpasties/english-pasties.htm>
39. Knead to Know A History of Baking 9781837731213, 9781837731220,
<https://dokumen.pub/knead-to-know-a-history-of-baking-9781837731213-9781837731220.html>

40. Uncategorized - UK PieFest, <https://www.ukpiefest.co.uk/?cat=1>
41. History and Types of Pies Explained | PDF | European Cuisine, <https://www.scribd.com/document/975258247/pie-wiki>
42. Medieval Recipes for Modern Cooks | PDF | Roasting | Breads - Scribd, <https://www.scribd.com/document/831942526/Pleyn-Delit-Medieval-Cookery-for-Modern-Cooks-Sharon-Butler-Constance-B-Hieatt-Etc-Z-Library>
43. Jagdschloss Grunewald, <https://www.freunde-psg.de/schloesser-und-gaerten/schloss-3>
44. Jagdschloss Grunewald - visitBerlin.de, <https://www.visitberlin.de/de/jagdschloss-grunewald>
45. Der Waldrapp. Historische Quellen - Version 1/1 - manuscripta.at, https://manuscripta.at/Ma-zu-Bu/wr/Roland-Martin_Waldrapp_Historische-Quellen_Version-2-1_2024.pdf
46. <https://sd458c9149fd0c72b.jimcontent.com/download/version/1401805352/module/9668720922/name/Renaissance.pdf>
47. Topografie der Gemengsel und Gehäcksel - Dieter Froelich, <http://www.dieterfroelich.de/restauration/topografie.pdf>
48. Friedrich-Wilhelm I. – der Soldatenkönig und seine Schlösser, <https://havelblog.de/friedrich-wilhelm-i-der-soldatenkoenig-und-seine-schloesse/r/>
49. Schloss Königs Wusterhausen, <https://www.koenigs-wusterhausen.de/schloss-koenigs-wusterhausen-a5fbd22575d82cd1>
50. Schloss Königs Wusterhausen, <https://kulturelle-bildung-brandenburg.de/projekte/schloss-koenigs-wusterhausen/>
51. SCHLOSS KÖNIGS WUSTERHAUSEN, https://www.spsg.de/fileadmin/user_upload/pdf/Schloesser/SPSG_Flyer_Schloss-Koenigs-Wusterhausen.pdf
52. Schloss Königs Wusterhausen Kavalierhäuser, <https://jungekuenstlerstellensichvor.com/schloss-koenigswusterhausen-kavalierhauser/>