BI-02 The two endocrine systems compared

Introduction: A Tripartite Framework for Endocrine Analysis

The biological foundations of human coexistence—how men and women interact, pair-bond, parent, and structure societies—are inextricably linked to the endocrine system. The endocrine network operates as a primary biological interface, translating environmental stimuli into physiological adaptations and shaping statistical predispositions in behaviour. However, discussions surrounding sex differences in endocrinology frequently succumb to either strict biological determinism or absolute social constructivism. To navigate this complex domain with scientific rigor, this analysis employs a tripartite framework, separating the phenomena into three interacting layers:

1. **Physiology (The Measurable Substrate):** The foundational biochemical and anatomical mechanisms, encompassing hormone concentrations, enzymatic pathways, and receptor distributions. At this layer, the male and female endocrine architectures are dimorphic and fundamentally distinct, characterized by non-overlapping temporal architectures and metabolic baselines.
2. **Behaviour (The Statistical Distribution):** The psychological and behavioural outcomes modulated by neuroendocrine signalling. At this layer, sex differences manifest as overlapping statistical distributions (Gaussian curves). While group averages differ—often yielding quantifiable effect sizes—the overlap dictates that the behavioural phenotype of an individual cannot be reliably deduced solely from their biological sex or hormonal profile.
3. **Culture (The Variable Environment):** The ecological, social, and cultural contexts that exert bidirectional influence on the endocrine system. Hormones do not operate in a vacuum; social scripts, hierarchies, and environmental stressors act as epigenetic and physiological modulators that upregulate or downregulate endocrine cascades.

This report comprehensively evaluates the male and female endocrine systems in adulthood, focusing on eight principal hormones: testosterone, estradiol, progesterone, oxytocin, vasopressin, prolactin, cortisol, and dehydroepiandrosterone (DHEA). It prioritizes peer-reviewed primary literature, systematic reviews, and meta-analyses to examine their concentrations, rhythms, receptor topography, and responses to physical and psychosocial stressors.

1. The Physiological Layer: Mechanisms, Axes, and Receptors

The human endocrine system is governed by complex feedback loops regulated primarily by the hypothalamus and the pituitary gland. Two main axes are paramount to this comparative analysis: the Hypothalamic-Pituitary-Gonadal (HPG) axis and the

Hypothalamic-Pituitary-Adrenal (HPA) axis.

The **HPG axis** is the central regulator of reproduction. In both sexes, the hypothalamus releases Gonadotropin-Releasing Hormone (GnRH) in a pulsatile fashion into the hypophyseal portal system. This stimulates the anterior pituitary to secrete Luteinizing Hormone (LH) and Follicle-Stimulating Hormone (FSH). In males, LH drives the Leydig cells of the testes to synthesize testosterone, while FSH targets Sertoli cells to regulate spermatogenesis. In females, LH and FSH orchestrate the ovarian synthesis of estradiol and progesterone, driving the menstrual cycle.

The **HPA axis** is the primary mediator of the physiological stress response. The hypothalamus releases Corticotropin-Releasing Hormone (CRH), stimulating the anterior pituitary to secrete Adrenocorticotropic Hormone (ACTH). ACTH travels systemically to the adrenal cortex, stimulating the release of glucocorticoids (primarily cortisol) and adrenal androgens (such as DHEA)[1].

These axes engage in profound crosstalk. During acute or chronic stress, hyperactivation of the HPA axis exerts an inhibitory effect on the HPG axis. Elevated glucocorticoids suppress GnRH pulsatility at the hypothalamus, decrease pituitary responsiveness to GnRH, and directly inhibit gonadal steroidogenesis. This mechanism evolved to prioritize immediate survival over reproduction during periods of environmental threat[3].

1.1 The Eight Primary Hormones: Concentrations and Neural Topography

Understanding the endocrine influence on behaviour requires mapping not just the circulating concentrations of hormones, but their specific receptor distributions within the central nervous system. Table 1 outlines the typical adult reference ranges, followed by a detailed analysis of each hormone's mechanisms.

Table 1: Typical Adult Serum/Plasma Concentrations by Sex

Hormone	Typical Adult Male Range	Typical Adult Female Range	Primary Source
Testosterone (Total)	10.4 – 34.7 nmol/L (300–1000 ng/dL)	0.5 – 2.4 nmol/L (15–70 ng/dL)	Gonads (Testes/Ovaries), Adrenals
Estradiol (E2)	36.7 – 146.8 pmol/L (10–40 pg/mL)	55 – 1468 pmol/L (15–400 pg/mL)*	Gonads, peripheral aromatization
Progesterone	< 3.18 nmol/L (< 1.0 ng/mL)	0.3 – 63.6 nmol/L (0.1–20 ng/mL)*	Corpus Luteum, Adrenals
DHEA-S	80 – 560 µg/dL	35 – 430 µg/dL	Adrenal Cortex (Zona Reticularis)

Cortisol (Total)	138 – 690 nmol/L (5–25 µg/dL)	138 – 690 nmol/L (5–25 µg/dL)	Adrenal Cortex (Zona Fasciculata)
Prolactin	< 20 ng/mL (Average: 5 ng/mL)	< 25 ng/mL (Average: 13 ng/mL)	Anterior Pituitary (Lactotrophs)
Oxytocin (Plasma)	1.53 ± 1.19 pg/mL**	4.53 ± 1.18 pg/mL**	Hypothalamus, Posterior Pituitary
Vasopressin (Plasma)	1.0 – 5.0 pg/mL	1.0 – 5.0 pg/mL	Hypothalamus, Posterior Pituitary

** Female ranges for estradiol and progesterone fluctuate drastically across the follicular and luteal phases of the menstrual cycle.* Plasma oxytocin measurements are highly assay-dependent and actively debated in the literature. Some studies, however, indicate significantly higher baseline averages in females[5].

1.1.1 The Androgens: Testosterone and DHEA

Testosterone is the primary male sex steroid, though it is present and vital in females. It exerts its effects via the androgen receptor (AR). In the brain, ARs are densely distributed in the hypothalamus, the amygdala, and the hippocampus. Beyond driving male secondary sexual characteristics and skeletal muscle hypertrophy, centrally acting testosterone modulates neural circuits associated with status-seeking, territoriality, and threat vigilance.

DHEA (Dehydroepiandrosterone) and its sulfated derivative **DHEA-S** are the most abundant circulating steroid hormones in the human body. Synthesized primarily in the zona reticularis of the adrenal cortex, they serve as crucial precursors for both peripheral androgens and estrogens. In the brain, DHEA operates as an independent neurosteroid. It modulates neuroplasticity and acts as an antagonist at GABA-A receptors while potentiating NMDA receptors, contributing to resilience against chronic stress and depressive symptomatology.

1.1.2 The Estrogens and Progestogens: Estradiol and Progesterone

Estradiol, the most potent endogenous estrogen, binds to estrogen receptors (ER-alpha and ER-beta). These receptors are highly concentrated in the hypothalamus, the amygdala, and the prefrontal cortex. In females, estradiol drives reproductive maturation and ovulation, maintains bone mineral density, and strongly modulates serotonergic, dopaminergic, and glutamatergic tone. In males, circulating estradiol—derived primarily from the peripheral aromatization of testosterone—is equally critical for maintaining bone density, vascular health, and libido[8].

Progesterone is essential for the preparation and maintenance of pregnancy, acting through the progesterone receptor (PR). In the central nervous system, progesterone is rapidly metabolized into allopregnanolone. Allopregnanolone is a highly potent positive allosteric modulator of the GABA-A receptor. It exerts profound anxiolytic, sedative, and anticonvulsant

effects. In vulnerable women, rapid fluctuations in allopregnanolone levels prior to menstruation trigger an abnormal central nervous system response, leading to Severe Premenstrual Dysphoric Disorder (PMDD), a condition hypothesized to stem from aberrant GABA-A receptor sensitivity rather than abnormal absolute hormone concentrations, as investigated by Bäckström and Epperson[9].

1.1.3 The Nonapeptides: Oxytocin and Vasopressin

Synthesized primarily in the paraventricular (PVN) and supraoptic nuclei of the hypothalamus, these two structurally similar nonapeptides project both centrally into the brain and peripherally via the posterior pituitary gland.

Oxytocin acts via the oxytocin receptor (OXTR), which is densely distributed in the amygdala, the striatum, the ventromedial hypothalamus, and the nucleus accumbens. Peripherally, oxytocin induces uterine contractions during parturition and milk ejection during nursing. Centrally, it is heavily implicated in social memory, the modulation of fear and anxiety in the amygdala, and the reinforcement of affiliative pair-bonding and maternal care[11].

Vasopressin (AVP) acts centrally via the V1a receptor, which shows prominent expression in the lateral septum and amygdala. While its well-known peripheral function is the endocrine regulation of renal water excretion (antidiuretic hormone) and blood pressure, its central actions modulate social recognition, territoriality, male-typical pair-bonding behaviours, and reactive aggression[11].

1.1.4 Stress and Lactation: Cortisol and Prolactin

Cortisol acts via glucocorticoid receptors (GR) and mineralocorticoid receptors (MR), which are distributed universally throughout the body. High densities of GRs and MRs in the hippocampus facilitate the negative feedback loop that shuts down the HPA axis following a stressor.

Prolactin is a polypeptide hormone synthesized by lactotrophs in the anterior pituitary. Its receptors (PRLR) are found in the mammary glands and heavily throughout the brain, notably in the choroid plexus. While its primary physiological role is to induce breast enlargement and stimulate postpartum lactation, it also functions as a neuropeptide that modulates the stress response and regulates maternal emotionality. Baseline prolactin levels average 13 ng/mL in women and 5 ng/mL in men, with hyperprolactinemia (often drug-induced via dopamine D2 antagonists) capable of causing profound hypogonadism in both sexes[7].

2. Two Fundamentally Different Temporal Architectures

The most profound physiological distinction between the male and female endocrine systems is not merely the absolute concentration of circulating hormones, but the temporal architecture governing their release.

2.1 The Male Circadian Paradigm

The adult male HPG axis operates on a stable, 24-hour circadian rhythm. Testosterone secretion is tightly coupled to sleep architecture, specifically REM sleep. In healthy young men, testosterone levels peak in the early morning (typically between 06:00 and 08:00) and steadily decline throughout the waking day, reaching their lowest point in the late evening[14].

This daily architectural reset provides a reliable, predictable, and tonic level of androgens. The male central nervous system is bathed in a relatively stable hormonal milieu day after day, without programmatic, wide-scale fluctuations. Consequently, male receptor sensitivities do not require the constant, drastic recalibration necessary in the female system.

2.2 The Female Circatrigintan (Menstrual) Paradigm

In stark contrast, the adult female HPG axis operates on a ~28-day circatrigintan cycle, characterized by massive, programmatic shifts in estradiol and progesterone.

During the follicular phase, rising FSH stimulates ovarian follicles, which begin secreting increasing amounts of estradiol. This gradual rise culminates in a massive estradiol peak, which triggers a positive feedback loop resulting in the LH surge, causing ovulation. Following ovulation, the system enters the luteal phase, where the newly formed corpus luteum secretes enormous quantities of progesterone alongside estradiol. If fertilization does not occur, the corpus luteum undergoes apoptosis, leading to a precipitous drop in both hormones, triggering menstruation.

This temporal architecture dictates that the female brain is exposed to a continuously shifting ratio of neuroactive steroids. The brain must constantly adapt to the varying serotonergic and dopaminergic tone dictated by estradiol, as well as the profound GABAergic fluctuations dictated by progesterone and allopregnanolone. The female endocrine system is inherently dynamic, requiring massive physiological flexibility.

2.3 Rhythms Across the Lifespan: Adrenopause, Andropause, and Menopause

Both sexes experience age-related hormonal declines, but the trajectories vary significantly by hormone and by sex.

The decline of adrenal androgens, specifically **DHEA and DHEA-S**, is linear and steep. DHEA levels peak in the early to mid-20s and subsequently drop by approximately 2% to 3% per year in both men and women. By age 70 to 80, individuals possess only 10% to 20% of their youthful peak levels, a phenomenon termed "adrenopause." Because cortisol production generally holds steady across the lifespan, the cortisol-to-DHEA ratio shifts dramatically with age, which researchers hypothesize contributes to age-related frailty and cognitive decline, though DHEA supplementation trials have yielded mixed results[2].

In women, ovarian function ceases abruptly around age 50, a discrete event known as **menopause**. The cessation of follicular depletion results in a rapid, severe decline in both estradiol and progesterone, initiating systemic changes in bone metabolism, cardiovascular risk profiles, and neuroendocrinology.

In men, the decline of testosterone—often colloquially termed **andropause** or Late-Onset Hypogonadism (LOH)—is not an abrupt cessation but a gradual, inexorable drift. The exact parameters of this decline were clarified by the landmark European Male Ageing Study (EMAS), a prospective cohort study conducted by Wu et al. (2010) and published in the *Journal of Clinical Endocrinology & Metabolism*. The study tracked 3,369 men aged 40–79 across eight European countries. EMAS revealed that total testosterone declines by an average of -0.04 nmol/L per year, and free testosterone by -3.12 pmol/L per year[18].

Crucially, however, the EMAS data demonstrated that this decline is not strictly a function of

chronological aging. It is heavily confounded by modifiable risk factors, particularly rising body mass index (BMI) and comorbid illnesses. The study found that obesity decreases total and free testosterone significantly more than age alone; obese men experienced reductions in free testosterone of -53.72 pmol/L compared to non-obese men. Healthy, non-obese aging men experience a much shallower decline in androgens, challenging the notion that severe testosterone deficiency is an inevitable consequence of male aging[19].

3. Endocrine Responses to Environmental Stressors

The endocrine system is not a closed loop; it is highly responsive to the external environment. Examining how the male and female systems respond to stress, sleep loss, and energy deficits reveals further sexually dimorphic traits.

3.1 Psychological Stress: The Trier Social Stress Test (TSST)

The Trier Social Stress Test (TSST), originally developed by Kirschbaum et al. (1993), is the gold standard protocol for inducing acute psychological stress in laboratory settings. It combines social-evaluative threat and uncontrollability, requiring participants to deliver a speech and perform mental arithmetic in front of a stoic, non-responsive panel. This protocol reliably activates the HPA axis, resulting in cortisol secretion[22].

A rigorous meta-analysis by Liu et al. (2017) in *Psychoneuroendocrinology* aggregated findings from 34 TSST studies comprising 1,350 individuals (640 women and 710 men). Using a random-effects model, the analysis found significant and replicable sex differences: men exhibited stronger, higher salivary cortisol values at peak reactivity and during the recovery phase compared to women[1].

The literature suggests these physiological differences may reflect divergent evolutionary coping strategies. Males exhibit greater physiological reactivity to achievement-based and status-threatening stressors (the traditional "fight-or-flight" response). In contrast, females may employ more socially oriented "tend-and-befriend" coping mechanisms, mediated in part by oxytocin, which attenuates the HPA axis response and reduces absolute cortisol output during social-evaluative stress[1].

3.2 Sleep Loss and Circadian Disruption

Because the male testosterone rhythm is intricately tied to sleep architecture, sleep deprivation acts as a severe endocrine disruptor. A controlled clinical trial conducted by Leproult and Van Cauter (2011) and published in *JAMA* investigated this phenomenon. Ten young, healthy men underwent one week of sleep restriction, limited to 5 hours of sleep per night.

The results were striking: one week of sleep restriction reduced daytime testosterone levels by 10% to 15%. To put this effect size into perspective, a 15% reduction in testosterone is equivalent to aging 10 to 15 years[25]. Sleep loss also elevated evening cortisol levels, disrupting the normal circadian decline of the HPA axis.

Furthermore, prolactin secretion, which is highly sensitive to the circadian rhythm and peaks roughly 4 hours after sleep onset, is deeply disrupted by sleep fragmentation or shift work. Chronic circadian desynchrony impairs reproductive capacity in both sexes, underscoring the absolute necessity of sleep architecture for baseline endocrine homeostasis[15].

3.3 Diet and Exercise: Relative Energy Deficiency in Sport (RED-S)

When physical energy expenditure chronically exceeds dietary energy intake, a state of Low Energy Availability (LEA) occurs. The brain detects this caloric deficit as starvation and initiates a physiological triage, suppressing non-essential physiological functions—most notably the HPG axis—to conserve energy for survival. Historically termed the "Female Athlete Triad," the International Olympic Committee (IOC) updated the framework in 2014 to "Relative Energy Deficiency in Sport" (RED-S) to acknowledge its multi-systemic consequences in both sexes[29]. The threshold for HPG disruption, however, differs significantly by sex.

Table 2: Estimated Thresholds for HPG Axis Disruption in RED-S

Sex	Energy Availability Threshold	Clinical Endocrine Consequence
Female	< 30 kcal/kg Fat-Free Mass (FFM) / day	Functional Hypothalamic Amenorrhea (FHA), profoundly low estradiol.
Male	< 25 kcal/kg Fat-Free Mass (FFM) / day	Functional Hypogonadotropic Hypogonadism, clinically low testosterone.

In females, dropping below 30 kcal/kg FFM per day rapidly inhibits GnRH pulsatility. This halts the menstrual cycle (amenorrhea) and causes a crash in circulating estradiol, which subsequently leads to rapid losses in bone mineral density and an increased risk of stress fractures. Males appear slightly more resilient to short-term caloric deficits, with the disruption threshold estimated closer to 25 kcal/kg FFM. However, when male endurance athletes cross this threshold, they experience severe secondary hypogonadism, with clinically low testosterone, reduced libido, elevated cortisol, and impaired musculoskeletal recovery[29].

4. Behavioural Modulations: Overlapping Distributions and Controversies

Transitioning to the second layer—behaviour—requires shifting from the binary absolutes of physiology to the statistical reality of overlapping normal distributions. While hormones bias neurocognitive networks toward specific behaviours, human cortical development allows for immense cognitive override. Consequently, no single hormone dictates behaviour in isolation.

4.1 Testosterone, Aggression, and the Dual-Hormone Hypothesis

The relationship between testosterone and human aggression is highly contested. In many non-human species, the correlation is robust and absolute; in humans, it is subtle and mediated by complex social factors.

Archer (2006) proposed the "Challenge Hypothesis" for humans, arguing that testosterone

levels rise during status-relevant conflicts and drive competitive, dominant behaviours[33]. A comprehensive meta-analysis by Geniole et al. (2020), published in *Hormones and Behavior*, aggregated 45 independent studies involving 9,760 participants to evaluate this link. The analysis found a statistically significant but weak positive correlation between baseline testosterone and aggression, with a mean weighted effect size of $r = 0.14$. This effect was found to be slightly stronger in men ($r = 0.071$) than in women ($r = 0.002$)[35].

To explain this remarkably weak correlation, Mehta and Josephs (2010) proposed the **Dual-Hormone Hypothesis**. They posited that testosterone only promotes status-seeking and aggression when cortisol levels are concurrently low. If cortisol is high—indicating acute psychological stress or anxiety—the behavioural effects of testosterone are entirely blocked[37].

However, the Dual-Hormone Hypothesis itself has recently faced intense scrutiny. A highly rigorous meta-analysis by Dekkers et al. (2019) in *Psychoneuroendocrinology* reviewed 33 studies containing 49 effect sizes and 8,538 participants. The analysis yielded only marginal support for the hypothesis: the interaction effect between testosterone and cortisol on status-relevant behaviour was statistically significant but practically minuscule ($r = -0.061$)[37].

The current consensus suggests that while testosterone plays a role in status-seeking, territoriality, and threat vigilance, actual human physical aggression is overwhelmingly mediated by prefrontal cortical control, trait impulsivity, and socio-economic variables, rather than mere androgen concentrations.

4.2 Testosterone, Fatherhood, and Social Context

Human males exhibit a profound degree of endocrine neuroplasticity in response to familial roles. While high testosterone facilitates mating effort, it is biologically antagonistic to the empathetic, nurturing demands of parenting effort.

In a landmark longitudinal study published in the *Proceedings of the National Academy of Sciences (PNAS)*, Gettler et al. (2011) tracked 465 men in the Philippines over 4.5 years. The study found that single men with higher baseline testosterone were significantly more likely to partner and become fathers. However, upon transitioning to fatherhood, these men experienced massive, statistically significant drops in their testosterone levels.

Men who became partnered fathers experienced a median decline in waking testosterone of 26%, and a median decline in evening testosterone of 34%—declines far steeper than the standard age-related drift observed in non-fathers of the same cohort[40]. Furthermore, the study demonstrated a dose-response relationship with caregiving: fathers who spent 3 or more hours a day actively engaging in childcare exhibited the lowest testosterone levels at follow-up[40].

This replicated finding demonstrates that the male endocrine system is highly responsive to social context. Male physiology adaptively downregulates competitive mating drives in favour of affiliative parenting drives when placed in a stable familial environment.

4.3 Oxytocin, Trust, and The Crisis of Replication

In 2005, a highly publicized study by Kosfeld et al. claimed that the intranasal administration of oxytocin significantly increased interpersonal trust in economic games. This ignited a decade of enthusiastic research branding oxytocin as the "cuddle hormone" and proposing it as a treatment for autism and social anxiety[43].

Today, the field of intranasal oxytocin research is embroiled in a severe replication crisis. In a devastating critique published in *Biological Psychiatry*, Leng and Ludwig (2016) argued that the premise of intranasal administration is fundamentally flawed. They demonstrated that very little, if any, intranasal oxytocin crosses the blood-brain barrier to reach the cerebrospinal fluid in pharmacologically relevant concentrations. They posited that any observed behavioural effects were likely due to peripheral actions or statistical noise[45].

This critique was corroborated by Walum et al. (2016) in the same journal, who demonstrated that the vast majority of intranasal oxytocin behavioural studies were severely underpowered (with an average statistical power around 12%). This low power, combined with publication bias and HARKing (Hypothesizing After Results are Known), led to a massive inflation of false-positive results in the literature[43]. While endogenous oxytocin unquestionably modulates social bonding and threat perception within the amygdala, the idea that a nasal spray can reliably alter complex human behaviours is currently viewed with profound skepticism by rigorous neuroendocrinologists[45].

Conversely, research investigating genetic variations in the endogenous oxytocin receptor gene (OXTR) yields more reliable, albeit modest, data. The most thoroughly investigated Single Nucleotide Polymorphism (SNP) is rs53576, involving an adenine (A) to guanine (G) transition. A comprehensive meta-analysis by Bakermans-Kranenburg and van IJzendoorn (N=4,955) found that individuals homozygous for the G allele (GG) exhibit slightly higher "general sociality" and self-reported empathy compared to A-allele carriers. However, the effect size is very small (Cohen's $d = 0.11$)[47]. Furthermore, Tost et al. (2010) in *PNAS* found that risk allele carriers (A) show structural neuroanatomical differences, including decreased hypothalamus volume, linking the variant to heightened stress reactivity and lower psychological resilience[48].

5. Distinguishing Averages from Individuals

A critical requirement in analyzing endocrine data is distinguishing between what is known about group averages and what can be said about specific individuals.

At the physiological layer, absolute hormone levels are heavily dimorphic. The average adult male possesses roughly 10 to 15 times more circulating testosterone than the average adult female, creating two distinct curves with virtually no overlap in healthy, non-pathological populations.

However, moving to the behavioural layer, the traits influenced by these hormones—such as physical aggression, spatial reasoning, empathy, or nurturing behaviour—form Gaussian distributions that overlap significantly between the sexes. If male physical aggression yields an effect size of $d = 0.5$ compared to females (a moderate effect in psychological research), it means that the average male is more aggressive than about 69% of females. Yet, this still leaves an immense overlap where hundreds of millions of women are more aggressive than hundreds of millions of men.

Therefore, while a specific hormone panel can identify a patient's biological sex with near-perfect accuracy, it holds virtually zero predictive validity regarding that individual's personality, moral character, or behavioural phenotype. The human prefrontal cortex exerts top-down executive control that can override subcortical, hormonally driven impulses.

6. The Cultural Layer: Biology is Not Destiny

The third layer of analysis is culture. The endocrine system does not dictate human social structures; rather, it interacts with them bidirectionally.

The longitudinal fatherhood studies (Gettler et al., 2011) and the data on acute stress reactivity provide unequivocal evidence that the environment dictates the hormonal baseline. A cultural environment or organizational structure that demands relentless male competition, sleep deprivation, and high-stakes social evaluation will chronologize high cortisol and fluctuating testosterone, driving hyper-vigilant and potentially aggressive behaviour. Conversely, a family structure or society that incentivizes active, hands-on fatherhood will literally alter the male endocrine architecture, suppressing testosterone and facilitating the neurological pathways required for patience, bonding, and child-rearing.

Culture, therefore, acts as an epigenetic architect. It determines which biological potentials within the male and female endocrine systems are activated, and which are suppressed.

Conclusion

Summary of Robust Findings

The male endocrine architecture is defined by a 24-hour circadian rhythm, maintaining tonic levels of testosterone that slowly decline with age and comorbidities, particularly obesity. The female architecture is defined by a 28-day circatrigintan cycle, characterized by massive, programmatic fluctuations in estradiol and progesterone that cease abruptly at menopause. Men experience significantly stronger and more prolonged peak cortisol responses to acute psychosocial stress (the TSST). Conversely, women are highly vulnerable to complete HPG-axis shutdown from dietary energy deficits (RED-S at <30 kcal/kg FFM). Testosterone shows a statistically reliable but remarkably weak correlation with human physical aggression (r = 0.14). The male endocrine system is highly neuroplastic in response to social roles; men transitioning into active fatherhood experience a precipitous, statistically massive drop in testosterone (up to 34%), facilitating child-rearing behaviors. Finally, early claims regarding intranasal oxytocin reliably increasing human trust have largely failed to replicate due to methodological flaws and the blood-brain barrier. Genetic variations in the oxytocin receptor (OXTR rs53576) do correlate with empathy and stress reactivity, though effect sizes remain small.

Contested or Open Questions

- **The Dual-Hormone Hypothesis:** Does cortisol truly moderate the relationship between testosterone and status-seeking behaviour? Recent large-scale meta-analyses (e.g., Dekkers et al., 2019) show the interaction effect is nearly zero, leaving the validity of the hypothesis highly debated.
- **Intranasal Neuropeptide Delivery:** Can oxytocin or vasopressin be effectively delivered to the central nervous system via the nasal cavity to treat social deficits (such as in autism spectrum disorder), or are all observed behavioural effects the result of peripheral action or placebo?
- **Andropause vs. Pathology:** Is the age-related decline in male testosterone a natural, inevitable artifact of chronological aging, or is it primarily a reversible symptom of modern

epidemics of obesity, sleep restriction, and metabolic syndrome?

- **Endogenous Oxytocin Baselines:** Do women truly possess higher baseline plasma oxytocin, or do current immunoassay methodologies fail to accurately capture the pulsatile, rapid-decay dynamics of peripheral and central neuropeptides in both sexes?

Essential Sources for Further Reading

1. **Gettler, L. T., et al. (2011).** *Longitudinal evidence that fatherhood decreases testosterone in human males.* Proceedings of the National Academy of Sciences (PNAS).
2. **Wu, F. C., et al. (2010).** *Hypothalamic-pituitary-testicular axis disruptions in older men are differentially linked to age and modifiable risk factors: the European Male Aging Study.* Journal of Clinical Endocrinology & Metabolism.
3. **Leng, G., & Ludwig, M. (2016).** *Intranasal Oxytocin: Myths and Delusions.* Biological Psychiatry.
4. **Liu, J. J. W., et al. (2017).** *Sex differences in salivary cortisol reactivity to the Trier Social Stress Test (TSST): A meta-analysis.* Psychoneuroendocrinology.
5. **Geniole, S. N., et al. (2020).** *Is testosterone linked to human aggression? A meta-analytic examination of the relationship between baseline, dynamic, and manipulated testosterone on human aggression.* Hormones and Behavior.
6. **Leproult, R., & Van Cauter, E. (2011).** *Effect of 1 week of sleep restriction on testosterone levels in young healthy men.* JAMA.
7. **Dekkers, T. J., et al. (2019).** *A meta-analytical evaluation of the dual-hormone hypothesis: Does cortisol moderate the relationship between testosterone and status, dominance, risk taking, aggression, and psychopathy?* Psychoneuroendocrinology.

Implications for Living Together

Applying endocrine research to the structuring of couples, families, and organizations offers practical utility, provided one avoids the naturalistic fallacy—the error of assuming that what is biologically "natural" is morally or socially "good." Biology describes what *is*, not what *ought* to be.

For couples and families, understanding the disparate temporal architectures of both sexes can foster profound interpersonal empathy. Recognizing that female mood, energy, and stress resilience fluctuate in tandem with intense, shifting ratios of estradiol and allopregnanolone across a 28-day cycle allows cyclical emotional shifts to be reframed as physiological realities rather than psychological failings. Conversely, recognizing that male testosterone production is deeply dependent on circadian sleep architecture highlights that chronic male irritability, low libido, or lethargy may be solved by prioritizing adequate sleep (treating the endocrine root) rather than treating it solely as a relational issue.

The data surrounding fatherhood holds immense implications for the structuring of families and societies. Biology explicitly rewards involved fatherhood. Because male testosterone downregulates in response to hands-on childcare—reducing competitive, mating-oriented drives and increasing affiliative capacities—societies and organizations that provide robust paternity leave and encourage fathers to be physically present with their infants are actively facilitating a biological shift that promotes familial stability and reduces male risk-taking.

Ultimately, biology establishes the boundaries of human potential, but culture and environment

dictate where within those boundaries we reside. By structuring organizations and family lives to reduce chronic stress (preventing HPA-axis hyperactivation) and support adequate sleep and nutrition, both men and women can optimize their endocrine health, fostering more cooperative, resilient, and cohesive communities.

Works cited

1. Sex differences in salivary cortisol reactivity to the Trier Social Stress, https://www.researchgate.net/publication/316444610_Sex_differences_in_salivary_cortisol_reactivity_to_the_Trier_Social_Stress_Test_TSST_A_meta-analysis
2. DHEA Supplement: Evidence, Studies & Why Doctors Are Cautious, https://acibademinternational.com/blog/dhea-supplements-what-the-hormone-does-who-the-studies-included-and-why-doctors-are-cautious/
3. Stress, hypothalamic-pituitary-adrenal axis ... - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC11535056/
4. Microecological modulators—new perspectives for treating sex, https://pmc.ncbi.nlm.nih.gov/articles/PMC13171582/
5. Mayo Clinic Biochemical Testing Handbook | PDF - Scribd, https://www.scribd.com/doc/122080381/Mayo-Clinic-Biochemical-testing-handbook
6. Sex-Related Differences in Plasma Oxytocin Levels in Humans, https://www.researchgate.net/publication/332056310_Sex-Related_Differences_in_Plasma_Oxytocin_Levels_in_Humans
7. Hyperprolactinemia - StatPearls - NCBI Bookshelf - NIH, https://www.ncbi.nlm.nih.gov/books/NBK537331/
8. What Causes High Estradiol Levels in Males? - Everlywell, https://www.everlywell.com/blog/testosterone/what-causes-high-estradiol-levels-in-males/
9. Allopregnanolone in premenstrual dysphoric disorder (PMDD) - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC7231988/
10. Role of allopregnanolone-mediated γ-aminobutyric acid A receptor, https://pmc.ncbi.nlm.nih.gov/articles/PMC10017536/
11. The Neurohypophysis: Endocrinology of Vasopressin and Oxytocin, https://www.ncbi.nlm.nih.gov/books/NBK279157/
12. Oxytocin and vasopressin in male forensic psychiatric patients with, https://www.tandfonline.com/doi/full/10.1080/14789949.2021.1985158
13. The Involvement of Prolactin in Stress-Related Disorders - MDPI, https://www.mdpi.com/1660-4601/20/4/3257
14. Left-Handedness and Male-Male Competition: Insights from Fighting, https://pmc.ncbi.nlm.nih.gov/articles/PMC10481021/
15. Treating symptomatic hyperprolactinemia in women with ... - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC3766216/
16. The Sex Hormone Precursors Dehydroepiandrosterone (DHEA) and, https://pmc.ncbi.nlm.nih.gov/articles/PMC12429729/
17. Hormonal and Metabolic Changes of Aging and the Influence ... - PMC,

https://pmc.ncbi.nlm.nih.gov/articles/PMC8020896/
18. The European Male Ageing Study (EMAS): design, methods and, https://www.researchgate.net/publication/5524289_The_European_Male_Ageing_Study_EMAS_design_methods_and_recruitment
19. Hypothalamic-Pituitary-Testicular Axis Disruptions in Older Men Are, https://www.researchgate.net/publication/5582341_Hypothalamic-Pituitary-Testicular_Axis_Disruptions_in_Older_Men_Are_Differentially_Linked_to_Age_and_Modifiable_Risk_Factors_The_European_Male_Aging_Study
20. Investigating the basis of sexual dysfunction during late-onset ... - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC6436191/
21. Age-Related Changes in the Male Reproductive System - NCBI - NIH, https://www.ncbi.nlm.nih.gov/books/NBK278998/
22. Full article: The TSST triggers self-reported stress and biological, https://www.tandfonline.com/doi/full/10.1080/10253890.2025.2503984
23. An introductory guide to conducting the Trier Social Stress Test, https://www.psychologie.uni-freiburg.de/abteilungen/psychobio/neuePublikationen/NeurosciBiobehavRev-TSSTProtocol19.pdf
24. Sex differences in salivary cortisol reactivity to the Trier Social Stress, https://d-nb.info/1185980415/34
25. Effect of 1 Week of Sleep Restriction on Testosterone Levels in, https://www.researchgate.net/publication/51183786_Effect_of_1_Week_of_Sleep_Restriction_on_Testosterone_Levels_in_Young_Healthy_Men
26. One Week of Bad Sleep Aged Your Testosterone by a Decade, https://fitchef.com/studies/sleep-deprivation-testosterone/
27. Low Testosterone or Bad Sleep? What Your Labs Might Really Be, https://www.precisionhormoneconsulting.com/blog/testosterone-sleep-problem
28. Neurobiological and Hormonal Mechanisms Regulating Women's, https://pmc.ncbi.nlm.nih.gov/articles/PMC7840832/
29. Relative energy deficiency in sport (RED-S) - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC9172228/
30. Relative Energy Deficiency in Sport (REDs) - Oxford Academic, https://academic.oup.com/edrv/article/45/5/676/7629683
31. Relative Energy Deficiency in Sports (RED-S), https://www.germanjournalsportsmedicine.com/archive/archive-2024/issue-4/relative-energy-deficiency-in-sports-red-s
32. Relative Energy Deficiency in Sport (RED-S): Scientific, Clinical, and, https://pmc.ncbi.nlm.nih.gov/articles/PMC9724109/
33. Comparison of clear and narrow outcomes on testosterone levels in, https://pmc.ncbi.nlm.nih.gov/articles/PMC5726082/
34. (PDF) Testosterone and Human Aggression: An Evaluation of the, https://www.researchgate.net/publication/7293021_Testosterone_and_Human_Aggression_An_Evaluation_of_the_Challenge_Hypothesis
35. Is testosterone linked to human aggression? A meta-analytic, https://www.researchgate.net/publication/337658220_Is_testosterone_linked_to_human_aggression_A_meta-analytic_examination_of_the_relationship_between_b

aseline_dynamic_and_manipulated_testosterone_on_human_aggression
36. The Relationship Between Testosterone and Aggression: A Meta, https://www.researchgate.net/publication/222323023_The_Relationship_Between_Testosterone_and_Aggression_A_Meta-Analysis
37. Beyond the Challenge Hypothesis: The Emergence of the Dual, https://pmc.ncbi.nlm.nih.gov/articles/PMC7311256/
38. Testosterone, Cortisol and Empathy - Behavioral Biology Lab, http://primate.uchicago.edu/2015AHBPb.pdf
39. A meta-analytical evaluation of the dual-hormone hypothesis, https://www.researchgate.net/publication/329845235_A_meta-analytical_evaluation_of_the_dual-hormone_hypothesis_Does_cortisol_moderate_the_relationship_between_testosterone_and_status_dominance_risk_taking_aggression_and_psychopathy
40. Longitudinal evidence that fatherhood decreases testosterone in, https://www.researchgate.net/publication/51631751_Longitudinal_evidence_that_fatherhood_decreases_testosterone_in_human_males
41. Longitudinal evidence that fatherhood decreases testosterone in, https://www.pnas.org/doi/pdf/10.1073/pnas.1105403108
42. Does Cosleeping Contribute to Lower Testosterone Levels in, https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0041559
43. Understanding the Oxytocin System and Its Relevance to Psychiatry, https://pmc.ncbi.nlm.nih.gov/articles/PMC4718566/
44. The active monitoring of oxytocin research evidence (AMORE, https://medpsychneuro.ruhr-uni-bochum.de/fileadmin/user_upload/The_active_monitoring_of_oxytocin_research_evidence__AMORE__platform.pdf
45. Intranasal Oxytocin: Myths and Delusions | Request PDF, https://www.researchgate.net/publication/276365841_Intranasal_Oxytocin_Myths_and_Delusions
46. Intranasal Oxytocin: Myths and Delusions - PubMed, https://pubmed.ncbi.nlm.nih.gov/26049207/
47. Association of Oxytocin Receptor Gene (OXTR) rs53576, https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0131820&type=printable
48. A common allele in the oxytocin receptor gene (OXTR) impacts, https://www.pnas.org/doi/10.1073/pnas.1003296107