

Advanced Biological and Physicochemical Dynamics of Organic Waste Valorization: Composting, Vermiculture, and NPK Fertilizer Generation

Introduction

The conversion of biological waste into nutrient-dense, stabilized agricultural amendments represents a foundational pillar of the global circular bioeconomy. As planetary agricultural systems face escalating pressures from severe soil degradation, absolute dependence on energy-intensive synthetic fertilizers, and the accumulation of environmentally disruptive waste streams, the imperative to recycle organic nutrients via biological pathways has never been more critical. The transformation of raw, highly putrescible organic residues—encompassing human excreta, livestock manure, and plant biomass—into pathogen-free, humified, and agronomically viable nitrogen-phosphorus-potassium (NPK) fertilizers relies on highly orchestrated, microbially mediated processes.

Aerobic composting and vermiculture serve as the primary biotechnological vectors for this stabilization and transformation. These processes transcend simple decomposition; they represent dynamic successions of microbial and invertebrate communities that depolymerize recalcitrant organic structures, mitigate the persistence of severe biological and chemical contaminants, and synthesize complex humic substances vital for soil structural integrity. The biochemical conversion of labile organic matter into stable humus entails a labyrinth of thermodynamics, enzymatic hydrolysis, and complex genetic regulation, ultimately dictating the bioavailability of essential macronutrients and the long-term sequestration of atmospheric carbon. By leveraging naturally occurring microbiota alongside engineered additives like biochar and inorganic catalysts, waste management paradigms can transition from mere disposal to the optimized generation of high-fidelity agricultural inputs. The following analysis exhaustively details the microbiological principles of composting and vermiculture, the thermodynamic kinetics of macronutrient generation, the mechanisms of contaminant passivation, and the integration of advanced additives designed to maximize the agronomic efficacy of organic fertilizers.

1. Microbiological Foundations and Thermodynamics of Aerobic Composting

Aerobic composting is a self-heating, highly dynamic biochemical process wherein alternating microbial consortia biodecompose organic matter in response to a continuously fluctuating physicochemical environment¹. The process is fundamentally driven by the metabolic exergonic

reactions of these microorganisms, which raise the internal temperature of the compost matrix, dictating a highly specific temporal succession of microbial taxa. The continuous availability of oxygen and moisture regulates the efficiency of this decomposition, ensuring that the process avoids anaerobic fermentation, which would otherwise lead to the production of phytotoxic organic acids and malodorous compounds.

1.1 Successional Dynamics: Mesophilic, Thermophilic, and Maturation Phases

The composting process proceeds through three distinct, temperature-driven successional phases: mesophilic, thermophilic, and maturation (or curing). Each stage is defined by specific thermal regimes and shifts in the dominant microbial phyla, reflecting a continuous evolutionary adaptation to the changing substrate¹.

During the initial mesophilic phase (ambient to approximately 45 °C), soluble, readily degradable compounds such as simple sugars, amino acids, and low-molecular-weight lipids are rapidly metabolized. This phase is characterized by an explosive proliferation of mesophilic bacteria, with Proteobacteria, Bacteroidetes, and Actinobacteria frequently dominating the community architecture¹. These microorganisms exhibit rapid growth rates and generate substantial metabolic heat through oxidative respiration, elevating the internal pile temperature at a rate of 3–5 °C per hour⁶. Genera such as *Pseudomonas* and *Klebsiella* are highly active during this early stage, initiating the preliminary hydrolysis of the feedstock⁶. Fungal populations are also prominent, utilizing the available moisture and optimal temperatures to colonize the substrate surface.

As the temperature breaches 50 °C, the process transitions into the thermophilic phase (peaking between 55 °C and 70 °C), which is arguably the most critical stage for the destruction of human and animal pathogens, the reduction of antibiotic resistance genes (ARGs), and the accelerated degradation of complex polymers¹. The extreme thermal environment suppresses or completely eliminates mesophilic populations, prompting a dramatic shift toward thermotolerant and extremophilic taxa. Firmicutes, particularly genera such as *Bacillus*, *Geobacillus*, *Thermus*, and *Saccharomonospora*, become overwhelmingly dominant⁵. This dominance is facilitated by their endospore-forming capabilities, allowing them to survive thermal shocks, and their production of highly thermostable enzymatic machinery capable of operating at elevated kinetic states⁵. Although sustained temperatures above 55 °C are generally considered lethal to most fungi, recent metagenomic analyses reveal that specific thermophilic fungi belonging to the phyla Ascomycota (e.g., *Thermomyces*, *Myriococcum*), Basidiomycota, and Rozellomycota persist¹. These thermophilic fungi are indispensable, as they release extracellular enzymes critical for the degradation of recalcitrant lignocellulosic structures that bacteria struggle to hydrolyze alone¹.

Following the depletion of easily metabolizable carbon sources, the overall microbial metabolic rate declines, initiating the cooling and maturation phase. The temperature gradually returns to ambient levels, allowing for the recolonization of mesophilic bacteria and a massive resurgence of fungal diversity². During this prolonged stabilization phase, microorganisms synthesize complex humic substances utilizing accumulated amino acids, volatile fatty acids, and modified

lignin nuclei as precursors⁸. The fungal-to-bacterial ratio typically increases significantly during maturation, contributing to the structural stabilization of the final compost and the formation of stable soil aggregates upon application³.

1.2 Enzymatic Depolymerization of Lignocellulose

The structural recalcitrance of plant-derived waste is primarily due to lignocellulose—a highly resistant, three-dimensional matrix composed of cellulose microfibrils, hemicellulose tethers, and a highly cross-linked amorphous lignin polymer⁴. The successful biotransformation of this material into bioavailable carbon relies on the synergistic secretion of hydrolytic and oxidative extracellular enzymes by both bacterial and fungal communities. Fungi, particularly white-rot basidiomycetes and various aggressive ascomycetes, are the most efficient degraders of the lignin shield¹¹.

Lignin degradation is a non-specific, oxidative process mediated by a suite of ligninolytic enzymes, predominantly laccase (Lac), lignin peroxidase (LiP), manganese peroxidase (MnP), and versatile peroxidase (VP)¹². Laccases are multicopper oxidases capable of oxidizing both phenolic and non-phenolic compounds through single-electron transfers. This oxidation generates highly reactive free radicals that lead to the spontaneous, non-enzymatic cleavage of carbon-carbon and ether bonds within the lignin polymer¹¹. Manganese peroxidases operate by oxidizing Mn(II) to Mn(III), which is then chelated by organic acids (such as oxalate) and acts as a diffusible redox mediator that can penetrate deep into the wood matrix to oxidize phenolic lignin structures at a distance from the fungal hyphae¹³.

Fungal and bacterial populations exhibit profound cooperative functional complementarity in this process. Fungi utilize their exploratory hyphal networks and robust oxidative enzymes to physically and chemically breach the protective lignin barrier, thereby exposing the underlying cellulose and hemicellulose¹⁴. Bacteria, equipped with intricate cellulosomes and free cellulases (including endoglucanases, exoglucanases, and β -glucosidases), subsequently colonize these exposed micro-sites, rapidly hydrolyzing the polysaccharides into fermentable sugars¹¹. Furthermore, specific thermophilic fungi, such as *Trametes troglia*, maintain continuous laccase production even at elevated temperatures (e.g., 50 °C) by synthesizing specialized heat shock proteins¹¹. This ensures that lignocellulose depolymerization—a highly endothermic barrier to compost maturation—persists uninterrupted throughout the most active thermal phases of the composting cycle.

2. Vermiculture: The Earthworm-Microbe Symbiosis

Vermiculture is a specialized, mesophilic biotransformation process that harnesses the synergistic, mutualistic interactions between earthworms (commonly epigeic species such as *Eisenia fetida* and *Eudrilus eugeniae*) and distinct microbial communities to stabilize organic residues⁷. Unlike conventional thermophilic composting, which relies on thermal destruction and oxidative stress, vermicomposting operates strictly at ambient temperatures. It relies on the intense mechanical and biochemical processing of waste within the earthworm digestive tract to achieve rapid stabilization and nutrient mineralization.

2.1 The Earthworm Gut as a Tubular Bioreactor

The earthworm gut functions as a highly efficient, small-scale, continuous-flow tubular bioreactor that fundamentally alters the physical, chemical, and biological properties of ingested organic matter¹⁷. As earthworms engulf substrates, the material is subjected to rigorous mechanical grinding in the muscular gizzard, which dramatically decreases particle size and exponentially increases the specific surface area available for microbial and enzymatic action¹⁸.

The internal gut environment provides optimal conditions for microbial metabolism: it maintains ideal moisture levels, a near-neutral pH buffered by calciferous glands, and high concentrations of easily assimilable carbon derived from mucopolysaccharides continuously secreted by the earthworm's intestinal lining. This nutrient-rich, protected environment collectively stimulates explosive microbial proliferation, often leading to microbial biomass densities far exceeding those found in the surrounding substrate¹⁷.

Simultaneously, the earthworm secretes a diverse array of endogenous digestive enzymes, including cellulases, proteases, amylases, and chitinases. These endogenous enzymes operate in tandem with exogenous enzymes produced by the ingested, transient, and resident gut microbiota¹⁷. This powerful enzymatic consortium rapidly hydrolyzes insoluble organic compounds into soluble, highly bioavailable nutrients. The transit of material through the gut selective pressure fundamentally modifies the microbial community structure. Specific bacterial and fungal taxa are digested and assimilated for earthworm nutrition, while beneficial, symbiotic microbes—particularly plant-growth-promoting rhizobacteria (PGPR)—often emerge in significantly higher concentrations in the resulting fecal castings (vermicompost)²¹. Furthermore, earthworm mucus has been shown to contain specific biochemical agents that disrupt bacterial communication, potentially inhibiting horizontal gene transfer mediated by quorum-sensing signaling molecules, thereby assisting in the suppression of pathogenic traits and antibiotic resistance¹⁹.

2.2 Synthesis of Phytohormones and Indole-3-Acetic Acid (IAA)

A defining hallmark of vermicompost quality, which elevates it above conventional aerobic compost, is its profound enrichment in plant growth-regulating substances, particularly phytohormones such as auxins, gibberellins, and cytokinins²³. Among these, Indole-3-acetic acid (IAA) is the most extensively characterized natural auxin, playing a paramount and irreplaceable role in regulating plant cell division, cellular elongation, vascular tissue differentiation, and adventitious root development²⁴.

The biosynthesis of IAA in vermicompost is not primarily performed by the earthworm itself, but rather driven by the intense metabolic activity of gut-associated microorganisms, particularly specific strains of *Bacillus*, *Pseudomonas*, and *Streptomyces* that proliferate within the castings²⁴. These specialized bacteria synthesize IAA predominantly via the indole-3-pyruvic acid pathway, utilizing tryptophan—an amino acid derived from the intense proteolytic degradation of organic proteins within the gut—as the primary biochemical precursor²⁶.

Advanced transcriptomic and genetic analyses of PGPR isolated from *Eisenia fetida* castings have identified the distinct upregulation of specific gene products (e.g., *dhaS*, *patB*, *yclC*, *yhcX*) that are directly involved in IAA biosynthesis²⁸.

Empirical evaluations unequivocally demonstrate that the concentration of exogenous IAA in

agricultural soils is directly proportional to the application rate of high-quality vermicompost. For example, controlled studies indicate that the incorporation of 1 kg of vermicompost into a highly sterile sand-based matrix increases the baseline concentration of exogenous IAA by 0.0470 mg/g to 0.0642 mg/g over a standard growing season. This biochemical enrichment translates directly into radically enhanced root morphometrics (e.g., maximum root development volume reaching 370 cm³ in augmented plots) and significantly increased agricultural yields compared to unamended controls²⁴. Additionally, specific vermi-bacteria like *Bacillus xiamenensis* strain VITMSJ3, isolated from *Eisenia fetida*, not only produce substantial IAA but also exhibit extreme resistance to heavy metals, tolerating lead up to 4000 ppm and cadmium up to 450 ppm, making vermicompost an exceptional tool for phytoremediation in marginal soils²⁵.

3. Feedstock Characterization and NPK Profiles

The primary agronomic value of processed organic waste lies in its capacity to serve as a comprehensive, slow-release source of essential macronutrients—nitrogen (N), phosphorus (P), and potassium (K)—while simultaneously replenishing depleted soil organic carbon pools. The ultimate efficiency of nutrient release and the fertilizer value of the final compost are strictly dictated by the elemental composition of the raw feedstock and the specific biogeochemical kinetics governing their mineralization.

Organic feedstocks exhibit extreme heterogeneity in their baseline elemental composition, necessitating strategic, scientifically formulated blending to optimize the initial carbon-to-nitrogen (C/N) ratio (ideally maintained between 20:1 and 30:1) and ensure a balanced final NPK profile³.

3.1 Human Waste: Humanure and Source-Separated Urine

Historically stigmatized, human excreta represents a highly concentrated, yet vastly underutilized, nutrient reservoir capable of significantly offsetting synthetic fertilizer demand. An average adult human produces approximately 90 g of organic matter (dry weight), 10–12 g of nitrogen, 2 g of phosphorus, and 3 g of potassium on a daily basis²⁹.

When separated at the source, urine and feces serve entirely different agronomic roles. Urine is the primary vector for soluble macronutrients, containing 70–80% of total excreted nitrogen (predominantly in the form of rapidly hydrolyzable urea) and the vast majority of excreted potassium. Median human urine generation rates approximate 1.42 liters per capita per day, yielding a dry solids content of 59 g/cap/day²⁹. Conversely, human feces contain the bulk of the organic carbon, undigested dietary fiber, and phosphorus. Bacterial biomass comprises 25–54% of the dry solids in raw feces³⁰. When solid human feces are subjected to aerobic composting for 30 days, the resulting substrate exhibits exceptionally rich nutritional parameters, with dry weight concentrations often reaching 15.62% carbon, 1.50% to 7.00% nitrogen, and remarkably

high phosphorus levels ranging from 3.00% to 6.45% (P_2O_5)³¹. Dehydrated night soil, historically referred to as "poudrette," can provide highly concentrated fertilization, typically

containing up to 5.5% N, 4% P_2O_5 , and 2% K_2O ³². Furthermore, large-scale municipal biosolids derived from human waste, such as "Milorganite" produced in Milwaukee, function as highly effective, heat-dried slow-release fertilizers that feed plant roots over an 8-10 week

period, averting the rapid nutrient crash associated with synthetic urea³³.

Human Waste Fraction	Nitrogen (N)	Phosphorus (P ₂ O ₅)	Potassium (K ₂ O)	Key Characteristics
Solid Feces (Raw)	4.8% - 7.0%	3.0% - 5.4%	1.0% - 2.5%	High pathogen load; requires thermophilic processing or extended curing. ³¹
Fecal Compost (30-day)	1.5% - 2.0%	~6.45%	~1.5%	Exceptional phosphorus source; high bacterial biomass contributing to stable organic N. ³⁰
Dehydrated Night Soil	~5.5%	~4.0%	~2.0%	Concentrated "poudrette"; high risk of ammonia volatilization if improperly stored. ³²

3.2 Animal Manures and Animal-Based Concentrates

Livestock and poultry manures are traditional agricultural feedstocks; however, their nutrient profiles are highly variable, dependent entirely on the animal species, gastrointestinal physiology, dietary intake, and the specific bedding materials utilized in the husbandry process³⁴.

Poultry manure is globally recognized for its exceptional nutrient density, averaging 3.03% N, 2.63% P_2O_5 , and 1.4% K_2O on a dry weight basis³⁵. This high concentration is due to the avian physiological mechanism of co-excreting highly concentrated uric acid with feces.

However, its low C/N ratio makes it highly susceptible to catastrophic ammonia (NH_3) volatilization if not properly bulked with high-carbon amendments. In stark contrast, well-decomposed ruminant farmyard manure (a mixture of cattle dung, urine, and lignocellulosic

litter) presents a much more dilute nutritional profile, averaging 0.5% N, 0.2% P_2O_5 , and 0.5% K_2O ³⁵. Despite its low macronutrient density, it is unparalleled as a precursor for stable humification and soil structural improvement. Sheep and goat droppings offer intermediate values of approximately 3% N, 1% P_2O_5 , and 2% K_2O , providing excellent physical aeration to compost piles due to their pelleted structure³⁵. For targeted agronomic application, animal processing byproducts serve as highly concentrated organic amendments. Blood meal offers an immense, rapidly mineralizable nitrogen source (10–12% N), while raw and steamed bone meals serve as the premier organic sources of slow-release phosphorus (20–30% P_2O_5)³⁵.

Animal-Derived Manures / Meals	Nitrogen (N) %	Phosphorus (P ₂ O ₅) %	Potassium (K ₂ O) %
Poultry Manure	3.03	2.63	1.40
Sheep/Goat Manure	3.00	1.00	2.00
Cattle Farmyard Manure	0.50	0.20	0.50
Blood Meal	10.0 - 12.0	1.0 - 2.0	1.00
Raw Bone Meal	3.0 - 4.0	20.0 - 25.0	Traces
Meat Meal	10.5	2.5	0.5

Data synthesized from agricultural reference standards³².

3.3 Plant Residues, Green Manures, and Oil Cakes

Agricultural residues and yard wastes are critical not primarily for their macronutrient content, but for providing the necessary carbon skeleton, porosity, and moisture regulation during composting. However, specific plant derivatives are utilized precisely for their fertilizing capacity. Green manures, particularly leguminous crops cultivated specifically for soil incorporation, contribute significant biologically fixed nitrogen. Species such as *Sesbania aculeata* (Dhaincha) and *Crotalaria juncea* (Sunnhemp) average 2.3–3.5% N, along with valuable potassium reserves ranging from 1.2% to 2.2% K_2O ³².

Following the extraction of botanical oils, the residual biomass—known as oil cakes—provides some of the most potent, plant-derived concentrated organic manures available. Edible oil cakes (e.g., groundnut, safflower, and decorticated cotton seed) decompose rapidly, releasing high concentrations of nitrogen (up to 7.9% N). Non-edible oil cakes (e.g., castor, neem, karanj), while also nutrient-dense (3.9–4.9% N), offer the added agronomic benefit of containing naturally occurring biochemical compounds that act as nematicides and nitrification inhibitors, thereby slowing the conversion of ammonium to nitrate and reducing leaching losses³².

Plant-Derived Concentrates (Oil Cakes)	Nitrogen (N) %	Phosphorus (P₂O₅) %	Potassium (K₂O) %
Groundnut Cake (Edible)	7.3	1.5	1.3
Safflower Cake (Decorticated)	7.9	2.2	1.9
Cotton Seed Cake (Decorticated)	6.4	2.9	2.2
Castor Cake (Non-edible)	4.3	1.8	1.3
Neem / Karanj Cake (Non-edible)	3.9 - 5.2	0.9 - 1.8	1.2 - 1.5

Data synthesized from agricultural reference standards³².

4. Nitrogen Transformations and Mineralization

Kinetics

Nitrogen within raw organic waste exists predominantly in complex, highly polymerized organic forms, primarily locked within proteins, amino acids, microbial cell walls, and nucleic acids. In this state, it is entirely unavailable for plant uptake. To become agronomically functional, the organic nitrogen must undergo a series of biologically mediated transformations collectively known as mineralization, encompassing ammonification and nitrification³⁶.

4.1 The Ammonification and Nitrification Pathways

The initial step, ammonification, involves the enzymatic hydrolysis of organic nitrogenous compounds by a broad spectrum of heterotrophic bacteria and fungi, resulting in the release of

ammonium (NH_4^+) and free ammonia (NH_3)³⁶. The equilibrium between these two compounds is strictly pH-dependent; at elevated alkaline pH levels commonly observed during

the active phase of composting, highly volatile NH_3 dominates, leading to severe nitrogen losses and the generation of noxious odors.

As composting proceeds, readily degradable carbon is exhausted, temperatures cool, and the pH stabilizes to near-neutral levels, establishing the optimal microenvironment for nitrification. Nitrification is a specialized, two-step aerobic chemolithoautotrophic oxidation process. First,

ammonia-oxidizing bacteria (AOB) and ammonia-oxidizing archaea (AOA) oxidize NH_3 / NH_4^+ to hydroxylamine (NH_2OH) and subsequently to nitrite (NO_2^-). This initial,

rate-limiting reaction is catalyzed by the transmembrane enzyme ammonia monooxygenase, which is encoded by the highly conserved *amoA* functional gene³⁴. High nitrogen inputs, such as chronic urea application, significantly upregulate *amoA* expression, increasing AOB population densities tenfold⁴⁰. Subsequently, nitrite-oxidizing bacteria (NOB) rapidly oxidize the

highly toxic nitrite intermediate into stable nitrate (NO_3^-). This second step is mediated by the nitrite oxidoreductase (NXR) enzyme, encoded by the *nxr* gene³⁴. The progressive accumulation

of nitrate and the corresponding decline of the NH_4^+ / NO_3^- ratio serves as a primary, quantifiable indicator of compost maturity and stability¹.

4.2 Mathematical Modeling of Mineralization Kinetics

When finished compost is applied to agricultural soils, the release of its residual organic nitrogen pool is a time-dependent, microbially driven process that must be mathematically modeled to accurately predict nutrient availability and synchronize it with crop demand.

Nitrogen mineralization kinetics most frequently adhere to a first-order exponential decay model, formulated as:

$$N_{min}(t) = N_0 (1 - e^{-kt})$$

where $N_{min}(t)$ is the cumulative amount of nitrogen mineralized at time t (days), N_0

represents the absolute size of the potentially mineralizable organic nitrogen pool, and k

represents the first-order mineralization rate constant (day^{-1})⁴².

The kinetics of mineralization are profoundly influenced by the biochemical quality of the compost and the ambient soil temperature. Poultry manure, due to its low C:N ratio and high concentration of labile nitrogen, typically exhibits a significantly higher mineralization rate

constant (k) and a larger mineralizable pool (N_0) compared to cattle manure or green waste compost⁴². Incubation studies demonstrate that poultry manure undergoes a rapid initial release

phase during the first 30 days, making the nitrogen immediately available, but also highly susceptible to leaching if crop root systems are not adequately established⁴⁴. In contrast, highly humified amendments like vermicompost and mature farmyard manure exhibit significantly slower, more stable mineralization rates. These materials possess a high proportion of stable, recalcitrant structures where nitrogen is complexed within organic matrices, resulting in low k values (e.g., $k = 0.009$)⁴².


Temperature acts as a critical kinetic accelerator. Across all manure types, increasing the incubation temperature from 15 °C to 35 °C dramatically amplifies the mineralizable nitrogen fraction (e.g., a 10–20% increase in mineralization rate at 35 °C compared to lower regimes), directly correlating with enhanced microbial metabolic and enzymatic activities⁴⁴.

5. Phosphorus Recovery and Biological Potassium Solubilization

While nitrogen is predominantly governed by biological oxidation and reduction, the management of phosphorus and potassium in organic waste valorization relies on targeted chemical precipitation and specialized microbial weathering mechanisms to transform insoluble mineral complexes into bioavailable plant nutrients.

5.1 Phosphorus Sequestration via Struvite Crystallization

Liquid and semi-solid effluents, such as source-separated human urine, swine slurry, and anaerobic digestates, are rich in soluble reactive phosphorus and ammonium, but are difficult to store and apply agronomically. To maximize nutrient retention from these streams, phosphorus and nitrogen can be co-precipitated as struvite (magnesium ammonium phosphate hexahydrate,



)⁴⁵. The synthesis of highly pure struvite crystals requires an exact equimolar ratio (1:1:1) of Mg^{2+} , NH_4^+ , and PO_4^{3-} under strictly controlled alkaline conditions (typically pH 8.0–9.0). When integrated with solid compost or biochar matrices (a practice known as co-composting), struvite acts as a premium, slow-release fertilizer that fundamentally alters the nitrogen loss dynamics of the pile, severely curtailing both ammonia volatilization and phosphate leaching⁴⁷. Recent innovations in crystallization technology have demonstrated that utilizing biochar as a "seeding" material provides a vast, highly porous surface area for nucleation. Biochar seeding promotes rapid struvite formation, intensifies overall nutrient recovery (exceeding 97% for phosphate and 94% for ammonium), and yields larger, higher-quality struvite crystals (e.g., increasing particle size from 70 µm to 100 µm)⁴⁶.

5.2 Microbial Potassium Solubilization (KSB)

Although organic wastes contain inherent potassium, over 90% of total soil potassium globally remains locked in highly insoluble primary silicate minerals, such as feldspar, mica, and illite, rendering it entirely unavailable for root uptake⁴⁸. To mobilize this vast structural reserve and reduce reliance on finite, salt-heavy synthetic potassium chloride (muriate of potash), organic

fertilizers and seed coats are increasingly inoculated with Potassium Solubilizing Bacteria (KSB), with elite strains including *Bacillus mucilaginosus*, *Pseudomonas putida*, and *Frateuria aurantia*⁴⁹.

These powerful rhizobacteria solubilize potassium through two primary, intertwined biochemical mechanisms:

1. **Acidolysis via Organic Acid Exudation:** The bacteria secrete substantial quantities of low-molecular-weight organic acids, primarily gluconic, citric, malic, oxalic, and tartaric acids. These acids donate protons (H^+), sharply lowering the micro-environmental pH of the rhizosphere. This localized acidification directly degrades the rigid aluminosilicate lattice of the minerals⁴⁸.
2. **Chelation and Complexolysis:** The dissociated anions of these organic acids act as powerful biological chelating agents. They form stable complex structures with structural cations such as Al^{3+} , Fe^{3+} , and Ca^{2+} that hold the mineral lattice together. By removing these stabilizing cations, the mineral structure destabilizes, facilitating the rapid release of the entrapped K^+ ions into the soil solution⁴⁸.

Field evaluations of KSB-enriched biofertilizers demonstrate their capacity to supply the

equivalent of 20–40 kg K_2O per hectare per season. Beyond simple nutrient provision, the application of KSB augments root proliferation, enhances sugar translocation, and significantly improves drought resilience and disease resistance by strengthening plant cell walls, all while actively avoiding the chloride-induced salinity stress associated with conventional mineral top-dressings⁴⁹.

6. Humification Pathways, Structural Evolution, and Phytotoxicity

The ultimate objective of processing organic waste is not merely the breakdown of material, but successful humification—the profound biochemical transformation of labile, putrescible organic matter into highly stable, recalcitrant humic substances (HS). These substances impart long-term structural, chemical, and biological benefits to soil, fundamentally driving cation exchange capacity and moisture retention⁵³.

Humic substances are operationally partitioned into three primary fractions based on their solubility at different pH levels: fulvic acids (FA; lower molecular weight, highly soluble at all pH levels), humic acids (HA; higher molecular weight, soluble in alkali but insoluble in acid), and humin (highly recalcitrant, insoluble at all pH levels)⁹.

6.1 Polymerization, Condensation, and Supramolecular Assembly

Humification is not a simple, linear degradation pathway; rather, it represents a highly complex, dynamic continuum of molecular transformations⁵⁵. As polysaccharides, aliphatic carbon chains, and structural proteins are enzymatically degraded during the aggressive mesophilic and thermophilic phases, vast quantities of reactive intermediates—such as polyphenols, reducing sugars, amino acids, and modified lignin nuclei—accumulate in the matrix.

During the cooling and maturation phase, these reactive intermediates undergo intricate abiotic and biotic condensation and polymerization reactions (e.g., Maillard reactions and quinone-amino acid couplings) to form high-molecular-weight, highly aromatic humic networks⁹. Modern chemical analysis also supports a supramolecular model of humus, suggesting that rather than forming massive, covalently bonded macro-polymers, humic substances are composed of relatively small, heterogeneous biomolecules bound tightly together by weak non-covalent intermolecular forces (hydrogen bonds, van der Waals forces) that exhibit emergent macro-polymer properties⁵⁵.

The progression of humification is typically quantified by the Degree of Polymerization (DP) and the ratio of humic acid carbon to fulvic acid carbon (HA/FA). A high-quality, mature compost is characterized by a decreasing concentration of FA and an increasing concentration of HA, as smaller fulvic components act as precursors and are integrated into larger, more stable humic acid structures¹⁰. The introduction of specific additives, such as biochar, clay minerals (e.g., bentonite, attapulgite), and iron-based catalysts, has been shown to significantly accelerate this transition. Iron-based additives, for instance, promote intense Fenton-like redox reactions, elevating final HA concentrations by 82% to over 260% relative to unamended controls⁵⁴. Vermicomposting also dramatically enhances humification; the continuous mechanical and enzymatic processing within the earthworm gut rapidly diminishes aliphatic, polypeptidic, and carbohydrate components while exponentially increasing the density of oxygenated, aromatic functional groups⁵⁷.

6.2 Spectroscopic Characterization of Compost Maturity

The structural evolution of organic matter during composting cannot be fully understood through wet chemistry alone; it requires rigorous monitoring utilizing advanced, non-destructive spectroscopic techniques:

- **UV-Visible Spectroscopy:** Evaluates the E4/E6 ratio (the ratio of absorbance at 465 nm to 665 nm). A declining E4/E6 ratio indicates increasing molecular weight, higher degrees of aromatic condensation, and advanced maturation of the humic acids⁵⁷.
- **Fourier-Transform Infrared (FTIR) Spectroscopy:** Tracks the progressive destruction of aliphatic bands and the distinct emergence of aromatic ($C = C$) and oxygenated functional groups (such as carboxyl and phenolic hydroxyl moieties) that are characteristic of mature, stable humus capable of heavy metal chelation⁹.
- **Raman Spectroscopy:** Specifically measures the degree of carbon aromatization and structural ordering (e.g., analyzing the D and G bands of carbonaceous structures). It provides high-resolution data indicating that advanced treatments like vermicomposting induce a markedly higher degree of aromatic ring polycondensation and carbon stability compared to passive windrow composting⁵⁷.
- **X-Ray Diffraction (XRD):** Utilized to investigate the progressive physical degradation of crystalline cellulose fractions as the composting time extends⁵⁸.

6.3 Phytotoxicity and the Germination Index (GI)

Applying immature, unstabilized compost to agricultural soils induces severe phytotoxicity, characterized by the catastrophic inhibition of seed germination, severe root necrosis, and

nutrient lock-up. This toxicity is primarily mediated by the presence of high concentrations of un-ionized ammonia (NH_3), heavy metals, and short-chain volatile fatty acids (VFAs)—particularly acetic, propionic, and highly toxic butyric acid—which are byproducts of anaerobic micro-sites within the pile⁶¹.

Phytotoxicity is empirically quantified utilizing the Germination Index (GI) bioassay, a highly sensitive biological standard most frequently performed using garden cress (*Lepidium sativum*) or cucumber (*Cucumis sativus*) seeds due to their rapid and sensitive response to aqueous compost extracts⁶². The GI integrates both the relative rate of seed germination and the vigor of radicle (root) elongation relative to a distilled water control. It is calculated using the formula:

$$GI = \left(\frac{\text{Germinated seeds}_{treatment}}{\text{Germinated seeds}_{control}} \right) \times \left(\frac{\text{Root length}_{treatment}}{\text{Root length}_{control}} \right) \times 100$$

As microbial activity exhausts volatile organic acids and nitrification successfully converts toxic NH_4^+ into benign NO_3^- , phytotoxicity steadily dissipates⁶³. A compost extract yielding a GI > 60% indicates acceptable baseline maturity, while a GI > 80% signifies the complete disappearance of phytotoxins and confirms the absolute agronomic safety of the final product⁶¹. Studies demonstrate that when optimum composting conditions are met (e.g., windrows turned every 4 days with precise moisture control), severe phytotoxicity can be entirely resolved within 56 days⁶³.

7. Contaminant Passivation and Ecological Risk Mitigation

The agronomic utilization of organic fertilizers, particularly those derived from human biosolids, poultry litter, and municipal solid waste, carries inherent toxicological risks. These risks stem from the presence of resilient biological pathogens, heavy metals, pharmaceutical residues, and emerging synthetic contaminants. Effective, scientifically validated bioprocessing strategies must reliably mitigate these hazards to prevent their introduction into the human food chain.

7.1 Pathogen Destruction and Thermal Inactivation

Human and animal excreta harbor highly concentrated, diverse arrays of enteric pathogens, including *Salmonella* spp., *Escherichia coli*, *Vibrio cholerae*, and highly resistant helminth ova⁶⁵. Regulatory frameworks, such as the US EPA's Process to Further Reduce Pathogens (PFRP), dictate strict operational parameters to ensure safety. To achieve PFRP status, composting systems must maintain internal pile temperatures of at least 55 °C for a minimum of 3 consecutive days in an Aerated Static Pile (ASP) system, or 15 consecutive days (incorporating at least five mechanical turnings) in an open windrow system⁶.

These exact time-temperature matrices are specifically engineered to inactivate the most physically resilient biological contaminants, particularly the thick-shelled ova of the human

roundworm, *Ascaris lumbricoides*⁶⁶. *Ascaris* eggs are highly resistant to standard chemical disinfection (e.g., chlorination) and desiccation, establishing them as the ultimate biological indicator for compost sanitization. Extended thermophilic conditions (e.g., maintaining 55 °C for several weeks) promote rapid, irreversible protein denaturation and lipid membrane disruption, achieving a >99% reduction in viable *Ascaris* eggs within 3 to 7 weeks⁶. Furthermore, mature, cured compost establishes a highly diverse, homeostatic microbial community that exerts intense competitive exclusion against transient enteric bacteria. This complex microbial ecology effectively suppresses the dangerous regrowth of opportunistic pathogens, such as *Salmonella typhimurium*, which could otherwise recolonize the nutrient-rich matrix if introduced post-treatment⁶⁸.

7.2 Heavy Metal Speciation and Bioavailability

Unlike complex organic contaminants, elemental heavy metals (e.g., Cu, Zn, Cd, Pb, Cr) cannot be biologically destroyed or degraded. However, the processes of composting and vermicomposting fundamentally alter their physicochemical speciation, thereby drastically reducing their biological mobility, environmental leachability, and subsequent phytotoxicity⁷⁰. The environmental risk of heavy metals is evaluated utilizing the sequential extraction procedure developed by the Community Bureau of Reference (BCR), which rigorously partitions metals into four distinct functional fractions based on their solubility:

1. **F1 (Acid-Soluble/Exchangeable):** Highly mobile, acutely bioavailable, and easily absorbed by plant roots.
2. **F2 (Reducible):** Bound to iron and manganese oxides; moderately mobile under reducing conditions.
3. **F3 (Oxidizable):** Bound to organic matter, humic substances, and sulfides; highly stable under standard non-oxidizing conditions.
4. **F4 (Residual):** Locked permanently within the primary mineral lattice; environmentally inert and non-toxic⁷⁰.

During the aggressive degradation of organic matter and the concomitant synthesis of humic acids, free metal cations in the compost matrix interact heavily with the negatively charged oxygenated functional groups (carboxyl, phenolic hydroxyl) of the newly formed humic structures. This intense complexation drives a distinct, quantifiable speciation shift, transforming dangerous metals from the highly bioavailable F1 and F2 fractions into the highly stable F3 and F4 fractions⁷².

Vermicomposting exerts an even more profound passivation effect on heavy metals. As contaminated material passes through the earthworm gut, the unique biochemical environment facilitates the rapid formation of organo-metallic complexes. Furthermore, earthworms inherently bioaccumulate substantial quantities of heavy metals directly into their tissue (with bioconcentration factors heavily favoring Cd, Ni, and Cu), effectively scrubbing the substrate. Consequently, the resulting earthworm castings exhibit a significantly lower total bioavailable concentration of heavy metals compared to the raw feedstock⁷⁰. The integration of highly porous additives like biochar, dolomite, and diatomite as bulking agents further amplifies metal immobilization through deep micro-pore sequestration, robust ion exchange, and surface electrostatic adsorption, reducing zinc and copper leaching potentials by over 60%⁵⁸.

7.3 Antibiotic Resistance Genes (ARGs) and the Rebound Phenomenon

Intensive livestock operations rely heavily on veterinary antibiotics, rendering animal manures a critical environmental reservoir for residual antibiotics, antibiotic-resistant bacteria (ARB), and antibiotic resistance genes (ARGs)⁷⁹. The unchecked environmental propagation of ARGs via horizontal gene transfer (HGT), mediated by mobile genetic elements (MGEs) such as plasmids, transposons, and Class 1 integrons (*int1*), poses a severe, escalating threat to global public health and ecological stability⁸⁰.

The thermophilic phase of aerobic composting serves as the primary physical mechanism for ARG attenuation. Sustained high temperatures physically denature resistant plasmids and systematically inactivate the mesophilic bacteria (e.g., Bacteroidetes and certain strains of Proteobacteria) that serve as the primary host vehicles for these genes⁶. However, a pervasive and highly concerning challenge in waste management is the "ARG rebound" phenomenon—a dramatic resurgence in both the absolute and relative abundance of ARGs (such as *sul1*, *tet(33)*, and *tetG*) during the cooling and maturation phases of compost⁷⁹.

This rebound is driven by the rapid proliferation of surviving, multidrug-resistant host bacteria as temperatures normalize, combined with the intense co-selection pressure exerted by persistent co-contaminants like heavy metals. Heavy metals and antibiotics frequently share identical efflux pump mechanisms or genetic loci on the same MGEs (co-resistance and cross-resistance), meaning the mere presence of heavy metals forces the microbial community to maintain ARGs even after the actual antibiotics have fully degraded⁸⁰.

To effectively suppress ARG rebound, advanced operational interventions are mandatory. Extending the duration of the thermophilic phase—such as through hyperthermophilic composting utilizing semi-permeable membrane covers—or incorporating specific additives like mature humus soil and biochar significantly limits ARG proliferation⁷⁹. Biochar is particularly effective; it physically adsorbs heavy metals and residual antibiotics (thereby neutralizing the co-selection triggers) and interferes with the cell-to-cell contact required for conjugative horizontal gene transfer, dramatically downregulating the abundance of *int1* and stifling the spread of resistance⁸⁰.

7.4 Emerging Contaminants: PFAS and Microplastics

While thermal and biological treatments successfully degrade most standard pharmaceuticals and bacterial pathogens, modern emerging synthetic contaminants present profound, unresolved challenges for the composting industry. Per- and polyfluoroalkyl substances (PFAS)—ubiquitous, highly fluorinated "forever chemicals"—and microplastics exhibit extreme chemical and thermal stability⁸³.

Current research conclusively indicates that PFAS compounds do not degrade during conventional aerobic composting, anaerobic digestion, or vermicomposting processes⁸³. When highly contaminated municipal biosolids or unsegregated industrial sludges are utilized as primary feedstocks, these indestructible contaminants persist entirely intact in the final organic fertilizer. This poses a catastrophic risk of direct bioaccumulation in soil invertebrates, such as earthworms, and subsequent, unimpeded transfer into higher agricultural food webs⁸³.

Consequently, rigorous source separation, strict legislative limits on municipal sludge inputs, and stringent feedstock screening remain the only currently viable defenses against the systematic accumulation of PFAS in compost-amended soils.

7.5 Greenhouse Gas Mitigation and Genetic Regulation

While the recycling of organic waste fundamentally prevents the massive methane generation associated with anaerobic landfilling, improperly managed aerobic composting can still

inadvertently emit potent greenhouse gases, specifically methane (CH_4) and nitrous oxide (N_2O).

Methane emissions in compost piles originate from localized anaerobic micro-sites that develop when moisture content exceeds 65% or when physical turning is neglected. The net release of methane into the atmosphere is strictly dictated by the dynamic biological equilibrium between

methanogenic archaea (which generate CH_4) and methanotrophic bacteria (which consume and oxidize CH_4 into CO_2)⁸⁵. The populations of these competing functional guilds are precisely quantified via qPCR assays targeting the *mcrA* (methyl coenzyme M reductase) gene for methanogens and the *pmoA* (particulate methane monooxygenase) gene for methanotrophs⁸⁵. Maintaining high matrix porosity and employing forced mechanical aeration (e.g., Aerated Static Piles, ASP) optimizes oxygen diffusion throughout the core. This physical intervention successfully suppresses *mcrA* expression and highly stimulates *pmoA* activity, allowing methanotrophs to biologically neutralize methane before it escapes the pile surface⁸⁵. Nitrous oxide, an immensely potent greenhouse gas possessing a 100-year global warming

potential 273 times greater than CO_2 , is primarily generated as an unstable intermediate byproduct of incomplete nitrification and denitrification pathways within the pile⁸⁷. Denitrifying

bacteria convert accumulated nitrate back into inert nitrogen gas (N_2), a process whose final, environmentally benign step is catalyzed by the nitrous oxide reductase enzyme, which is encoded by the *nosZ* functional gene⁸⁸.

The incorporation of biochar into the compost matrix acts as a master variable for emission mitigation. Biochar physically adsorbs ammonium and nitrate, preventing their rapid conversion into volatile intermediates. More importantly, comprehensive meta-analyses reveal that biochar amendments significantly and consistently upregulate the abundance and expression of the *nosZ* gene within the compost microbiome⁸⁹. By genetically promoting the complete reduction of N_2O to harmless N_2 gas, biochar co-composting drastically curtails nitrous oxide emissions, effectively closing the nitrogen loop and yielding a superior, highly climate-resilient organic fertilizer⁸⁹.

Conclusion

The biological conversion of crude organic waste into high-value, stabilized NPK fertilizers represents a highly advanced intersection of microbiology, biochemistry, and thermodynamics.

The synergistic action of extremophilic bacteria and highly evolved lignocellulolytic fungi drives the rapid depolymerization of recalcitrant feedstocks, while the complex tubular bioreactor of the earthworm gut fundamentally alters the bio-architecture of the matrix, stimulating the targeted synthesis of vital plant phytohormones like Indole-3-acetic acid. Optimization of these systems—through precise kinetic modeling, C/N balancing, forced mechanical aeration, and the strategic addition of functional materials like biochar and potassium-solubilizing bacteria—strictly dictates the efficiency of nutrient mineralization. Furthermore, these engineered biological pathways are essential for suppressing the proliferation of dangerous enteric pathogens, reducing the toxic bioavailability of heavy metals, and mitigating the grave ecological risk of antibiotic resistance gene rebound. By rigorously controlling fundamental operational parameters, modern composting and vermiculture transcend simple waste disposal, establishing highly engineered, closed-loop biological technologies that are absolutely essential for ensuring sustainable global soil fertility and safeguarding long-term environmental health.

Works cited

1. Microbial community dynamics in the mesophilic and thermophilic, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8654937/>
2. Microbial Community Succession and Response to Environmental, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6431636/>
3. Bacterial and fungal communities and contribution of ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5727367/>
4. Characterization of Thermophilic Lignocellulolytic Microorganisms in, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8385673/>
5. Succession of Microbial Community during the Co-Composting of, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9407818/>
6. A Review on the Degradation of Antibiotic Resistance Genes During, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12389969/>
7. Changes in Bacterial and Fungal Communities across Compost, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3836849/>
8. Study on dynamic changes of microbial community and ... - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9077819/>
9. Food waste humification: a process analysis - ResearchGate, https://www.researchgate.net/publication/365434740_Food_waste_humification_a_process_analysis
10. Sustainable fertilizers from wastes: a strategy to enhance soil carbon, <https://www.oaepublish.com/articles/cf.2025.91>
11. Microbial Degradation of Lignocellulose for Sustainable Biomass, <https://www.mdpi.com/2071-1050/17/9/4223>
12. The Role of Ligninolytic Enzymes in Sustainable Agriculture - MDPI, <https://www.mdpi.com/2073-4395/15/2/451>
13. Role of lignin peroxidase, manganese peroxidase, and laccase, https://www.researchgate.net/publication/244753256_Microbial_degradation_of_lignin_Role_of_lignin_peroxidase_manganese_peroxidase_and_laccase
14. Lignocellulose dissociation with biological pretreatment towards the,

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC9535326/>
15. Fungal–bacterial interaction networks in lignocellulose degradation,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC13159211/>
 16. A Comparative Analysis of the *Eisenia fetida* Gut and Compost Matrix,
<https://www.mdpi.com/2036-7481/16/8/177>
 17. (PDF) Comparative studies on the digestive enzymes in the gut of,
https://www.researchgate.net/publication/372492684_Comparative_studies_on_the_digestive_enzymes_in_the_gut_of_earthworms_Eudrilus_eugeniae_and_Eisenia_fetida
 18. Earthworms strongly modify microbial biomass and activity triggering,
https://www.researchgate.net/publication/6208854_Earthworms_strongly_modify_microbial_biomass_and_activity_triggering_enzymatic_activities_during_vermicomposting_independently_of_the_application_rates_of_pig_slurry
 19. Mechanisms and challenges in reducing antibiotic resistance genes,
<https://www.maxapress.com/article/doi/10.48130/biocontam-0025-0021>
 20. Changes in microbial biomass C, N, and P and enzyme activities in,
https://www.researchgate.net/publication/222181078_Changes_in_microbial_biomass_C_N_and_P_and_enzyme_activities_in_soil_incubated_with_the_earthworms_Metaphire_guillelmi_or_Eisenia_fetida
 21. Enzymatic dynamics into the *Eisenia fetida* (Savigny, 1826) gut,
https://www.researchgate.net/publication/320827565_Enzymatic_dynamics_into_the_Eisenia_fetida_Savigny_1826_gut_during_vermicomposting_of_coffee_husk_and_market_waste_in_a_tropical_environment
 22. Plant growth promoting bacteria drive food security - SciELO,
<https://www.scielo.br/j/bjb/a/8RkkPpSkWFtTHghcy8VHM9t/?lang=en>
 23. Microbial diversity of vermicompost bacteria that exhibit useful,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3725894/>
 24. Influence of Vermicompost on the Concentration of Exogenous,
https://www.researchgate.net/publication/381515581_Influence_of_Vermicompost_on_the_Concentration_of_Exogenous_Indole-3-Acetic_Acid_and_Its_Effect_on_the_Development_of_Tomato_Plants_Lycopersicum_esculentum_L
 25. Deciphering the enhanced translocation of Pb, Ni and Cd from,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11178750/>
 26. Role of plant growth promoting microorganisms in sustainable,
<https://dokumen.pub/role-of-plant-growth-promoting-microorganisms-in-sustainable-agriculture-and-nanotechnology-9780128170045-0128170042-9780128170052.html>
 27. Characterization of Plant Growth Promoting Rhizobacteria from,
https://www.researchgate.net/publication/399901158_Characterization_of_Plant_Growth_Promoting_Rhizobacteria_from_Dominant_Weeds_Inhabited_in_Kaligara_ng_River_Delta_Indonesia
 28. Silicon and plant growth promoting rhizobacteria differentially,
https://www.researchgate.net/publication/337698637_Silicon_and_plant_growth_promoting_rhizobacteria_differentially_regulate_AgNP-induced_toxicity_in_Brassica_juncea_Implication_of_nitric_oxide

29. Nutrient Recovery from Human Excreta for urban and ... - CGSpace,
<https://cgspace.cgiar.org/bitstreams/abdd6a74-f2e6-4d7c-aae6-4f6fb05fa0e1/download>
30. The Characterization of Feces and Urine: A Review of the Literature,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4500995/>
31. human feces compost as an organic fertilizer for sweet corn - AJFAND,
<https://ajfand.net/Volume23/No9/Gunawan20990.pdf>
32. nutrient composition of different organic manures and analysis,
https://www.researchgate.net/publication/353071857_NUTRIENT_COMPOSITION_OF_DIFFERENT_ORGANIC_MANURES_AND_ANALYSIS_METHODS
33. For 100 years, Milwaukee has been turning human waste into lawn fertiliser; 10.5 billion pounds diverted from landfills,
<https://timesofindia.indiatimes.com/world/us/for-100-years-milwaukee-has-been-turning-human-waste-into-lawn-fertiliser-10-5-billion-pounds-diverted-from-landfills/articleshow/133907835.cms>
34. Applied and Environmental Chemistry of Animal Manure: A Review,
https://www.researchgate.net/publication/309660882_Applied_and_Environmental_Chemistry_of_Animal_Manure_A_Review
35. ORGANIC FARMING :: Manures - TNAU Agritech Portal,
https://agritech.tnau.ac.in/org_farm/orgfarm_manure.html
36. Advances in Plant Microbiome and Sustainable Agriculture,
<https://dokumen.pub/advances-in-plant-microbiome-and-sustainable-agriculture-diversity-and-biotechnological-applications-1st-ed-9789811532078-9789811532085.html>
37. Biodiversity, and biotechnological contribution of beneficial soil,
https://www.researchgate.net/publication/351088468_Biodiversity_and_biotechnological_contribution_of_beneficial_soil_microbiomes_for_nutrient_cycling_plant_growth_improvement_and_nutrient_uptake
38. Microbial Dynamics and Functionality - ResearchGate,
https://www.researchgate.net/profile/Sudhir-Shekhar-2/publication/337309547_Environmental_Remediation_Microbial_and_Nonmicrobial_Prospects/links/5e39493b458515072d7d2cfd/Environmental-Remediation-Microbial-and-Nonmicrobial-Prospect.pdf
39. Factors Influencing Soil Nitrification Process and the Effect on,
<https://www.frontiersin.org/journals/sustainable-food-systems/articles/10.3389/fsufs.2022.821994/full>
40. Humic acids buffer the effects of urea on soil ammonia oxidizers and,
https://www.researchgate.net/publication/221768886_Humic_acids_buffer_the_effects_of_urea_on_soil_ammonia_oxidizers_and_potential_nitrification
41. Sustainable Intensification of Maize in the Industrial Revolution,
<https://www.frontiersin.org/journals/sustainable-food-systems/articles/10.3389/fsufs.2022.827477/full>
42. Nitrogen mineralization kinetics in soil supplemented with organic,
https://www.researchgate.net/publication/374743439_Nitrogen_mineralization_kinetics_in_soil_supplemented_with_organic_amendments

43. Vermicompost enrichment using organic wastes: Nitrogen content,
https://www.researchgate.net/publication/351194506_Vermicompost_enrichment_using_organic_wastes_Nitrogen_content_and_mineralization
44. Nitrogen Dynamics from Conventional Organic Manures as ... - MDPI,
<https://www.mdpi.com/2504-3129/5/3/49>
45. A comparative study of phosphorus recovery as struvite from cow,
https://www.researchgate.net/publication/351697187_A_comparative_study_of_phosphorus_recovery_as_struvite_from_cow_and_human_urine
46. Biochar seeding promotes struvite formation, but accelerates heavy,
https://www.researchgate.net/publication/328451513_Biochar_seeding_promotes_struvite_formation_but_accelerates_heavy_metal_accumulation
47. The Optimal Cyanobacterial Sludge Incorporation Balances Nutrient,
<https://www.mdpi.com/2073-4395/15/10/2429>
48. Biodiversity and functional role of Potassium Solubilizing Bacteria,
[https://ciasjournal.com/articles/Biodiversity-and-functional-role-of-Potassium-Solubilizing-Bacteria-\(KSB\)-in-sustainable-crop-production-A-review](https://ciasjournal.com/articles/Biodiversity-and-functional-role-of-Potassium-Solubilizing-Bacteria-(KSB)-in-sustainable-crop-production-A-review)
49. Potash-Mobilizing Biofertilizers: Complete Guide - BulkAgroChem,
<https://bulkagrochem.com/potash-mobilizing-biofertilizers>
50. Influence of potassium solubilizing bacteria on crop productivity and,
<https://academicjournals.org/journal/AJAR/article-full-text-pdf/15EA84831760>
51. The use of soil microbial potassium solubilizers in ... - Academia.edu,
https://www.academia.edu/99689261/The_use_of_soil_microbial_potassium_solubilizers_in_potassium_nutrient_availability_in_soil_and_its_dynamics
52. (PDF) Potassium Solubilizing Microorganisms: A New PGPM,
https://www.researchgate.net/publication/401134243_Potassium_Solubilizing_Microorganisms_A_New_PGPM
53. Methods Used for Artificial Humification of Animal Wastes,
<https://encyclopedia.pub/entry/52867>
54. Progress in Additives That Promote Humification During Agricultural,
<https://www.mdpi.com/2311-5637/12/8/392>
55. Optimizing the Cow Manure-Straw Ratio to Promote Organic Matter,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12987250/>
56. Study of the Humification Process and Humic Acid-like Structure,
<https://www.mdpi.com/2073-4395/13/2/465>
57. Humic acid-like fractions in raw and vermicomposted winery and,
https://www.researchgate.net/publication/229417264_Humic_acid-like_fractions_in_raw_and_vermicomposted_winery_and_distillery_wastes
58. Spectroscopic characterization of organic matter transformation,
https://www.researchgate.net/publication/343172723_Spectroscopic_characterization_of_organic_matter_transformation_during_composting_of_textile_solid_waste_using_UV-Visible_spectroscopy_Infrared_spectroscopy_and_X-ray_diffraction_XRD
59. Study on Molecular Structure Model and Visualization of Humic,
https://www.researchgate.net/publication/360908604_Study_on_Molecular_Structure_Model_and_Visualization_of_Humic_Acids_in_Different_Soils_Based_on_Sta

tistics

60. Advanced Characterization of Organic Matter Decaying during,
https://www.researchgate.net/publication/353686530_Advanced_Characterization_of_Organic_Matter_Decaying_during_Composting_of_Industrial_Waste_Using_Spectral_Methods
61. The Evaluation of Compost Maturity and Ammonium Toxicity Using,
<https://www.mdpi.com/2073-4395/14/11/2636>
62. Evaluation of Phytotoxicity of Organic Acids by Germination Test,
<https://hrcak.srce.hr/file/503286>
63. (PDF) Reduction of compost phytotoxicity during the process of,
https://www.researchgate.net/publication/42344659_Reduction_of_compost_phytoxicity_during_the_process_of_decomposition
64. Evaluation of Compost Maturity and Ammonium Toxicity Using,
<https://www.preprints.org/manuscript/202407.1148>
65. Occurrence and survival of viruses in composted human faeces,
https://www2.mst.dk/udgiv/publications/2003/87-7972-715-8/html/helepubl_eng.htm
66. A Review of the Literature on the Occurrence and Survival of,
https://www.researchgate.net/publication/266583967_A_Review_of_the_Literature_on_the_Occurrence_and_Survival_of_Pathogens_of_Animals_and_Humans_in_Garden_Compost
67. Sanitary Parasitology - IRIS,
<https://iris.who.int/bitstreams/f0b9c45d-fa1d-498a-8185-4f9d353a0f2e/download>
68. Salmonella regrowth in compost as influenced by substrate,
<https://pubmed.ncbi.nlm.nih.gov/24202718/>
69. Occurrence, growth, and suppression of salmonellae in composted,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC291765/>
70. (PDF) Use of Additives in Composting Promotes Passivation and,
https://www.researchgate.net/publication/376246923_Use_of_Additives_in_Composting_Promotes_Passivation_and_Reduction_in_Bioavailability_of_Heavy_Metals_HMs_in_Compost
71. Speciation Analysis Method of Heavy Metals in Organic Fertilizers,
https://www.researchgate.net/publication/366315654_Speciation_Analysis_Method_of_Heavy_Metals_in_Organic_Fertilizers_A_Review
72. Effects of substrate types on the transformation of heavy metal,
https://www.researchgate.net/publication/345126229_Effects_of_substrate_types_on_the_transformation_of_heavy_metal_speciation_and_bioavailability_in_an_aerobic_digestion_system
73. Composting as a Sustainable Solution for the Management of Plant,
<https://www.mdpi.com/2071-1050/16/22/9940>
74. Application of Rice Husk Biochar and Earthworm on Concentration,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9603306/>
75. Comparative assessment of heavy metals content during the,
https://www.researchgate.net/publication/272843662_Comparative_assessment_of_heavy_metals_content_during_the_composting_and_vermicomposting_of_Municipal_Solid_Waste

[cipal_Solid_Waste_employing_Eudrilus_eugeniae](#)

76. The effect of different additives on the immobilization of heavy,
<https://journals.agh.edu.pl/geol/article/download/7351/3371/40257>
77. Full article: Effects of adding biochar on the preservation of nitrogen,
<https://www.tandfonline.com/doi/full/10.1080/10962247.2022.2095055>
78. Effect of Biochar Addition on Mechanism of Heavy Metal Migration,
<https://www.mdpi.com/2311-5637/8/10/523>
79. Humus Soil Inhibits Antibiotic Resistance Gene Rebound in Swine,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11944299/>
80. Biochar can mitigate co-selection and control antibiotic resistant,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9160353/>
81. The Fate of Antibiotic Resistance Genes and Their Influential Factors,
<https://www.mdpi.com/2305-6304/14/5/428>
82. Effect of Extending High-Temperature Duration on ARG Rebound in,
<https://www.mdpi.com/2071-1050/16/13/5284>
83. Compostable Products in Aotearoa New Zealand - Bioplastics NEWS,
<https://bioplasticsnews.com/wp-content/uploads/2024/07/compostable-products-in-aotearoa-nz.pdf>
84. Microplastics in ecosystems: ecotoxicological threats and strategies,
<https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2025.1672484/full>
85. Real-time quantification of mcrA, pmoA for methanogen ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/21488508/>
86. Forced Aeration Composting,
<https://compostsystems.com/forced-aeration-composting/>
87. Mitigating Nitrous Oxide Emissions from Agricultural Soils with Biochar,
<https://www.mdpi.com/2073-4395/15/5/1115>
88. Relationships between different types of biochar and N₂O emissions,
https://www.researchgate.net/publication/376620234_Relationships_between_different_types_of_biochar_and_N2O_emissions_during_composting_based_on_roles_of_nosZ-carrying_denitrifying_bacterial_communities_enriched_on_compost_and_biochar_particles
89. Biochar mitigates the N₂O emissions from acidic soil by increasing,
<https://pubmed.ncbi.nlm.nih.gov/32063300/>
90. Biochar alters nitrogen transformations but has minimal effects on,
<https://www.research-collection.ethz.ch/server/api/core/bitstreams/0a845369-c38c-4d74-99aa-8dc978db6d38/content>
91. Linking N₂O emissions and nosZ gene abundance: A meta-analysis,
https://www.researchgate.net/publication/392496090_Linking_N2O_emissions_and_nosZ_gene_abundance_A_meta-analysis_of_organic_carbon_amendments_in_agricultural_soils