

Advanced Agronomic and Post-Harvest Dynamics of *Cicer arietinum* and *Phaseolus vulgaris* in the Temperate Berlin-Brandenburg Climate

Introduction to the Geo-Climatic and Pedological Context

The architectural design of a sustainable, high-yielding crop rotation in Central Europe must be inextricably linked to the localized climatic boundaries and pedological realities of the target geography. The cultivation of high-protein pulse crops, specifically the garbanzo bean or chickpea (*Cicer arietinum* L.) and the black bean, a morphological variant of the common bean (*Phaseolus vulgaris* L.), represents a central pillar of the transition toward closed-loop agricultural systems¹. While both species belong to the Fabaceae family and share the capacity for symbiotic biological nitrogen fixation, they exhibit highly divergent evolutionary adaptations, physiological temperature thresholds, and harvest mechanics².

The Berlin and wider Brandenburg region presents a highly specific, moderately challenging agronomic environment. Situated within USDA Hardiness Zone 7b, the region experiences average annual extreme minimum temperatures ranging between $-15^{\circ}C$ and $-12.2^{\circ}C$, dictating that warm-season legumes must be strictly cultivated as summer annuals¹. Historical meteorological data indicates that the last spring air frost ($0^{\circ}C$) typically occurs around May 8, while the first autumn air frost arrives near September 22, delineating a reliable frost-free window of approximately 136 to 137 days¹. When observing light ground frost ($-1^{\circ}C$), the agricultural window expands from April 26 to October 6, offering up to 163 operational days for resilient crops¹.

Frost Threshold	Last Spring Occurrence	First Autumn Occurrence	Resulting Frost-Free Days
Air Frost ($0^{\circ}C$)	May 8	September 22	~ 137 days ¹
Light Ground Frost ($-1^{\circ}C$)	April 26	October 6	~ 163 days ¹

Hard Ground Frost (−2° C)	April 5	October 28	~ 206 days ¹
-------------------------------	---------	------------	-------------------------

The defining limitation of the Brandenburg agricultural landscape is its pedology. The region is dominated by Pleistocene-derived glacial sands and sandy loams (Arenosols), frequently exhibiting critically low agricultural yield indices, commonly rated between 25 and 40 Bodenkunde¹. These coarse-textured soils are characterized by an abundance of macropores, which ensures excellent aeration and ease of mechanical cultivation under all moisture regimes, but simultaneously results in an exceptionally low Available Water Capacity¹. Consequently, these soils are highly susceptible to acute drought stress during the summer months and suffer from severe vulnerability to the downward leaching of water-soluble macronutrients during winter precipitation events¹. Compounding this physical limitation is the long-term decline in localized groundwater recharge, which has dropped significantly by up to 40% since 1980 due to shifting precipitation patterns and increased evapotranspiration¹.

To achieve agronomic stability and the preservation of soil organic carbon in this fragile environment, the implementation of a rigorously structured crop rotation is mandatory. In sandy Arenosols where native organic matter degrades rapidly due to high microbial oxidation rates and high oxygen diffusion, the rotation must be calibrated using the Humus Balance framework established by the Association of German Agricultural Analytic and Research Institutes (VDLUFA)¹. Grain legumes, such as chickpeas and beans, contribute approximately +160 kg Humus-C/ha, serving as vital restorative phases between aggressive, humus-depleting heavy feeders like winter brassicas or potatoes¹.

Thermal Kinetics and Growing Degree Days (GDD)

Modeling

The temporal sequencing of crop development from seedling emergence to physiological maturity is governed by the accumulation of thermal energy, quantified as Growing Degree Days (GDD). The mathematical model for GDD assumes that the plant development rate is directly proportional to temperature within a specific linear range, bounded by a base

temperature (T_{base}) below which development ceases, and an upper temperature (T_{upper}) where heat stress inhibits metabolism⁵. The standard formulation is expressed as:

$$GDD = \sum \left(\frac{T_{max} + T_{min}}{2} \right) - T_{base}$$

The calculation of thermal time highlights the divergent evolutionary origins and climatic requirements of chickpeas and black beans.

Cicer arietinum evolved in the Fertile Crescent and is genetically adapted as a cool-season

crop⁷. It utilizes a lower T_{base} of $5^{\circ}C$ for phenological modeling⁹. Modern chickpea varieties require between 1,287 and 1,612 GDD (base $5^{\circ}C$) to reach physiological maturity⁹. Studies conducted at similar European latitudes (52°N) demonstrate that the grain-filling GDD requirement is remarkably stable across genotypes, typically demanding 220 to 227 GDD⁹. The long-term April–September thermal sum in temperate European climates averages approximately 1,480 GDD (base $5^{\circ}C$), comfortably accommodating early-to-medium maturing chickpea genotypes while leaving a safety margin of roughly 200 GDD before autumn frosts threaten the harvest⁹. Modern desi and kabuli types possess moderate cold tolerance, capable of surviving brief frosts to $-2^{\circ}C$, allowing for early spring sowing strategies⁹. Conversely, *Phaseolus vulgaris* originated in the warm, neotropical regions of Mesoamerica and the Andes, operating as an obligate warm-season crop³. Black beans utilize a higher T_{base} of $10^{\circ}C$ for agronomic modeling⁵. They are exquisitely sensitive to chilling injury; soil temperatures must consistently exceed $15.5^{\circ}C$ to $18^{\circ}C$ to prevent seed rot by soil-borne oomycetes and to ensure uniform hypocotyl emergence¹⁴. Sowing in Berlin must consequently be delayed until late May, capitalizing on rapidly warming soils¹⁴.

Phenological Parameter	Chickpea (<i>Cicer arietinum</i>)	Black Bean (<i>Phaseolus vulgaris</i>)
Base Temperature (T_{base})	$5^{\circ}C$ [cite: 9]	$10^{\circ}C$ [cite: 11, 12]
Optimal Germination Temp.	$10^{\circ}C$ –	$15.5^{\circ}C$ – [cite: 14, 15]
Target Sowing Window (Berlin)	Late April - Early May ¹⁶	Mid to Late May ¹⁴
GDD to Maturity	1, 287 – (base $5^{\circ}C$) ⁹	Cultivar dependent (base $10^{\circ}C$)
Frost Tolerance (Vegetative)	Survives brief exposure to $-2^{\circ}C$ [cite: 9]	Lethal tissue damage at $0^{\circ}C$ [cite: 14, 15]

High-Temperature Stress and Reproductive Microsporogenesis

While the Berlin climate presents frost risks at the margins of the season, mid-summer heatwaves present a severe physiological threat during the critical reproductive phases of both crops. Although black beans require warm soils for germination, they are paradoxically highly sensitive to supra-optimal atmospheric temperatures during anthesis.

When ambient temperatures exceed 30°C —a threshold increasingly breached during Brandenburg summers—*Phaseolus vulgaris* experiences catastrophic reproductive failure, with total seed weight frequently decreasing by 50% to 71% in heat-sensitive accessions¹⁷.

High-temperature stress disrupts microsporogenesis (pollen development) by inducing the premature degeneration of the tapetal cell layer within the anther¹⁸. This tapetal collapse starves the developing microspores, leading to anther indehiscence and the production of

sterile, aborted pollen¹⁸. Furthermore, temperatures exceeding 32°C during the day and 27°C at night inhibit pollen tube elongation and impair stigma receptivity¹⁸. The resulting

fertilization failure triggers massive flower abscission and the abortion of developing ovules¹⁸. To mitigate this, agronomists must utilize fast-maturing cultivars that complete anthesis before the absolute peak heat of late July, or select specifically bred heat-tolerant accessions that maintain pollen viability under thermal load¹⁸.

Chickpeas similarly suffer under terminal heat stress. While vegetative growth can tolerate high

temperatures, exposure to temperatures exceeding 30°C during the pod-filling stage accelerates senescence and truncates the grain-filling duration, resulting in reduced seed size and diminished total yield¹⁷. Early sowing strategies in April are therefore highly recommended for chickpeas to ensure that the reproductive phase coincides with the cooler temperatures of early summer, maximizing the photothermal quotient and ensuring optimal sink-source assimilate partitioning²¹.

Pedology, pH Management, and Aluminum Toxicity Mitigation

The optimization of the rhizosphere pH is the paramount physiochemical intervention required in the Berlin-Brandenburg Arenosols. The pH directly dictates the bioavailability of essential mineral nutrients and governs the metabolic efficacy of the symbiotic microbiome²².

Chickpeas exhibit optimal physiological performance in well-drained, friable loams with a neutral to slightly alkaline pH, ideally ranging between 6.0 and 8.0⁷. The high macroporosity of Berlin's sandy soils is highly advantageous for *Cicer arietinum*, as the species is acutely sensitive to hypoxic waterlogging⁷. Transient anaerobic conditions in the root zone physically suffocate the obligate aerobic rhizobia, depress the activity of the nitrogenase enzyme complex, and drastically increase susceptibility to soil-borne vascular wilt pathogens,

particularly *Fusarium oxysporum* f. sp. *ciceris*⁷.

Black beans demonstrate a narrower and slightly lower optimal pH window, preferring mildly acidic to neutral soils ranging from pH 5.8 to 6.5¹⁴. However, the native pH of poorly buffered glacial sands in the Brandenburg region can frequently drop below 5.5 due to the continuous leaching of basic cations (Ca^{2+} , Mg^{2+} , K^{+}) driven by winter precipitation¹. When the soil pH falls below 5.5, structural aluminosilicate minerals in the clay fraction undergo rapid acid hydrolysis, releasing highly phytotoxic trivalent aluminum cations (Al^{3+}) into the soil solution²⁵.

Aluminum toxicity is catastrophic for *Phaseolus vulgaris* and *Cicer arietinum*. The Al^{3+} cations physically bind to the negatively charged pectins in the root apex, instantly ceasing cell division and elongation²⁶. This results in stunted, coralloid root systems that are entirely incapable of scavenging deep moisture reserves during summer droughts²⁶. Furthermore, high Al^{3+} activity paralyzes the chemotactic flavonoid signaling between the legume root hairs and the rhizobial symbionts, effectively preventing nodulation and biological nitrogen fixation²⁴. To mobilize bound phosphorus and counteract these toxicities, legume roots actively exude low-molecular-weight organic acids, predominantly citrate²⁶. Citrate functions as a potent biological chelating agent; it replaces phosphorus adsorbed on the surface of iron and aluminum oxides, complexing with Al^{3+} to reduce its phytotoxicity while simultaneously increasing the bioavailability of phosphate ions in the rhizosphere²⁶. To further ameliorate acidic Arenosols, the incorporation of engineered biochar is highly recommended. Biochar physically adsorbs labile Al^{3+} , buffers the localized rhizosphere pH, and permanently elevates the cation exchange capacity of the sandy matrix, providing a hospitable microenvironment for root expansion and rhizobial proliferation²⁹.

Microbiome Engineering and Biological Nitrogen Fixation (BNF)

The profound agronomic value of cultivating grain legumes lies in their capacity to execute biological nitrogen fixation, an energetically expensive biochemical process that converts inert atmospheric dinitrogen (N_2) into bioavailable ammonia (NH_3)⁴. This reaction is catalyzed by the nitrogenase enzyme complex—a highly oxygen-sensitive metalloenzyme composed of a molybdenum-iron (MoFe) dinitrogenase and an iron (Fe) dinitrogenase reductase⁴. Because the Haber-Bosch process requires immense fossil fuel inputs (temperatures of $400^{\circ}C$ and pressures of 200 bar), maximizing BNF is essential for sustainable agriculture⁴. Because *C. arietinum* and *P. vulgaris* have divergent evolutionary histories, they exhibit strict host-specificity regarding their microbial symbionts. Chickpeas exclusively form nodules with the genus *Mesorhizobium*, predominantly *Mesorhizobium ciceri* and *M. mediterraneum*⁸. Elite

strains, such as *M. ciceri* CC1192, contain specialized integrative and conjugative elements (ICEs), such as ICEMcSym1192, which harbor the *nif* and *fix* gene clusters required for highly efficient nitrogen fixation³⁷. These genetic elements encode for the high-oxygen-affinity *cbb₃*-type cytochrome oxidase, allowing the bacteria to respire in the microaerobic environment of the root nodule while protecting the delicate nitrogenase enzyme from oxygen degradation³⁸. Conversely, black beans partner specifically with *Rhizobium leguminosarum* bv. *phaseoli*²⁴. Because native European soils lack endemic populations of these highly specific rhizobia, artificial seed inoculation prior to sowing is an absolute requirement to achieve nitrogen self-sufficiency¹⁶. The formulation and application methodology of the inoculant critically impact rhizobial survival. While peat-based slurries coated directly onto the seed are a historical standard, direct exposure to desiccation, ultraviolet radiation, and acidic seed exudates can rapidly decimate the bacterial population⁴³. Furthermore, the concurrent application of fungicidal seed treatments (such as thiram-based compounds) is highly toxic to *Mesorhizobium*, leading to catastrophic failures in nodulation⁴⁵. Advanced agronomic systems increasingly utilize liquid or granular inoculants injected directly into the furrow below the seedbed. This spatial separation physically shields the fragile bacteria from toxic seed treatments and harsh surface conditions, maximizing nodule occupancy rates and boosting nitrogen uptake by up to 19.4% in sandy soils⁴¹. It is also critical to avoid mixing rhizobial inoculants with liquid micronutrient fertilizers. High concentrations of copper sulfate ($CuSO_4$) and zinc sulfate ($ZnSO_4$) exhibit acute bactericidal toxicity; exposing *M. ciceri* to 200 $\mu\text{mol/L}$ $CuSO_4$ causes populations to crash below detection limits within minutes, underscoring the need to physically separate inoculants from in-furrow mineral fertilizers⁴³.

Spatial Geometry and Sowing Mechanics

Plant architecture, canopy closure dynamics, and localized humidity microclimates dictate the optimal spatial geometry for both pulse species. Achieving the correct plant density is a delicate thermodynamic balance; dense canopies suppress weeds but trap humidity, escalating disease risk, while sparse canopies waste photosynthetic radiation and allow aggressive weed proliferation¹⁴.

Chickpea Sowing Dynamics

Chickpeas present a semi-erect, branching morphology¹⁶. In the Berlin climate, early spring sowing is targeted as soon as soil temperatures reach 10°C to 15°C ¹⁶. Seeds are precision-drilled to a depth of 2.5 to 4.0 cm, adjusting slightly deeper in coarse, sandy beds to ensure capillary moisture contact, as shallow seeds rapidly desiccate and stall germination¹⁶. The targeted planting density for medium-seeded "Desi" type chickpeas (test weight of 21-23 g per 100 seeds) is highly specific⁴⁸. Agronomic trials consistently demonstrate that while high seeding rates (100 to 120 kg/ha) accelerate early canopy closure, they induce intense interplant competition for the limited moisture available in Arenosols, resulting in poor grain filling, reduced pod numbers, and lower net economic returns⁴⁸. The optimal seed rate is precisely

calibrated to 80 to 84 kg/ha, targeting a final plant stand of 33 to 44 seedlings per square meter⁴⁸. This architecture is achieved via an in-row spacing of 8 to 15 cm and an inter-row spacing of 45 to 60 cm¹⁶. Utilizing twin rows on a 76 cm bed is highly effective, perfectly balancing root architectural expansion with the spatial requirements for mechanical weed cultivation¹⁶.

Black Bean Sowing Dynamics

Black beans demand significantly warmer germination conditions. Sowing in Berlin must be delayed until mid-to-late May, ensuring the absolute avoidance of late spring air frosts and confirming soil temperatures steadily exceed $15.5^{\circ}C$ at a 5 cm depth¹⁴. Planting into cold soils ($< 10^{\circ}C$) results in imbibitional chilling injury, where the rapid influx of cold water disrupts cellular membrane reorganization, leading to electrolyte leakage, rotting, and catastrophic stand failure¹⁴.

To optimize light interception while maintaining sufficient airflow to prevent foliar pathogens, black bean seeds are planted 2.5 to 5.0 cm deep¹⁴. The optimal geometry utilizes a tight in-row spacing of 5 to 8 cm within rows spaced 45 to 60 cm apart¹⁴. Because *Phaseolus vulgaris* is exceptionally sensitive to weed competition during its early vegetative stages, maintaining uniform inter-row spacing is an absolute structural necessity to facilitate precision mechanical weed control⁵².

Parameter	Chickpea (<i>Cicer arietinum</i>)	Black Bean (<i>Phaseolus vulgaris</i>)
Sowing Depth	2.5 - 4.0 cm ¹⁶	2.5 - 5.0 cm ¹⁴
In-Row Spacing	8 - 15 cm ¹⁶	5 - 8 cm ¹⁴
Inter-Row Spacing	45 - 60 cm ¹⁶	45 - 60 cm ¹⁴
Target Plant Stand	33 — seedlings/ m^2 [cite: 50]	Cultivar dependent (bush vs. vine)
Optimal Seed Rate	80 — kg/ha ⁴⁸	Varies by seed size

Weed Competition and Mechanical Cultivation

In the absence of broad-spectrum synthetic herbicides—particularly under organic management systems prevalent in the EU—weed control relies entirely on mechanical disruption and canopy competition⁵⁴. The Critical Period of Weed Control (CPWC) defines the exact temporal window during which the crop must remain weed-free to prevent unacceptable

yield loss.

For *Phaseolus vulgaris*, weed interference models based on Gompertz and logistic equations indicate that the CPWC begins at 50 GDD (approximately 11 days after emergence) and ends at 284 GDD (approximately 28 days after emergence)¹³. If aggressive weed populations are permitted to compete for the entirety of the growing season, fresh pod and dry seed yields are depressed by up to 60% to 65%¹³. Even highly localized competition from volunteer crops, such as corn or cereals from previous rotations, exerts massive penalties; dry bean yield losses increase by 1.2% to 1.8% for every 100 GDD that volunteer weed control is delayed¹¹.

To manage weeds during this highly vulnerable phase without damaging the delicate legume hypocotyls, blind cultivation techniques are deployed⁵⁶. Using a rotary hoe, the soil surface is aggressively disturbed just prior to or immediately following bean emergence, physically uprooting small, white-thread stage weeds while the deeply rooted legume seeds remain secure⁵³. Once the beans establish their first true leaves, precision inter-row cultivation via specialized sweeps or tines is utilized to slice weed roots between the 45 to 60 cm rows, ceasing operation once the legume canopy begins to close to prevent mechanical damage to the flowering branches⁵².

Phytopathological Constraints and Epidemiological Modeling

The successful maturation of grain legumes requires navigating devastating fungal pathogens, the life cycles of which are intrinsically linked to the localized climatic parameters of Brandenburg.

Ascochyta Blight in Chickpeas

The paramount biological threat to chickpea cultivation globally is Ascochyta blight, a highly destructive foliar disease caused by the necrotrophic fungus *Ascochyta rabiei*⁵⁸. The pathogen survives the winter on infested crop residues on the soil surface⁵⁸. In the spring, as

temperatures approach 20°C and relative humidity exceeds 95%, pycnidia absorb moisture and exude millions of conidia enclosed in a mucilaginous cirrus⁵⁸.

The primary vector for disease transmission is rain-splash dispersal⁶⁰. Epidemiological modeling and open-channel rainfall simulator studies demonstrate that the kinetic energy of falling raindrops physically ruptures this cirrus, aerosolizing the spore-laden droplets and propelling them upward into the chickpea canopy⁵⁸. While historical models relied on a coarse 2.0 mm daily rainfall threshold to predict infection events, advanced segmented regression analyses reveal that rain-splash dispersal is highly sensitive to micro-precipitation, exhibiting a critical breakpoint at an event rainfall of just 1.1 mm⁶². Once this 1.1 mm threshold is exceeded, the high frequency of droplet impacts drastically increases the trajectory and volume of secondary splash droplets, raising infection probabilities across the upper canopy and facilitating multi-step dispersal between rows⁶².

Infection requires a leaf wetness duration of at least 6 to 10 hours⁵⁸. In dense canopies typical of excessive seeding rates, restricted airflow prevents the evaporation of morning dew,

allowing the conidia to germinate and rapidly girdle stems and pods, frequently resulting in 100% yield loss in susceptible cultivars⁵⁸. Maintaining wide inter-row spacing (45–60 cm) is an architectural necessity to facilitate rapid canopy drying, physically disrupting the pathogen's required environmental parameters¹⁶.

Soil-Borne Pathogens and Rotational Hygiene

To suppress the buildup of soil-borne inoculum, strict temporal crop rotation intervals are legally and agronomically mandated¹. *Rhizoctonia solani*, a devastating basidiomycete fungus that causes severe stem canker and root rot in legumes and potatoes, thrives in compacted, structurally degraded sandy soils¹. Furthermore, if brassica cover crops (such as white mustard) are utilized inappropriately within the rotation, they serve as asymptomatic multiplication vectors for *Plasmodiophora brassicae* (clubroot), necessitating strict 4 to 6-year cultivation pauses between susceptible botanical families to ensure soil hygiene¹. The inclusion of chickpeas and black beans, coupled with non-cruciferous cover crops like phacelia and bristle oat, serves to starve these specialized pathogens through temporal isolation¹.

Maturation, Desiccation, and the Genetics of Pod Shattering

As the legumes transition into late summer, the focus shifts from primary biomass accumulation to the physiological desiccation of the seed and the preservation of mechanical pod integrity. A fundamental domestication syndrome distinguishing modern grain legumes from their wild progenitors is the loss of natural seed dispersal, commonly known as pod shattering or dehiscence⁶⁶.

The Biomechanics and Genetics of Pod Shattering

Pod dehiscence is a highly calibrated biomechanical process driven by differential cellular desiccation. As the pod wall dries, variations in the cellular water content and the structural rigidity of varying tissues create extreme physical tension⁶⁹. Specifically, the pod valves are connected along the dorsal and ventral sutures by specialized dehiscence zones containing vascular bundles⁶⁹. During maturation, the "endocarp b" layer undergoes heavy lignification, developing thickened secondary cell walls, while the adjacent "endocarp a" layer undergoes programmed cell death and remains non-lignified⁶⁹. The differential contraction of these tissue layers upon atmospheric drying creates a massive twisting force. When this force overcomes the mechanical load limit of the dehiscence zone, the pod explosively shatters, scattering the seed⁶⁹.

In both *Phaseolus* and *Cicer*, modern genomics has elucidated the underlying genetic architecture governing this trait. The *POD DEHISCENCE 1 (PDH1)* gene, which encodes a dirigent-like protein responsible for regulating precise lignin deposition in the sclerenchyma and fiber cap cells, plays a central evolutionary role⁶⁷. Furthermore, quantitative trait loci (QTLs) such as the *PvMYB26* locus in common beans dictate the formation of the abscission layer in the ventral suture; non-shattering domesticates possess mutations that render this abscission layer non-functional¹⁰. Despite thousands of years of selective breeding for indehiscence, dry bean classes retain a dangerous phenotypic susceptibility to pod shattering, particularly in the

semi-arid conditions of a hot Brandenburg August, where rapid atmospheric drying renders the pods highly brittle⁶⁶.

Chemical Desiccation vs. European Regulatory Reality

In conventional, large-scale agriculture globally, chemical desiccation is frequently employed to kill the foliage, force uniform crop dry-down, and eliminate late-season weeds, typically utilizing systemic herbicides like glyphosate or contact desiccants like diquat and saflufenacil⁷⁴.

However, within the European Union, the regulatory landscape strictly dictates harvest protocols.

The application of glyphosate as a pre-harvest desiccant results in the systemic translocation of the chemical directly into the maturing seed, locking high concentrations of the residue and its primary metabolite, AMPA, into the cotyledons⁷⁴. Aggressive consumer pushback and strict Maximum Residue Limits (MRLs) established by the European Commission render pre-harvest chemical desiccation a high-risk practice for pulses destined for human consumption, frequently resulting in entire shipments being rejected at market borders⁷⁷. Furthermore, under strict EU Organic regulations (Regulation (EU) 2018/848), synthetic chemical desiccants are absolutely forbidden; any detection of residues triggers immediate decertification⁷⁹.

Consequently, European producers must rely entirely on physical maturation and mechanical field-curing processes to prepare the crop for storage.

Mechanical Curing and Drying Processes on the Vine

Because chemical desiccation is heavily restricted, and because leaving the crop to dry perfectly standing in the field invites catastrophic pod shatter losses from autumn wind events, the traditional and highly effective method of vine-curing is utilized.

Identifying Physiological Maturity

The decision to initiate physical harvest procedures relies on precise visual and physiological cues. Harvesting via undercutting or swathings should commence when the crop has reached 80% to 90% leaf drop, the stems are transitioning from green to brown, and approximately 80% of the pods have changed from green to a dry, "buckskin" or tan color⁶⁴. At this stage, the seeds have reached physiological maturity—defined as the point of maximum dry matter accumulation—and possess an internal moisture content of roughly 30%⁶⁴. If a seed is extracted and split open, it should exhibit zero internal green coloration, indicating the complete degradation of chlorophyll⁷⁷.

The Mechanics of Vine-Curing

Harvesting begins by physically severing the root system from the soil utilizing specialized undercutting blades, or by hand-pulling the entire plant¹⁶. The plants are then laid into windrows upon the soil surface or, to guarantee premium market quality and protect against autumn rains, gathered and hung on racks or mesh screens in a dry, well-ventilated, covered space¹⁶. This curing process, extending for 7 to 14 days depending on ambient humidity, is not merely a mechanism for moisture reduction; it is a vital physiological transition¹⁶.

1. **Assimilate Translocation:** By leaving the pods attached to the intact vine during the initial drying phase, residual complex carbohydrates and vital nutrients located in the

vascular tissues of the stem continue to slowly translocate into the seed, maximizing ultimate seed weight and nutritional density.

- 2. **Moisture Equilibration:** Natural atmospheric curing forces a slow, highly synchronized reduction in moisture. This prevents the rapid, uneven contraction of the cotyledons and the delicate seed coat that frequently occurs during artificial thermal drying⁸⁵. Rapid thermal gradients lead to irreversible seed coat cracking (Mechanical Damage Susceptibility), a severe cosmetic defect that critically downgrades the crop for canning markets⁶⁴.
- 3. **Pathogen Evasion:** Elevating the vines off the soil surface onto racks prevents the maturing pods from coming into physical contact with wet soil (earth-tag) or latent soil-borne fungi, entirely avoiding the rapid onset of white mold or pod rot that occurs if windrows are subjected to unpredictable late-summer rainfall⁶⁴.

Threshing Dynamics

Once the atmospheric curing process successfully lowers the internal seed moisture content to a critical window between 16% and 22%, the crop is ready for mechanical threshing⁶⁴. Attempting to process the beans below 14% moisture is highly dangerous, as the ultra-dry cotyledons become excessively brittle; the mechanical impact of the combine's steel components will cause massive internal splitting, transverse micro-fractures, and seed coat shattering⁶⁴. Conversely, threshing above 22% results in mashed seeds and frequently gums up the separation sieves.

To preserve the structural integrity of the beans, the combine harvester must be meticulously calibrated. The threshing cylinder speed must be reduced to the lowest functional limit, typically between 150 and 300 revolutions per minute (rpm), ensuring the peripheral speed of the cylinder remains below 914 meters per minute⁸⁷. Simultaneously, the concave clearance must be opened widely (e.g., 5/16 to 3/4 inch, or approximately 1.6 cm) to allow the large, dense seeds to pass through the threshing mechanism without being crushed or abraded against the metal plates⁸⁷.

Threshing Parameter	Target Specification	Consequence of Deviation
Optimal Moisture	16% - 22% ⁶⁴	<14% causes shattering; >22% causes mashing ⁶⁴
Cylinder Speed	150 - 300 rpm ⁸⁸	High speeds induce severe seed coat cracking ⁸⁷
Concave Clearance	1.6 cm (5/16 - 3/4 inch) ⁸⁷	Narrow clearance crushes seed; wide leaves unthreshed pods

Post-Harvest Thermodynamics and Quality Preservation

The agronomic lifecycle does not conclude at threshing; the preservation of the harvested grain requires strict thermodynamic control. Legume seeds are highly hygroscopic biological entities that will continuously interact with ambient atmospheric humidity, absorbing or desorbing moisture until equilibrium is reached⁹².

Moisture and Water Activity (a_w)

For long-term storage spanning up to 12 months, the internal moisture content must be carefully brought down to a stable 13% to 14%¹⁶. The critical biochemical metric governing storage stability is water activity (a_w), which measures the thermodynamic ratio of unbound, free water available for microbial metabolism⁹⁴. If the a_w of the stored beans rises above 0.65, the environment becomes highly conducive to the proliferation of xerophilic molds, such as *Aspergillus* and *Penicillium* species⁹⁴. These resilient fungi synthesize dangerous secondary metabolites known as mycotoxins (e.g., aflatoxins and ochratoxins), which permanently contaminate the crop, surviving standard cooking temperatures and rendering the batch entirely unfit for human consumption⁹⁸.

Seed Coat Darkening and the Hard-To-Cook (HTC) Defect

The physical environment of the storage facility—specifically temperature and relative humidity—directly impacts the culinary and economic quality of the pulses. When chickpeas or black beans are subjected to elevated temperatures (consistently $> 25^{\circ}C$) and high relative humidity ($> 65\%$) for prolonged periods, they undergo severe physiological deterioration⁹². First, the enzymatic oxidation of proanthocyanidins and polyphenolic compounds located within the seed coat causes progressive seed coat darkening⁹³. This biochemical browning transforms highly valued, bright Kabuli chickpeas into an unmarketable, muddy brown, destroying their visual appeal for the premium canning and hummus markets⁹³. Second, and far more critically, these warm, humid storage conditions induce the Hard-to-Cook (HTC) defect⁹⁹. The HTC phenomenon is a complex physiochemical hardening of the seed driven by the secondary lignification of cotyledon cells and the enzymatic cross-linking of pectic substances in the middle lamella, frequently mediated by phytase activity releasing calcium and magnesium ions that form insoluble pectate bridges⁹⁹. This biochemical locking of the cell walls severely restricts water imbibition during post-harvest processing, requiring vastly extended boiling times to achieve cellular separation and edibility, resulting in massive energy costs for processors and profound consumer dissatisfaction⁹⁹. To prevent this debilitating defect, beans must be stored in cool, dark environments (optimally below $20^{\circ}C$) with strictly regulated ambient humidity, utilizing hermetically sealed packaging or actively aerated silos to maintain absolute dormancy⁹².

Conclusion

The successful cultivation of *Cicer arietinum* and *Phaseolus vulgaris* in the Berlin-Brandenburg region requires the precise synthesis of complex agronomic, physiological, and genetic sciences. The region's defining parameters—a constrained frost-free window and highly permeable, low-AWC Arenosols—dictate that crop phenology must be perfectly aligned with environmental realities via meticulous GDD modeling. By manipulating soil pH to mitigate catastrophic aluminum toxicity, leveraging highly specific *Mesorhizobium* and *Rhizobium* inoculants to ensure absolute nitrogen self-sufficiency, and optimizing spatial geometry to suppress weeds and disrupt the rain-splash dispersal physics of *Ascochyta rabiei*, agronomists can secure robust yields in a challenging landscape.

Crucially, the intersection of strict European regulatory frameworks forbidding chemical desiccation with the innate genetic susceptibility of these legumes to explosive pod shattering demands specialized harvest mechanics. The implementation of physical vine-curing processes—relying on slow, atmospheric moisture equilibration—is imperative not merely for regulatory compliance, but for the preservation of delicate seed coat integrity. Ultimately, mastering the thermodynamics of post-harvest storage to prevent water-activity-driven mycotoxin synthesis and the physiochemical Hard-to-Cook defect ensures that the resulting crop meets the exacting nutritional and cosmetic standards of the premium European pulse market.

Works cited

1. 24 Berlin Crop Rotation Schedule.pdf
2. Plant Resources of South-East Asia ^ 2^bi Sb - WUR eDepot, <https://edepot.wur.nl/281901>
3. Functional Genomics: From Soybean to Legume - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12249971/>
4. Exploiting Biological Nitrogen Fixation: A Route Towards a ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7464700/>
5. Base and upper temperature thresholds to support the calculation of, https://repositori.irta.cat/bitstream/20.500.12327/4862/1/Paredes_Base_2025.pdf
6. Modeling seedling emergence | Request PDF - ResearchGate, https://www.researchgate.net/publication/222677466_Modeling_seedling_emergence
7. Chickpea · AgricultureID, <https://agricultureid.com/crops/chickpea>
8. Rhizobial diversity is associated with inoculation history at a two, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9329089/>
9. Thermal time requirements for phenological development of, <https://www.agronomyjournals.com/archives/2024/vol7issue2/PartB/9-2-114-560.pdf>
10. Pod indehiscence in common bean is associated with the fine ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7921299/>
11. Volunteer Corn (Zea mays) Interference in Dry Edible Bean, <https://bioone.org/journals/weed-technology/volume-30/issue-4/WT-D-16-0003>

[7.1/Volunteer-Corn-Zea-mays-Interference-in-Dry-Edible-Bean-Phaseolus/10.1614/WT-D-16-00037.1.pdf](#)

12. Vegetable production course | PPT - Slideshare,
<https://www.slideshare.net/slideshow/vegetable-production-course/124702609>
13. The critical period for weed competition in French bean (Phaseolus,
https://www.researchgate.net/publication/251540027_The_critical_period_for_weed_competition_in_French_bean_Phaseolus_vulgaris_L_in_Mediterranean_areas
14. Growing Black Beans for a Healthy and Flavorful Harvest | Taim.io,
<https://www.taim.io/plants/how-to-grow-black-beans>
15. Bean Seed Production ver_1pt7,
https://www.savingourseeds.org/pubs/bean_seed_production_ver_1pt7.pdf
16. Growing Chickpeas: Your Simple Step-by-Step Garden Guide | Taim.io,
<https://www.taim.io/plants/growing-chickpeas-a-step-by-step-guide>
17. (PDF) Thermal Stress Impacts on Reproductive Development and,
https://www.researchgate.net/publication/328046935_Thermal_Stress_Impacts_on_Reproductive_Development_and_Grain_Yield_in_Grain_Legumes
18. Effects of high-temperature stress on microsporogenesis in heat,
https://www.researchgate.net/publication/227738773_Effects_of_high-temperature_stress_on_microsporogenesis_in_heat-sensitive_and_heat-tolerant_genotypes_of_Phaseolus_vulgaris
19. (PDF) Adaptation to High Temperature and Water Deficit in the,
https://www.researchgate.net/publication/258387397_Adaptation_to_High_Temperature_and_Water_Deficit_in_the_Common_Bean_Phaseolus_vulgaris_L_during_the_Reproductive_Period
20. Developing Climate-Resilient Chickpea Involving Physiological and,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7052492/>
21. Sowing Date Affects the Timing and Duration of Key Chickpea (Cicer,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7598726/>
22. Natural Farming Training Toolkit & Best Practices Guide - NITI Aayog,
<https://www.niti.gov.in/sites/default/files/2026-03/Training-Manual-English.pdf>
23. Rhizobium Species: Role in Plant Nutrition, Crop Quality, Soil,
<https://www.indogulfbioag.com/post/rhizobium-species-plant-nutrition>
24. Competitiveness and Nitrogen Fixation Efficiency Analysis of ... - MDPI,
<https://www.mdpi.com/2077-0472/15/16/1784>
25. Potential role of biochars in decreasing soil acidification - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/28063658/>
26. Root exudates contribute to belowground ecosystem hotspots - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9581264/>
27. Rhizobium-Legume Symbiosis and Nitrogen Fixation under Severe,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC98982/>
28. Exudation of organic acids in common bean as related to,
https://www.researchgate.net/publication/222689733_Exudation_of_organic_acids_in_common_bean_as_related_to_mobilization_of_aluminum-and_iron-bound_phosphates
29. "Assessing the potential of biochar and aged biochar to alleviate,

- <https://pubmed.ncbi.nlm.nih.gov/29890430/>
30. Aluminum toxicity in plants and its possible mitigation in acid soils by, <https://pubmed.ncbi.nlm.nih.gov/33092837/>
 31. Importance of soil amendments with biochar and/or Arbuscular, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8886007/>
 32. Symbiotic Effectiveness, Rhizosphere Competence and Nodule, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11902108/>
 33. Biological nitrogen fixation and prospects for ecological ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9133800/>
 34. symbiotic effectiveness of rhizobia from chickpea - N2Africa, <https://www.n2africa.org/sites/default/files/MScThesis%20Tadele%20Ereso.pdf>
 35. Symbiotic Effectiveness, Rhizosphere Competence and Nodule, <https://www.mdpi.com/2223-7747/14/5/809>
 36. Phenotypic and Genotypic Diversity Among Symbiotic and Non, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6759536/>
 37. Significance of Plant Growth Promoting Rhizobacteria in Grain, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7698556/>
 38. Evolution of Diverse Effective N₂-Fixing Microsymbionts of Cicer, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8090884/>
 39. Comparative Genome Analysis of Polar Mesorhizobium sp ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10820077/>
 40. Physiological and Biochemical Responses of Two Major Legume, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13259165/>
 41. Performance of liquid and carrier-based inoculants ... - ResearchGate, https://www.researchgate.net/publication/289321518_Performance_of_liquid_and_carrier-based_inoculants_of_Mesorhizobium_ciceri_and_PGPR_Pseudomonas_d_iminuta_in_chickpea_Cicer_arietinum_L_on_nodulation_yield_and_soil_properties
 42. Nitrogen fixation by common beans in crop mixtures is influenced by, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10184335/>
 43. Seed-applied micronutrient toxicity to rhizobia and impaired legume, <https://academic.oup.com/jambio/article/136/4/lxaf067/8085148>
 44. effect of inoculation methods of mesorhizobium ciceri and pgpr in, https://www.researchgate.net/publication/290919909_EFFECT_OF_INOCULATION_METHODS_OF_MESORHIZOBIUM_CICERI_AND_PGPR_IN_CHICKPEA_CICER_AR_IETINUM_L_ON_SYMBIOTIC_TRAITS_YIELDS_NUTRIENT_UPTAKE_AND_SOIL_PRO_PERTIES
 45. The impact of fungicides with contrasting toxicities to rhizobia on the, <https://connectsci.au/cp/article/76/9/CP24381/238048/The-impact-of-fungicides-with-contrasting>
 46. (PDF) Crop density affects weed suppression in organically, https://www.researchgate.net/publication/338530393_Crop_density_affects_wee_d_suppression_in_organically_managed_Sunflower
 47. Chickpea (Cicer arietinum L.) Biology and Biotechnology - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9654780/>
 48. Standardization of optimal seed rate for maximizing seed yield,

- <https://www.agronomyjournals.com/archives/2025/vol8issue12/PartK/8-12-81-395.pdf>
49. Optimizing seed rate of bold seeded varieties of chickpea (Cicer, <https://pub.isprd.in/index.php/jfl/article/download/22/18>)
 50. Chickpea Plant Population Trial - Saskatchewan Pulse Growers, <https://saskpulse.com/resources/chickpea-plant-population-2024-profit-results/>
 51. Assessing seed characteristics for improved winter survival of late, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13083018/>
 52. GUIDE TO WEED CONTROL 2016–2017 Publication 75, https://www.ovma.ca/wp-content/uploads/2022/02/pub75_2016.pdf
 53. Proceedings of the 73rd Annual Meeting of the North Central Weed, <https://ncwss.org/wp-content/uploads/NCWSS-Annual-Meeting-Proceedings-2018.pdf>
 54. Effectiveness of alternative herbicides on three Conyza species from, https://www.researchgate.net/publication/326736230_Effectiveness_of_alternative_herbicides_on_three_Conyza_species_from_Europe_with_and_without_glyphosate_resistance
 55. Revision of permitted substances for use in organic production (2025), <https://agrinfo.eu/book-of-reports/revision-of-permitted-substances-for-use-in-organic-production-2025/pdf/>
 56. Guide to weed control - Field Crop News, https://fieldcropnews.com/wp-content/uploads/2023/04/Pub_75A_2021_Event.pdf
 57. Agsc 102. Principles of Crop Production. Lecture Notes. 05-02-2025, <https://www.scribd.com/document/997598979/Agsc-102-Principles-of-Crop-Production-Lecture-Notes-05-02-2025>
 58. The relationship between natural rain intensity and Ascochyta blight, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9510387/>
 59. Ascochyta Blight in Chickpea: An Update - MDPI, <https://www.mdpi.com/2309-608X/9/2/203>
 60. The relationship between natural rain intensity and Ascochyta blight, <https://www.cabidigitallibrary.org/doi/pdf/10.31220/agriRxiv.2021.00063>
 61. A Mechanistic Weather-Driven Model for Ascochyta rabiei Infection, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8000037/>
 62. Quantifying Rainfall-Driven Splash Dispersal of Ascochyta rabiei in, https://www.researchgate.net/publication/399686038_Quantifying_Rainfall-Driven_Splash_Dispersal_of_Ascocyta_rabiei_in_Chickpea_Experimental_Evidence_for_Intensity_Thresholds_and_Epidemiological_Model_Improvement
 63. Agronomy Guide for Field Crops - Government of Ontario, <https://www.ontario.ca/files/2022-10/omafra-agronomy-guide-for-field-crops-en-2022-10-13.pdf>
 64. Dry Bean - Harvesting - Alberta Pulse Growers, <https://albertapulse.com/dry-bean-harvesting/>
 65. Package of Practices (Crops) 2016 - Kerala Agricultural University, <https://kau.in/sites/default/files/documents/pop2016.pdf>

66. From domestication syndrome to breeding objective - ConnectSci, <https://connectsci.au/cp/article/74/11/944%20/28413/From-domestication-syndro-me-to-breeding-objective>
67. Pod Shattering: A Homologous Series of Variation Underlying, <https://www.mdpi.com/2073-4395/8/8/137>
68. Advancing Grain Legumes Domestication and Evolution Studies, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9680859/>
69. Microscopical Analysis of Autofluorescence as a Complementary, <https://www.preprints.org/manuscript/202305.0960>
70. Paralogous Gene Recruitment in Multiple Families Constitutes, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11652722/>
71. Pod shattering in grain legumes: emerging genetic and environment, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8136915/>
72. Pod Dehiscence and Seed Dormancy in Pea (*Pisum* sp.) - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5404241/>
73. A DIRIGENT Gene GmDIR26 Regulates Pod Dehiscence in Soybean, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11074831/>
74. Glyphosate in Food 2026: What to Avoid + the Certified Free List, <https://plasticdetox.org/articles/glyphosate-detox-guide.html>
75. Crop desiccation - Wikipedia, https://en.wikipedia.org/wiki/Crop_desiccation
76. Dry Bean Desiccation and Harvest Considerations, <https://manitobapulse.ca/2019/03/dry-bean-desiccation-and-harvest-considerations/>
77. Dry Bean Desiccation and Harvest Guide, <https://saskpulse.com/resources/dry-bean-desiccation-and-harvest-guide/>
78. Prevalence of pesticide residues in foods of plant origin imported, https://www.researchgate.net/publication/399632918_Prevalence_of_pesticide_residues_in_foods_of_plant_origin_imported_from_Mercosur_countries_evidence_from_EU_border_control
79. Entering the European market for canned beans - CBI. EU, <https://www.cbi.eu/market-information/processed-fruit-vegetables-edible-nuts/canned-beans-0/market-entry>
80. Brief Introduction to Requirements for Processing of Organic Food, https://www.ceres-cert.de/docs/transfer/3-2-6_EN.pdf
81. Organic farming - Umweltbundesamt, <https://www.umweltbundesamt.de/en/topics/agriculture/toward-ecofriendly-farming/organic-farming>
82. (PDF) Seed Development and Maturation - ResearchGate, https://www.researchgate.net/publication/368919370_Seed_Development_and_Maturation
83. Seed Development and Maturation - WUR eDepot, <https://edepot.wur.nl/635777>
84. black sicilian garbanzo seeds | Nutrient-dense, drought-tolerant, https://www.allianceofnativeseedkeepers.com/collections/all-heirloom-seeds/products/black-sicilian-garbanzo?_pos=513&_fid=94e3a00ce&_ss=c
85. Thin layer drying characteristics and chemical properties of two local,

- <https://academic.oup.com/ijfst/article/61/1/vvag090/8672025>
86. Genetic variability for susceptibility to seed coat mechanical damage,
https://www.researchgate.net/publication/374638017_Genetic_variability_for_susceptibility_to_seed_coat_mechanical_damage_and_relationship_to_end-use_quality_in_kidney_beans
 87. i SAFFLOWER PRODUCTION - RUFORUM Institutional Repository,
http://repository.ruforum.org/sites/default/files/Safflower%20Manual_BUAN.compressed.pdf
 88. HANDBOOK FOR BETTER EDIBLE BEAN PRODUCTION - ADM,
<https://www.adm.com/globalassets/products--services/services/farmer-services/seedwest-seed-profile-catalog/adm-edible-bean-2018.pdf>
 89. Tips For Dry Beans Alberta 2021 | PDF | Crop Rotation - Scribd,
<https://www.scribd.com/document/1010116903/Tips-for-Dry-Beans-Alberta-2021>
 90. agronomy news - Colorado Department of Education,
<https://spl.cde.state.co.us/artemis/ucsuserials/ucsu20653internet/ucsu20653199208internet.pdf>
 91. Unraveling On-Farm Wheat Loss in Fars Province, Iran - MDPI,
<https://www.mdpi.com/2071-1050/15/16/12569>
 92. Chickpea seed storage in different packagings, environments and,
<https://www.scielo.br/j/rbeaa/a/kP8WfmV3QJjSwNBCqKsy4DS/?lang=en>
 93. Chickpea - Storage and Grading - Alberta Pulse Growers,
<https://albertapulse.com/chickpea-harvesting-storage/chickpea-storage-and-grading/>
 94. Flour & Cereal Quality Control Equipment | Lab Instruments – Bastak,
<https://www.bastak.com/blog>
 95. Scientific Opinion on the risk posed by pathogens in food of non,
<https://backend.orbit.dtu.dk/ws/files/52965039/jan13%20FoNAO.pdf>
 96. Long-Term Fungal Inhibition by Pisum sativum Flour Hydrolysate,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4524146/>
 97. PhD Thesis Juan Felipe Sandoval Rueda 1 Protein from UK,
https://research.aber.ac.uk/files/103084922/Rueda_Felipe_Sandoval.pdf
 98. mycotoxin levels and impacts on chemical and functional,
https://www.researchgate.net/profile/Chinaza-Awuchi/publication/348351464_Mycotoxin_Levels_and_Impacts_on_Chemical_and_Functional_Properties_of_some_Grains_Sold_in_Owerri_Imo_State_Nigeria/links/5ff97a8892851c13fefb42e7/Mycotoxin-Levels-and-Impacts-on-Chemical-and-Functional-Properties-of-some-Grains-Sold-in-Owerri-Imo-State-Nigeria.pdf
 99. Effects of storage temperature and relative humidity on cooking time,
https://www.researchgate.net/publication/367526743_Effects_of_storage_temperature_and_relative_humidity_on_cooking_time_and_moisture_uptake_of_selected_common_bean_Phaseolus_vulgaris_L_varieties_grown_in_Ethiopia
 100. Impact of Roasting on Functional Properties of Hard-to-Cook Beans,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11817523/>
 101. Induction of seed coat darkening in common beans (Phaseolus,
https://www.cropj.com/melo_14_1_2020_21_27.pdf

102. Consumer traits of common beans: a global and regional ... - Frontiers,
[https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1658338/f
ull](https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1658338/full)