

Agrotechnological Protocols for Oilseed Cultivation, Post-Harvest Processing, and Wedge-Press Expression in the Arenosols of Berlin-Brandenburg

Regional Agronomic Heritage and Historical Precedents

The production of edible plant lipids across the North German Glacial Basin has historically contended with coarse-textured outwash sands, erratic spring moisture, and compressed frost-free windows¹. In the Margraviate and later Province of Brandenburg, domestic oilseed husbandry and mechanical oil expression were essential pillars of both municipal illumination and agrarian autarky².

Albrecht Daniel Thaer (1752–1828), operating from his experimental estate and Royal Prussian Academic Institute of Agriculture at Möglin on the boundary between the Barnim plateau and the Oderbruch from 1804 onward, systematically categorized the agricultural capacities of these glacial formations⁴. In his four-volume work, *Grundsätze der rationellen Landwirthschaft* (1809–1812), Thaer established the *Bodenwertzahl* (soil rating system), evaluating arable land on a scale of 0 to 100 points (*Bodenpunkte*)⁴. Thaer demonstrated that the glacial outwash sands—yielding low ratings between 20 and 40 *Bodenpunkte*—could not sustain demanding oilseeds like winter colza (*Brassica napus*) without intensive organic manuring and structured crop rotations¹. Where fertile Oderbruch alluvial clays permitted high-yielding winter rapeseed, the lighter sands of the Barnim, Fläming, and Lusatia forced Thaer to champion hardier, fast-maturing alternatives: *Sommerrübsen* (summer turnip rape, *Brassica rapa* subsp. *oleifera*), *Öllein* (oil flax, *Linum usitatissimum*), and *Leindotter* (false flax, *Camelina sativa*)⁸. Thaer showed that attempting to force heavy-feeding oil crops onto dry sand resulted in crop failure, whereas matching crop physiology to localized ground moisture and cation-exchange capacity stabilized yields⁹.

In the alluvial lowlands of the Spreewald and along the Spree, Havel, and Oder river courses, localized microclimates and elevated water tables supported specialized lipid production. Wendish and German smallholders cultivated oil flax as a dual-purpose textile and culinary crop, establishing the regional standard of *Spreewälder Leinöl*¹¹. Paired with boiled potatoes and fresh lactic curd cheese (*Pellkartoffeln mit Quark und Leinöl*), this cold-pressed, unrefined oil provided complete essential fatty acids and protein to the rural labor force¹¹.

The mechanical expression of these seeds reached industrial maturation at the *Holländerwindmühle Straupitz* in Lower Lusatia¹¹. Originally constructed as an open-timber post mill (*Bockwindmühle*) around 1640 and reconstructed in 1850 by Master Millwright Gottfried Nitschke as a brick tower Dutch mill, the Straupitz facility was expanded in 1910 with a

complete commercial oil mill¹³. It remains the last operational triple-gearred windmill in Europe, housing grain milling, timber sawing, and oil pressing within a single kinetic powertrain¹³. The extraction technology at Straupitz preserves the historic Dutch wedge press (*Holländer Keilpresse*): vertical oak stamping poles, lifted via rotary cams driven by wind sails, fall onto acute hardwood driving wedges². These wedges transmit kinetic impact horizontally, compressing flaxseed meal wrapped in woolen filters and horsehair mats against heavy iron containment plates². Historical operational records indicate that an oil miller at Straupitz manipulated 3.5 to 4 metric tons of mechanical mass per shift, demonstrating that non-electric, kinetic timber machines can maintain high-throughput oil expression².

North of the Spree, Prussian agrarian edicts systematically reinforced oilseed cultivation. King Frederick William I and Frederick II (the Great) issued repeated decrees (*Königliche Hanf- und Flachsbaum-Edicten*) compelling agricultural estates to plant mandatory quotas of hemp (*Cannabis sativa*) and flax³. These crops provided essential naval cordage and canvas, while the residual seed was processed in municipal wedge-press oil mills (*Schlagmühlen*) for culinary lipids and lamp oil². Meanwhile, *Camelina sativa* was maintained across the sands of Brandenburg as an insurance crop⁸. Known as *Leindotter* due to its historical presence within flax stands, camelina was prized on 20-point drift sands where cruciferous rapes suffered complete floral abortion from early-summer moisture deficits⁸.

The mechanical lineage of the timber wedge press traces directly to the Dutch milling district of the Zaan, north of Amsterdam¹⁷. Following Cornelis Corneliszoon's 1592 patent for the industrial crankshaft, the Zaan region developed over 1,200 industrial windmills, including hundreds of specialized *oliemolens* operating wedge presses by the year 1700¹⁷. The design leveraged the mechanical advantage of double-inclined planes and structural hardwood resistance to generate cellular rupture pressures rivaling modern hydraulic expellers, establishing the technological foundation for regional oil production².

Pedological Constraints and Agro-Climatic Dynamics of Brandenburg Sands

The cultivation of oil-bearing crops in the Berlin-Brandenburg region operates within rigid geological and atmospheric boundaries. The pedological matrix is dominated by Pleistocene glacial drift, outwash plains (*Sander*), ground moraines, and glacial spillways (*Urstromtäler*) formed during the Saalian and Weichselian glaciations¹.

The dominant agricultural soils are classified as Arenosols (pleistocene sands and loamy sands), with localized Luvisols and Histosols in topographic depressions¹. These soils routinely exhibit yield indices between 25 and 40 *Bodenpunkte*¹. Arenosols feature a coarse sand fraction exceeding 75%, an active clay fraction below 8%, and low native soil organic matter (typically 1.0% to 1.8% humus in the arable epipedon)¹. This physical composition restricts the Available Water Capacity (AWC) to between 60 and 90 millimeters per meter of soil depth¹.

Consequently, these soils have virtually no buffer against dry spells; precipitation events drain rapidly through macropores into deep subsoil horizons¹. Furthermore, the potential

cation-exchange capacity is low ($\text{CEC}_{\text{pot}} < 10 \text{ cmol}_c/\text{kg}$), causing non-adsorbed nutrient anions and weakly held cations—principally nitrate (NO_3^-), sulfate (SO_4^{2-}), and potassium (K^+)—to leach into regional aquifers during winter drainage and early spring snowmelt¹.

Climatically, Berlin-Brandenburg falls within USDA Hardiness Zone 7b, situated in a transitional zone between oceanic maritime weather systems from Western Europe and continental anticyclones from the Eurasian landmass¹. This positioning creates three major agronomic constraints:

1. The average frost-free period spans approximately 136 to 137 days¹. The median date for the final spring air frost (0°C) occurs around May 8 (coinciding with the historical *Eisheiligen*), while the first killing autumn air frost arrives around September 22¹. Ground frosts (-1°C to -2°C) extend this hazard window, ending in late April and returning by early October¹. Thermal energy accumulation averages 2,600 to 2,800 Growing Degree Days (GDD, base 5°C) or 1,400 to 1,600 GDD (base 10°C)¹.
2. Mean annual precipitation ranges between 500 and 600 millimeters, with localized pockets in the rain shadow of the Fläming receiving less than 480 millimeters¹. The distribution is poorly matched to the water demands of spring-sown annuals: May and June, which coincide with critical stages of floral differentiation and cell division in oilseeds, frequently feature acute droughts driven by stable continental high-pressure blocks (*Ostlagen*)¹.
3. The macroporous aeration and warm surface temperatures of Arenosols accelerate the microbial mineralization of soil organic carbon (SOC)¹. Based on the VDLUFA (Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten) humus balancing framework, row crop cultivation and brassica production run heavy structural carbon deficits (potatoes at $-760 \text{ kg Humus-C/ha}$; intensive brassicas at $-1,000 \text{ kg Humus-C/ha}$)¹.

Without access to manufactured mineral fertilizers or grid-powered deep-well irrigation, soil productivity depends on closed-loop biological management. Nitrogen must be fixed symbiotically through leguminous leys, structural potassium must be solubilized from feldspars using deep-rooting catch crops, and the soil water matrix must be stabilized through the systematic incorporation of biochar, aged livestock manure, and continuous cover crops¹.

Agronomic Profiles and Field Husbandry of Target Crops

The following agrotechnological specifications govern the field cultivation, variety selection, rotational sequencing, and hand-tool husbandry for the seven designated oil-producing crops

in Brandenburg Arenosols (25 to 40 *Bodenpunkte*).

Crop Species	Optimal Open-Pollinated Cultivars	Sowing Window	Sowing Rate & Depth	Row Spacing & Field Density	Rotational Position & Interval	Primary Pathogens & Organic Countermeasures	Seed Saving Isolation Distance
Winter Rapeseed (<i>Brassica napus</i>)	'Lembkes Winterraps' (traditional), 'Bienvenue', 'Bristol' (00-type) ²⁰	Aug 20 – Aug 30	3.5 – ; 1.5 –	Rows 25 – ; 40 –	Follows early potatoes or fallow; precedes spring cereals; 5 – pause ¹	<i>Psylliodes chryscephala</i> , <i>Brassicogethes aeneus</i> , <i>Plasmiodiophora brassicae</i> . Liming, trap crops, strict pause ¹ .	1,000 – (entomophilous/anemophilous)
Spring Rapeseed / Turnip Rape (<i>Brassica rapa / napus</i>)	'Liho' (00-napus), historical 'Sommerrübsen' (<i>B. rapa</i> ssp. <i>oleifera</i>) ¹⁰	Mar 20 – Apr 10	5.0 – ; 1.0 –	Rows 15 – ; 80 –	Follows winter rye; precedes leguminous green manure; 5 –	<i>Phyllotreta</i> spp. (flea beetles), <i>B. aeneus</i> . Early planting, rolled firm seedbed	1,000 – (crosse s with wild brassicas)

					pause ²¹	d, wood-ash dusting.	
Oil Sunflower (<i>Helianthus annuus</i>)	'Peredovik' (high oil), 'Iregi Szürke Csíkos'	Apr 25 – May 10 (soil >)	5.0 – ; 3.0 –	Rows 50 – ; 6 –	Follows winter rye; precedes spring pulses; 4 – pause ²²	<i>Sclerotinia sclerotiorum</i> , <i>Botrytis cinerea</i> . Wide plant spacing , open canopy airflow, no fresh manure ²² .	1, 500 – (insect- vectored cross-pollination)
Linseed / Oil Flax (<i>Linum usitatissimum</i>)	'Lirina', 'Hella', regional Wendish landraces ²³	Apr 01 – Apr 20	60 – ; 1.0 –	Rows 10 – ; 600 –	Follows root crops or weed-free cereals; precedes rye; 6 – pause ⁹	<i>Fusarium oxysporum</i> f. sp. <i>lini</i> , <i>Aphthona euphorbiae</i> . Strict 7-year rotation , early sowing, clean seed.	10 – (autogamous; low outcrossing)
Industrial	'Finola' (early	Apr 20 – May	25 – ;	Rows 20 –	Follows clover-	<i>Botrytis cinerea</i> ,	2, 000 – (anemo

Hemp Seed (<i>Cannabis sativa</i>)	dwarf), 'Fedora 17', 'USO 31'	10	2.0 —	; 100 —	grass leys; precedes winter grains; 3 — pause	passerine bird predation. Rapid canopy shading suppresses weeds; acoustic hazing.	philous wind-pollinated)
Camelina / False Flax (<i>Camelina sativa</i>)	'Sonnenwende', 'Celine', regional ecotypes ⁸	Mar 15 – Apr 10	6.0 — ; 0.5 —	Rows 12 — ; 350 —	Pioneer crop on poor sand; flexible rotation ; 3 yr pause ⁸	Downy mildew (<i>Hyaloperonospora camelinae</i>). High natural glucanase resistance to pests ⁸ .	20 — (predominantly autogamous)
Persian Walnut (<i>Juglans regia</i>)	'Geisenheim 120', 'Geisenheim 26', 'Mars', 'Wunder von Monrepos' ²⁴	Plant Nov (dormant) or Mar	Bare-root graft; collar at ground level	Spacing 10 × 10 to 12 × 12 (70 —)	Field borders , farmstead shelterbelts; non-rotational perennial	<i>Gnomonia leptostyla</i> (anthracnose), spring radiation frost. Late-budding cultivars	1,000 — (wind-pollinated; graft scions)

						mandat ory ²⁴ .	
Europe an Hazeln ut (<i>Corylu s avellana</i>)	'Hallesc he Riesenn uss', 'Wunde r aus Bollweil er', 'Cosfor d' ²⁶	Plant Oct – Nov	Bare-ro ot shrub; collar at ground level	Spacing 4 × or 5 × (500 –)	Hedger ows, silvopas ture edges; non-rot ational perenni al	<i>Curculi o nucum</i> (hazeln ut weevil), catkin freezing . Interpla nt cross-p ollinator s; poultry scratchi ng ²⁶ .	500 – (anemo philous cross-p ollinator) ²⁷

Detailed Crop Husbandry and Biological Interventions

1. Winter Rapeseed (*Brassica napus*) and Summer Turnip Rape (*Brassica rapa*)

Winter rapeseed is the most nutrient-demanding annual oilseed cultivated in Northern Europe⁷. In Brandenburg Arenosols, winter survival depends on the crop achieving the "8-leaf stage," an 8 mm root collar diameter, and a taproot penetration depth greater than 15 cm before the first hard ground freezes in late October¹. This physiological threshold requires sowing strictly between August 20 and August 30. Earlier sowing risks excessive stem elongation (*Schossen*) before winter, leaving the apical meristem exposed to frost desiccation; later sowing prevents the taproot from reaching moisture beneath the frost line¹.

On marginal sands below 30 *Bodenpunkte*, historical records favor *Sommerrübsen* (*B. rapa* ssp. *oleifera*) over winter napus¹⁰. Turnip rape matures within 85 to 95 days, completing flowering before the onset of the July dry spells¹.

Cruciferous crops face several key phytosanitary pressures:

- **Pollen Beetle (*Brassicogethes aeneus*):** Emerging when spring temperatures exceed 12°C, the adult beetle bores into green bud clusters to access pollen, destroying floral organs. In organic systems, a perimeter trap crop of *Sommerrübsen* (which blooms 7 to 10 days earlier than winter rape) intercepts migrating beetles. Dusting stands with a 3 : 1 volumetric blend of screened wood ash and slaked lime at dawn, when plants are damp with dew, provides an alkaline physical deterrent.

- **Cabbage Stem Flea Beetle (*Psylliodes chrysocephala*) and Crucifer Flea Beetle (*Phyllotreta* spp.):** Flea beetles feed on emerging cotyledons during dry spring weather. Successful establishment requires a firm seedbed consolidated with wooden hand rollers or sheep treading, ensuring seeds make contact with subsoil capillary water for rapid germination.
- **Clubroot (*Plasmodiophora brassicae*):** This soil-borne plasmodiophorid forms persistent resting spores that can survive in acidic sand for over 15 years¹. Cruciferous crops must observe a strict 5- to 6-year cultivation interval and cannot follow cruciferous catch crops like *Sinapis alba*¹. Field soils should be amended with wood ash or slaked lime to maintain a pH above 7.2¹.

2. Oil Sunflower (*Helianthus annuus*)

The sunflower is well suited to the sandy soils of Brandenburg due to its deep taproot, which can penetrate 2.0 to 3.0 meters into glacial sands to recover leached potassium and subsoil water¹. The plant requires approximately 400 mm of cumulative water throughout the growing season, with peak uptake occurring during floral head expansion and anthesis in July²².

Sowing occurs between April 25 and May 10, once soil temperatures at a 5 cm depth stabilize above 8°C²². Sowing depth on sand must reach 4.0 to 5.0 cm to secure moist capillary contact and reduce seed predation by corvids²². The open-pollinated cultivar 'Peredovik' provides an optimal balance of oil yield (42% to 45%) and early maturation, enabling harvest before the arrival of autumn rains.

The primary fungal pathogen is White Mold (*Sclerotinia sclerotiorum*), which induces both basal stem rot and mid-stem/head collapse²². Because fungal sclerotia persist in sandy soils, stand densities must not exceed 6 to 8 plants per square meter, with row spacings widened to 60 or 70 cm to promote wind ventilation through the canopy²². Sunflowers must not share close rotational sequences with other susceptible broadleaf hosts, such as field peas or potatoes¹.

3. Linseed / Oil Flax (*Linum usitatissimum*)

Linseed has deep cultural and agronomic roots in Brandenburg, particularly in the Spreewald⁹.

Flax roots are shallow and fibrous, concentrating primarily within the upper 40 cm of the soil profile, which makes the crop vulnerable to early-season drought⁸. Sowing must be completed between April 1 and April 20 to capture residual winter soil moisture⁸. The seedbed requires fine consolidation; seeds placed deeper than 1.5 cm exhaust their hypocotyl reserves before emergence⁸.

The main agronomic challenge is "flax sickness" (*Leinmüdigkeit*), caused by the accumulation of *Fusarium oxysporum* f. sp. *lini*. Thaer recorded that attempting to crop flax at short intervals produced stunted, chlorotic stands⁹. A rotational interval of at least 6 to 7 years between flax crops is mandatory⁹. Because flax has minimal leaf area during early growth, weed competition can quickly overwhelm stands. Hand-tine weeding with wooden rakes must take place when

seedlings are 3 to 5 cm tall, followed by manual rogueing.

4. Industrial Hemp Seed (*Cannabis sativa*)

Hemp seed produces high-value dietary lipids, while the plant's biomass improves soil structure¹. On sandy Arenosols, hemp requires substantial base fertility, typically requiring 25 to 35 metric tons of composted farmyard manure per hectare to sustain rapid vegetative growth. Sown between April 20 and May 10, hemp quickly closes its canopy, suppressing problem weeds such as couch grass (*Elymus repens*) and fat hen (*Chenopodium album*).

For grain production, monoecious, short-stature cultivars like 'Finola' are preferred over tall fiber types. 'Finola' is daylight-insensitive, matures within 100 to 110 days, and maintains an

average height under 1.8 meters, which simplifies manual sickle harvesting. Seed shatter remains an ongoing risk: the plant flowers indeterminately, causing upper achenes to ripen and drop while lower parts of the floral spike are still immature. Harvest must be initiated when 70% to 75% of the seeds on the main stems have developed hard, marbled coats. Passerine birds, particularly finches and doves, can strip standing heads during the final two weeks of ripening, requiring active scaring or field monitoring.

5. Camelina / False Flax (*Camelina sativa*)

Camelina sativa is exceptionally reliable on poor sands⁸. Characterized by high water-use efficiency and a short cycle (80 to 95 days for spring types), camelina yields reliably on marginal sands with ratings as low as 20 to 25 *Bodenpunkte*⁸. Seedlings are cold-hardy down to -8°C , permitting sowing as early as mid-March⁸. This early establishment allows the crop to flower in May and fill seed ahead of summer heat stress¹.

Camelina synthesizes specialized glucosinolates and phytoalexins that make it unpalatable to crucifer flea beetles (*Phyllotreta* spp.) and pollen beetles⁸. Its root exudates also exert allelopathic suppression on annual weeds⁸. Because the thousand-grain weight is only 0.8 to 1.2 grams, the seed must be sown very shallowly (0.5 – 1.0 cm) into a rolled, firm seedbed⁸.

Camelina is also well-suited to intercropping (*Mischfruchtanbau*) with grain legumes like lentils (*Lens culinaris*) or field peas (*Pisum sativum*)³². The rigid camelina stems act as a physical trellis (*Stützfrucht*), preventing legume lodging and creating a resilient mixed canopy¹. After manual threshing, the small camelina seeds are readily separated from the larger legume seeds using round-wire hand sieves³³.

6. Persian Walnut (*Juglans regia*)

As a perennial tree crop, the Persian walnut provides an enduring lipid source that does not require annual field preparation. In Berlin-Brandenburg, the primary production risks are winter cold snaps (below -25°C) and late spring radiation frosts (*Spätfröste*) during mid-May, which destroy tender shoots and flowering catkins¹. Planting ungrafted seedlings (*Zufallssämlinge*) is unviable for reliable production due to erratic budbreak and prolonged juvenile latency (12 to 15 years).

Reliable production requires grafted, late-leaving cultivars:

- **'Geisenheim 120' and 'Geisenheim 26':** Selected for late spring budbreak (mid-to-late May), avoiding frost windows while consistently setting thin-shelled, easily cracked nuts²⁴.
- **'Mars':** A cold-hardy cultivar of Czech origin with late foliation, good anthracnose resistance, and early autumn nut maturity²⁴.
- **'Wunder von Monrepos':** Exceptional frost evasion, high crack-out ratio (>48%), and light-colored kernels²⁴.

On sandy Arenosols, walnut trees develop deep root architectures, but require supplemental hand watering during their first three establishment seasons³⁶. Trees should be spaced $10 \times 10 \text{ m}$ to $12 \times 12 \text{ m}$ along farmstead perimeters or within agroforestry windbreaks outside cold-air drainage pockets²⁴. Grafted stock begins bearing in years 4 to 5, reaching full yields (20 to 40 kg of in-shell nuts per tree) by year 12.

7. European Hazelnut (*Corylus avellana*)

The European hazelnut is a native deciduous shrub well-suited to glacial moraines, provided the topsoil maintains a minimum humus content²⁶. Hazelnuts flower in late winter (January to March)²⁷. While dormant catkins can survive -15°C , fully extended flowers suffer damage below -8°C . Cross-pollination is required to secure fruit set; compatible, frost-hardy varieties must be planted in close proximity:

- **'Hallesche Riesennuss':** A historic regional variety from Halle an der Saale; vigorous growth, thick-shelled large nuts, cold-hardy, and late-blooming²⁷.
- **'Wunder aus Bollweiler':** A late-blooming variety from Alsace; resistant to late winter frosts, serving as a reliable pollenizer for 'Hallesche Riesennuss' and 'Cosford'²⁶.
- **'Webbs Preisnuss':** A reliable English cultivar yielding elongated, high-flavor kernels with crack-out ratios above 45%³⁷.

Hazelnuts are managed as multi-stemmed stools containing 5 to 7 primary boles²⁶. To control the Hazelnut Weevil (*Curculio nucum*)—which oviposits inside young nuts in June, leaving hollow, damaged shells—the soil beneath the bushes should be scratched shallowly with hand hoes in late autumn to expose overwintering pupae to frosts and poultry foraging²⁶.

Post-Harvest Processing: Threshing, Cleaning, and Desiccation Thermodynamics

Harvesting, threshing, and cleaning oilseeds using hand tools requires precise execution. Premature cutting leaves immature, moisture-laden seeds with elevated chlorophyll and high free fatty acid levels, while delayed harvesting causes heavy losses from pod shattering, bird feeding, and mold colonization.

Manual Harvest Protocols and Shatter Control

Cruciferous pods become brittle as they mature. When seeds reach physiological maturity

(seed coats turn brown/black, seed moisture drops to ~15%), harvesting must occur exclusively in early morning (04:30 to 08:00). High nocturnal humidity and dew plasticize the outer pod valves, suppressing shatter losses⁴¹. Stems are severed using hand sickles or short scythes fitted with collection cradles, and bound into small sheaves (*Garben*). Sheaves are shocked upright in the field for 2 to 4 days or transferred to linen sheets to capture fallen seed. Sunflower heads are severed individually using hand saws or heavy curved knives once the reverse side of the floral disk turns lemon-yellow, the outer bracts wither to brown, and the achenes loosen under thumb pressure. The cut heads are transferred to airy lofts for curing. Linseed is pulled entirely by the roots (*Raufen*) rather than cut. Pulling keeps the long bast fibers intact for spinning and prevents sticky stem sap from gumming up the harvesting tools. Flax bundles are shocked in open-sided sheds until the seed capsules (*Bolls*) rattle clearly when shaken.

Camelina pods are naturally shatter-resistant⁸. Pods can remain standing in the field for over a week after turning golden-yellow without substantial pod drop, permitting harvesting during midday dry conditions⁸. Stems are cut with scythes and raked onto sheets.

Hemp stalks are cut with heavy sickles when the bracts surrounding the seed clusters display dark, fully developed achenes. Stalk bundles are inverted and suspended under open barn roofs over collection tarpaulins.

Walnuts and hazelnuts are harvested exclusively from ground falls²⁶. Fallen nuts must be gathered every 24 to 48 hours to prevent green outer husks (*Pericarp*) from rotting and staining the inner nutmeat, and to reduce colonization by soil fungi. The green husks are stripped by hand or tumbled in slatted wooden barrels with coarse gravel and water.

Manual Threshing and Aerodynamic Cleaning

Threshing is conducted on a level, swept timber or packed-clay threshing floor (*Tenne*) lined with tightly woven linen or hemp tarpaulins. Traditional jointed flails consist of a seasoned ash handle (*Hauhechel*) joined to a heavy hornbeam or oak beater (*Klöppel*) by a rawhide leather swivel. Sheaves are laid heads-inward in two parallel rows. Flailers deliver glancing, rhythmic strikes rather than heavy, perpendicular crushing blows; glancing strikes pop open siliques and capsules without crushing seed coats or releasing intracellular lipases. A two-worker team can

process 100 to 150 kg of dry brassica or flax sheaves per hour, recovering 25 to 45 kg of uncleaned seed⁴¹. Sunflower heads are threshed by rubbing them against slatted wooden washboards or coarse expanded-metal screens suspended over tubs.

Threshed seed mixtures contain chaff, pod fragments, dust, and weed seeds. Cleaning utilizes a two-stage manual separation process:

1. **Aerodynamic Winnowing:** The threshed mass is tossed into an ambient breeze ($3 - 5 \text{ m/s}$) using wide, shallow willow winnowing baskets (*Worfschaufeln*). Chaff, dust, and hollow husks are carried off by air resistance, while sound seeds fall back onto the capture mats.
2. **Tiered Sieve Grading:** The winnowed grain is passed through sets of circular wooden riddle sieves fitted with woven brass or iron wire screens. The first screen (scalper) retains large stem fragments and debris, allowing seeds to drop through. The lower

screen (sifter) retains the seed crop while passing fine sand, cracked particles, and micro-weed seeds.

Desiccation Thermodynamics and Safe Storage Moisture

Because seed oil is completely hydrophobic, all moisture within an oilseed is concentrated in the hydrophilic protein and carbohydrate fractions. As a result, an oilseed containing 40% oil and 10% total moisture actually holds that water at an effective concentration of roughly 16.6% within its active cellular tissue.

If seed moisture exceeds critical water activity thresholds ($a_w > 0.65$ to 0.70), degradation occurs along three biochemical pathways:

1. Hydrolysis of triglycerides by endogenous seed lipases, generating free fatty acids (FFA) that reduce culinary smoke points and ruin flavor stability.
2. Respiration of storage fungi (*Aspergillus glaucus*, *Penicillium* spp.), producing localized hot spots and generating toxic mycotoxins.
3. Uncontrolled lipid autoxidation triggered by metabolic heat, destroying polyunsaturated fatty acids.

To preserve quality, seeds must be dried to their safe equilibrium moisture limits within 48 hours of harvest:

- **Solar Drying:** Seeds are spread thinly ($1 - 2$ cm depth) across canvas stretchers elevated 30 cm off the ground, sheltered from direct sun by linen cloths, and stirred hourly with wooden rakes.
- **Hypocaust Kiln Drying:** During humid, overcast autumn weather, drying relies on indirect wood heat. A masonry hypocaust or slatted drying floor over a brick flue directs warm air ($35 -$) up through the seed beds. The seed mass temperature must **never exceed** 40°C ; higher temperatures denature functional proteins, degrade natural tocopherol antioxidants, and accelerate the oxidation of delicate omega-3 and omega-6 fatty acids.
- **Sealed Storage:** Dried seeds are transferred to glazed stoneware pots, charred oak barrels, or dry metal tins. Containers are sealed with beeswax-impregnated linen covers, elevated on wooden skids above dirt floors, and kept in a cool pantry ($4 - 12^\circ\text{C}$) with relative humidity below 65%.

Oilseed Crop	Harvest Moisture Window	Safe Storage Moisture Content	Scalper Screen Sieve Size	Sifter Screen Sieve Size	Maximum Drying Temperature	Storage Longevity (Safe Form)
Winter Rapeseed	14 —	7.0 —	3.0 mm	1.0 mm	40°C	2 —

d			round	slot		(whole, glazed pot) ²
Turnip Rape / Rübsen	13 —	7.0 —	2.8 mm round	1.0 mm slot	40° C	2 — (whole, glazed pot) ¹⁰
Oil Sunflower	16 —	7.5 —	8.0 mm round	3.0 mm round	40° C	1 — (unhulled, dry bins) ²
Linseed / Flax	12 —	8.0 —	3.5 mm round	1.2 mm mesh	40° C	2 years (whole, dark airtight) ²
Hemp Seed	16 —	8.0 —	4.5 mm round	1.8 mm mesh	35° C	1 year (whole seed, airtight) ²
Camelina	11 —	6.5 —	2.0 mm mesh	0.6 mm mesh	40° C	2 — (whole seed, dry pot) ³¹
Persian Walnut	30 — (fresh)	5.0 — (kernel)	Manual sorting	Manual sorting	35° C	1 — (in-shell, mesh sack) ²⁶
European Hazelnut	20 — (fresh)	6.0 — (kernel)	Manual sorting	Manual sorting	35° C	1 — (in-shell, slatted box) ²⁶

Mechanical Expression in the Wooden Wedge Press

Mechanical extraction via a traditional wooden wedge press combines timber mechanics, classical wedge statics, fluid dynamics, and cellular shear². By relying on the kinetic energy of falling stampers driving acute, self-locking wedges, this mechanism achieves cellular rupture pressures that exceed the capabilities of levers or direct weights².

Mechanical Pretreatment: Dehulling and Rolling

Within intact seeds, lipids reside inside sub-micron organelles known as oleosomes, which are surrounded by protective phospholipid monolayers and structural oleosin proteins¹⁷.

Compressive load alone cannot express oil from uncrushed seeds; uncrushed seeds simply consolidate into a rubbery, impenetrable mass².

1. **Dehulling (Decortication):** Mandatory for sunflowers, walnuts, and hazelnuts; optional for hemp¹⁷.
 - *Sunflowers:* Cracked through a manual mill with wide stone clearance, followed by winnowing to strip hulls. Retaining more than 12% to 15% hulls reduces oil recovery, as dry hull fragments reabsorb expressed oil within the press.
 - *Tree Nuts:* Shelled manually or with lever crackers; kernels are separated from internal woody septa (*Walnusscheide*)²⁶.
2. **Flaking:** Cleaned seed meat is rolled through a set of manually driven, smooth-surfaced stone or cast-iron rollers¹⁷. The gap is set to produce flakes between 0.25 and 0.35 mm thick¹⁷. This rolling shears the cell walls and exposes the internal lipid bodies, while preserving macroscopic drainage pathways for fluid escape under compression².

Conditioning Thermodynamics: Roasting and Viscosity

Flaked seed meal requires controlled thermodynamic conditioning before pressing². Heating accomplishes three vital structural tasks:

1. **Protein Denaturation:** Thermal energy denatures protective oleosin membranes, allowing fine oil droplets to coalesce into continuous liquid pools¹⁷. It also makes the parenchymal cell walls brittle, reducing the elasticity of the solid matrix and facilitating fracture under compressive strain².
2. **Moisture Flashing:** Internal trace water vaporizes, producing micro-pressures that crack surviving cell walls from within².
3. **Viscosity Reduction:** Fluid mechanics indicate that vegetable oil dynamic viscosity (η) falls exponentially as temperature increases, governed by the Andrade relationship:

$$\ln(\eta) = A + \frac{B}{T}$$

Where A is an empirical fluid constant and $B = E_a/R$ represents the activation energy of flow divided by the gas constant¹⁷. Warming meal from ambient temperature (

15°C) to 75°C reduces oil viscosity by more than 70%, facilitating Darcy flow through the microscopic pore networks of the pressed cake¹⁷. For culinary table oils requiring high antioxidant retention and light flavors, seed temperatures must stay strictly between 38°C and 42°C (cold-pressed parameters)². When pressing for maximum caloric yield, meal can be roasted to $85 - 105^{\circ}\text{C}$ ². Heavy cast-iron plates preheated in hearth embers to $70 - 80^{\circ}\text{C}$ are inserted into the press trough adjacent to the seed packets to prevent cold press timbers from chilling the meal and increasing oil viscosity².

Wedge Statics and the Self-Locking Boundary Condition

The horizontal compressive force (W) delivered by the falling stamper's vertical impact (P) is dictated by static equilibrium²:

$$P = W \left[\frac{2\mu_s \cos \alpha + (1 - \mu_s^2) \sin \alpha}{\cos \alpha - 2\mu_s \sin \alpha} \right]$$

Where α is the wedge half-angle and μ_s is the coefficient of static friction between the sliding wood faces².

A vital requirement of wedge design is the self-locking angle of friction ($\phi = \arctan \mu_s$)².

The wedge remains securely locked in position only if $\alpha < \phi$ ². While dry oak sliding on oak features $\mu_s \approx 0.62$ ($\phi \approx 31.8^{\circ}$), pressing seeds introduces expressed oil across all wooden surfaces². Vegetable oils act as boundary lubricants, reducing the friction coefficient of hardwood interfaces to $\mu_s = 0.35 - 0.39$ ². Under continuous oil lubrication:

$$\phi_{\text{lubricated}} = \arctan(0.35) \approx 19.2^{\circ}$$

If a driving wedge is shaped with a half-angle $\alpha \geq 19.2^{\circ}$, the elastic springback of the compacted seed cake will violently eject the wedge upward once the vertical stamper lifts². To ensure a dependable safety factor against operational shock and vibration, the driving wedge half-angle must be engineered strictly between $\alpha = 10^{\circ}$ and 12° (a total inclusive taper of 20° to 24°)².

Terzaghi Consolidation and Operational Dwell Cycles

Liquid expression from porous seed matrices follows Terzaghi's consolidation theory and Biot's poroelasticity principles¹⁷. Total applied mechanical stress (σ) is shared between effective stress (σ') carried by the solid seed cell skeleton and pore fluid pressure (u) borne by the liquid oil¹⁷:

$$\sigma = \sigma' + u$$

The dissipation of pore fluid pressure over time (t) and cake thickness (z) is governed by the one-dimensional consolidation equation:

$$\frac{\partial u}{\partial t} = C_v \frac{\partial^2 u}{\partial z^2}$$

Where C_v is the consolidation coefficient, directly proportional to cake permeability (k) and inversely proportional to oil viscosity (η)¹⁷.

As oil drains through the filtration layers, cake void ratio (e) collapses, causing permeability to drop substantially¹⁷. Striking the wedge continuously without pauses crushes the timber without releasing additional oil². The operator must follow a structured pressing sequence:

1. **Initial Seating:** 20 to 30 moderate blows drive the 10° wedge down, purging air and establishing fluid pressure².
2. **Primary Dwell Cycle:** The press is held under full static load for **3 to 5 minutes**, allowing viscous oil to flow through the porous cake².
3. **Secondary Drive:** As cake volume contracts, 15 to 20 firm blows re-establish maximum internal pressure ($15 - \quad$)².
4. **Final Dwell Cycle:** The press is held under load for **5 to 8 minutes** until oil drainage ceases².
5. **Releasing Blow:** The primary stamper is lifted, and a secondary stamper strikes the inverted releasing wedge, collapsing the horizontal assembly, releasing static strain, and freeing the compacted cake discs².

Filtration Membrane Construction

Conditioned meal is packed into double-layer filter assemblies². The inner envelope consists of unbleached linen or cotton canvas duck ($450 - 550 \text{ g/m}^2$)². The outer envelope is a heavy horsehair mat (*Presshaarmatte*), hand-woven from equine tail hair and bound with leather

straps². This horsehair exterior contains the extreme lateral bursting stresses generated in the trough, while its open pores allow clear oil to flow freely into the wooden collection pans below².

Crop Seed Type	Shelling Degree	Flake Thickness	Conditioning Moisture	Conditioning Temp	Expression Pressure	Dwell Duration	Cold-Press Extraction Efficiency
OO-Rapeseed	Intact seed	0.20 —	8.0 —	40 —	15 —	8 —	68 — of total oil ²
Turnip Rape / Rübsen	Intact seed	0.20 —	8.0 —	40 —	15 —	8 —	65 — of total oil
Oil Sunflower	> dehusked ¹⁷	0.25 — [cite: 17]	8.5 — [cite: 17]	38 — [cite: 17]	16 — [cite: 17]	10 — [cite: 17]	70 — of total oil ¹⁷
Linseed / Flax	Intact seed	0.15 —	8.5 —	35 —	18 —	10 —	65 — of total oil
Hemp Seed	Intact or light hull	0.25 —	8.0 —	35 —	15 —	8 —	62 — of total oil
Camelina	Intact seed	0.15 —	7.5 —	38 —	16 —	8 —	66 — of total oil
Persian Walnut	100% shelled	0.40 —	6.0 —	Ambient (	12 —	6 —	75 — of total

				25°C)			oil
Europe an Hazeln ut	100% shelled	0.40 —	6.0 —	30 —	14 —	6 —	74 — of total oil

Lipid Biochemistry, Food Safety, and Fine-Dining Culinary Utilization

Cold-expressed plant lipids consist of triacylglycerols along with minor fractions of free fatty acids, mono- and diacylglycerols, phosphatides, phytosterols, tocopherols, and volatile aromatic terpenes. In fine-dining cookery, these oils are treated as delicate finishing fats, stable emulsion bases, and preservation mediums, each with specific biochemical constraints.

Autoxidation Pathways and Clarification Racking

The susceptibility of an oil to autoxidation depends on its fatty acid unsaturation, measured by its Iodine Value (IV):

- Monounsaturated Oleic Acid (C₁₈ : 1 cis — 9) oxidizes relatively slowly.
- Polyunsaturated Linoleic Acid (C₁₈ : 2 cis — 9, 12) oxidizes roughly 10 to 40 times faster than oleic acid.
- Alpha-Linolenic Acid (ALA, C₁₈ : 3 cis — 9, 12, 15), containing three conjugated double bonds with labile bis-allylic methylene carbons, oxidizes 100 to 200 times faster than oleic acid.

Raw, unrefined oil from the wedge press contains suspended cellular fragments, water droplets, and mucilaginous gums¹⁷. If left unclarified, native seed enzymes and environmental yeasts trigger rapid lipolytic breakdown¹⁷.

- **Gravity Sedimentation (Racking):** Crude pressed oil is transferred to tall, narrow-necked glazed stoneware jars (*Setzkrüge*) and held in a cold cellar (8 — 12°C) for 10 to 14 days. Gravitational settling draws suspended seed particles, cellular debris, and heavy gums to the bottom.
- **Clarification Decanting:** The clear surface oil is siphoned off the sediment using curved copper or glass racking tubes into dry stoneware bottles. Bottles are filled to within 1 cm of the rim to minimize oxygen contact, corked, and sealed with molten beeswax.

Toxicological Hazards and Antinutritional Compounds

Maintaining food safety requires strict management of endogenous toxins:

1. **Erucic Acid** (C₂₂ : 1 cis — 13): Found in high concentrations (40% to 50% of total fatty acids) in older landrace *Brassica napus* and *Brassica rapa*⁴⁴. Dietary erucic acid

causes myocardial lipidosis (lipid droplet deposition in heart muscle tissue) and impaired myocardial contractility. European Union food safety standards mandate that erucic acid in culinary oils **must not exceed** 20.0 g/kg (2.0%) of total fatty acids. The community must isolate low-erucic "00" cultivars (such as 'Bienvenue' or 'Bristol') from legacy brassicas, maintaining a spatial separation of at least 1,000 meters . Traditional high-erucic rapeseed should be restricted to lamp fuel, timber preservative, or press lubrication.

2. **Glucosinolates:** Characteristic sulfur metabolites of the Brassicaceae⁴⁵. When seed cells are crushed with water present, endogenous myrosinase (β -thioglucoside glucohydrolase) enzymes break down glucosinolates into toxic isothiocyanates, nitriles, and goitrogens⁴⁵. Goitrogens block iodine uptake in the thyroid, depressing metabolic rates. Because intact glucosinolates are water-soluble polar compounds, they partition into the press cake rather than the pure oil⁴⁵. Careful racking ensures the culinary oil remains free of meal particles carrying these compounds⁴⁵.
3. **Cyanogenic Glycosides in Linseed:** Raw flaxseed contains significant levels (100 – 300 mg HCN equivalents/kg) of cyanogenic glycosides, predominantly linamarin, lotaustralin, linustatin, and neolinustatin⁴⁶. When seeds are crushed, endogenous linamarase enzymes hydrolyze these glycosides into acetone cyanohydrin, which breaks down into toxic hydrogen cyanide (HCN)⁴⁶. However, cyanogenic glycosides and linamarase enzymes are water-soluble macromolecules and are insoluble in non-polar triglycerides⁴⁶. Clean, properly racked linseed oil contains no cyanogenic glycosides or active HCN ; the toxic payload remains entirely within the solid press cake⁴⁶.
- *Press Cake Detoxification Protocol:* To render linseed press cake safe for human consumption (such as Spreewald *Mühlenbrot* flour) or livestock feed, the cake must be crushed and mixed with water (1 : 3 weight ratio) at 30°C for 12 to 18 hours¹¹. This hydration allows native linamarase to cleave the glycosides, releasing gaseous HCN ⁴⁶. The mixture is then subjected to a rolling boil (> 85 – 100°C) in an open iron pot for 30 to 45 minutes⁴⁶. Because hydrogen cyanide has a boiling point of 25.6°C , the toxin vaporizes safely into the outdoor air, while the heating denatures all remaining enzymes⁴⁶.

Oil Type	Dominant Fatty Acids	Smoke Point (Cold-Pressed	Oxidative Shelf-Life	Culinary Profile and
----------	----------------------	---------------------------	----------------------	----------------------

		)		Table Use
Winter Rapeseed (00)	62% Oleic (C18 :); 20% Linoleic (C18 :); 9% ALA (C18 :)	175 —	9 — (cool & dark)	Buttery, mild nutty flavor; universal cooking base, cold sauces, light sautéing.
Turnip Rape / Rübsen	58% Oleic; 22% Linoleic; 10% ALA	165 —	6 —	Fresh cabbage aroma; suited to root vegetable braises, pickles, and slaws.
Oil Sunflower	65% Linoleic; 22% Oleic; < ALA	160 —	6 —	Clean, light seed flavor; excellent base for egg-yolk mayonnaise and shallow frying.
Linseed / Flax	55% ALA (C18 :); 18% Oleic; 15% Linoleic ¹²	105 — (DO NOT HEAT)	3 — (keep refrigerated) ¹²	Deep nutty taste; cold dressing for hot <i>Pellkartoffeln</i> , quark, and warm rye bread ¹² .
Hemp Seed	56% Linoleic; 20% ALA; 3% GLA	110 — (DO NOT HEAT)	2 — (dark ceramic)	Vivid herbal green; grassy, nutty flavor; finishing oil for roast venison,

				poached pike, lentils.
Camelina	38% ALA; 18% Gondoic (C20 :); 17% Linoleic ⁵¹	145 —	12 — (high γ -tocopherols) ⁵²	Aromatic mustard-green s finish; stable table oil for autumn mushroom sautéing and dressings.
Persian Walnut	60% Linoleic; 18% Oleic; 12% ALA	150 —	3 —	Rich, roasted nutmeat finish; pairs with bitter chicory, poached pears, and goat curd.
European Hazelnut	78% Oleic (C18 :); 12% Linoleic; < ALA	160 —	10 — (low unsaturation)	Fine praline aroma; baking, sweet emulsions, delicate poultry dressings, trout finishing.

Quantitative Modeling: Subsistence Acreage and Community Scaling

Agronomic planning must accommodate two operational scales:

1. **The Household Unit (4 Adults):** Requiring 40 kg of pure edible vegetable oil per year¹⁸. Animal fats (lard from two pigs, rendered suet, and goat butterfat) furnish the remaining 90 to 100 kg of the household's annual dietary lipid requirement (130 — 140 kg total lipid pool, delivering roughly 30% of daily calories)¹⁸.
2. **The Agrarian Micro-Community (300 Individuals):** Requiring 3,000 kg of culinary vegetable oil annually (10 kg per capita), complemented by community livestock fats.

Household Cultivation Footprint (4 Persons)

To hedge against drought-induced crop failures on Arenosols, the household diversifies its oil cropping across three complementary plants: Camelina (drought/pest insurance), Sunflower (caloric density), and Linseed (traditional finishing oil)¹.

Performance metrics on 25–40 *Bodenpunkte* sands under manual cultivation:

- **Camelina:** Seed yield = 1,200 kg/ha ; seed oil content = 38% ; wedge press efficiency = 70%⁵¹. Oil expressed per kg seed = 0.266 kg . Pure oil yield = 319.2 kg/ha .
- **Sunflower:** Dehulled seed yield = 1,400 kg/ha ; oil content = 42% ; press efficiency = 72%¹⁷. Oil expressed per kg seed = 0.302 kg . Pure oil yield = 422.8 kg/ha .
- **Linseed:** Clean seed yield = 900 kg/ha ; oil content = 40% ; press efficiency = 68%³⁴. Oil expressed per kg seed = 0.272 kg . Pure oil yield = 244.8 kg/ha .

To harvest 40 kg of net oil, the household cultivates:

- **Camelina Plot:** 0.05 ha (500 m²) yielding 15.96 kg oil.
- **Sunflower Plot:** 0.04 ha (400 m²) yielding 16.91 kg oil.
- **Linseed Plot:** 0.04 ha (400 m²) yielding 9.79 kg oil.
- **Total Net Cropped Area:** 0.13 ha (1,300 m²).
- **Resilience Buffer Allocation:** Adding a 10% seed-saving retention, a 5% cleaning loss, and a 35% climate buffer against summer drought establishes a gross household footprint of 0.20 ha (2,000 m²)¹⁸. This corresponds precisely to the 0.2-hectare rotational block within the autarkic 1.5-hectare family farm system¹⁸.

Micro-Community Cultivation Footprint (300 Persons)

For 300 people requiring 3,000 kg of culinary oil, oilseed cultivation forms one block within a 6-year rotational cycle across 75 hectares of arable Arenosol land¹. One

12.50-hectare block is planted to oilseeds each year:

- 4.00 ha **Camelina:** Sown on the lightest, sandiest tracts (20–25 *Bodenpunkte*). Clean seed yield = 4,800 kg ; pure oil expressed = 1,276 kg⁸.
- 3.50 ha **Oil Sunflower:** Sown on deeper loamy sands following winter rye. Clean seed

yield = 4,900 kg ; pure oil expressed = 1,479 kg ²².

- 2.50 ha **Linseed**: Sown on low-lying moist fields. Clean seed yield = 2,250 kg ; pure oil expressed = 612 kg ³⁴.
- 2.50 ha **Winter 00-Rapeseed**: Sown after manured early potatoes¹. Clean seed yield = 3,750 kg ; pure oil expressed = 1,080 kg .
- **Total Community Seed Harvest**: 12.50 ha produces 4,447 kg of crude pressed oil.
- Deducting a combined 15% margin for seed stock (10%) and decanting sediment (5%) yields 3,780 kg **of clear edible table oil**, satisfying the 3,000 kg baseline while providing a 26% safety cushion against severe regional droughts¹⁸.

Operational Metric	Household Scale (4 Persons)	Micro-Community Scale (300 Persons)
Annual Table Oil Requirement	40.0 kg [cite: 18]	3,000.0 kg
Gross Seed Mass to Be Pressed	155.0 kg	11,500.0 kg
Net Cropped Land Area	0.13 ha (1,300 m ²)	10.00 ha
Gross Cropped Area (with 35% Climate Buffer + Seed Stock)	0.20 ha (2,000 m ²) ¹⁸	12.50 ha
Total Arable Estate Dedicated to Crop Rotation	1.00 ha (5-year rotation) ¹⁸	75.00 ha (6-year rotation)
Labor Hours: Soil Prep & Sowing (Hand / Oxen)	18.0 hours	450.0 hours
Labor Hours: Sickle/Scythe Harvesting	24.0 hours	625.0 hours

Labor Hours: Flail Threshing & Winnowing	32.0 hours [cite: 41, 42]	800.0 hours
Labor Hours: Dehulling, Flaking & Roasting	16.0 hours	400.0 hours
Labor Hours: Wedge Press Extraction	25.0 hours	575.0 hours
Total Annual Labor Hours	115.0 hours	2, 850.0 hours
Press Cake Byproduct Recovered	115.0 kg	8, 500.0 kg

Nutritional and Metabolic Feeding Value of Press Cake

The defatted press cake (*Ölkuchen*) represents 65% to 75% of the total seed weight processed in the wedge press⁴⁵. Because mechanical extraction leaves 8% to 14% residual oil within the cake, this byproduct provides both concentrated crude protein and dense metabolizable energy for livestock rations⁴⁵.

1. **Dairy Goats:** The primary milk producers in the farm system¹⁸. Feeding ²⁰⁰ to 350 g/day of cold-pressed rapeseed or camelina cake per doe increases the energy density of forage rations, raising milk yields by 12% to 18% and boosting fat-corrected milk solids for hard cheese production¹⁸.
2. **Swine:** Swine need complete amino acid profiles with adequate lysine, methionine, and threonine⁴⁴. Sunflower and hemp cakes can supply up to 15% of the daily ration for growing and finishing pigs⁵⁵. Rapeseed cake must be limited to 5% to 8% in finishing diets to prevent unsaturated lipids from softening the lard and to avoid liver stress from trace glucosinolates⁴⁵. Linseed cake must be boiled before feeding to volatilize hydrogen cyanide⁴⁶.
3. **Poultry (Laying Hens):** Hens efficiently convert oil cake into eggs, which provide essential zinc and selenium¹⁸. Camelina and boiled linseed cakes fed at 5% to 8% of the diet enrich egg yolks with anti-inflammatory omega-3 fatty acids⁴⁴. However, feeding raw rapeseed meal to brown-egg laying hens carrying the mutant FMO3 gene must be avoided; gut bacteria metabolize seed sinapine into trimethylamine (TMA), imparting a fishy odor to the eggs.

Press Cake	Crude Protein	Residual Oil	Crude Fiber	Ruminant ME	Swine ME	Max Safe	Feeding
------------	---------------	--------------	-------------	-------------	----------	----------	---------

Source	(% DM)	(% DM)	(% DM)	(MJ/kg DM)	(MJ/kg DM)	Inclusion Rate	Directives & Antinutritional Factors
00-Rapeseed Cake	32 — [cite: 45]	10 — [cite: 45]	11 — [cite: 45]	12.8	11.5	Goats: 20%; Pigs: 8%; Poultry: 5% [cite: 45]	Glucosinolates (<). Restrict in swine finishing rations ^{4 5} .
Sunflower Cake (Dehulled)	34 — [cite: 45]	11 — [cite: 45]	14 — [cite: 45]	12.2	10.8	Goats: 25%; Pigs: 15%; Poultry: 10%	Low lysine; high methionine. Excellent, safe feed without antinutritional risks.
Linseed Cake	31 — [cite: 45]	9 — [cite: 45]	8 — [cite: 45]	13.5	12.1	Goats: 10%; Pigs: 10%; Poultry: 8% [cite: 45, 56]	High cyanogenic glycosides (150 —). Must be boiled

							for 30 min before feeding ⁴⁶ .
Hemp Seed Cake	30 — [cite: 45]	9 — [cite: 45]	24 — [cite: 45]	10.5	8.8	Goats: 20%; Pigs: 8%; Poultry: 5% [cite: 45]	High woody fiber. Excellent goat ration; low digestibility for young weaner pigs ⁴⁵ .
Camelina Cake	33 — [cite: 45, 51]	11 — [cite: 45, 51]	10 — [cite: 45, 51]	13.2 [cite: 51]	11.8 [cite: 45]	Goats: 15%; Pigs: 5%; Poultry: 8% [cite: 44, 55]	Glucosinolates (25 —). Increases tissue n-3; restrict in swine to keep lard firm ⁴⁴ .
Walnut / Hazelnut Cake	22 —	12 —	6 —	14.5	13.5	Goats: 15%; Pigs: 15%; Poultry: 15%	High tannins and polyphenols. Palatable

							conditioning feed for breeding stock.
--	--	--	--	--	--	--	---------------------------------------

Operational Failure Modes, Diagnostic Detection, and Safety Hazards

Three Most Common Operational Failure Modes

1. Dynamic Loss of Wedge Self-Locking (Kinetic Wedge Ejection)

The static equilibrium of the wedge mechanism requires that the taper half-angle α remain strictly less than the friction angle ($\phi = \arctan \mu_s$)². As expression proceeds, expressed oil coats all contacting timber faces, driving μ_s down from 0.62 to roughly 0.35². If the wedge taper was planed too steep ($\alpha > 19^\circ$), or if repetitive stamper pounding mushrooms the top of the wedge, the normal reactive force from the compressed seed cake exceeds the resisting friction vector².

The operator detects this condition when the driving wedge bounces upward following stamper impacts rather than remaining locked in place, accompanied by high-pitched timber groaning.

The operator must immediately back out the wedge using the releasing stamper². The driving wedge faces must be re-planed to a true $10^\circ - 12^\circ$ angle². Powdered pine rosin (*Kolophonium*) or fine dry quartz sand must be dusted across the contacting faces to restore boundary friction before resuming expression.

2. Hydrolytic and Oxidative Rancidity (Free Fatty Acid Spikes)

If seed crops are harvested wet ($> 14\%$ moisture), stored in humid structures, or rolled and left unpressed for extended periods, native seed lipases hydrolyze triglycerides into free fatty acids¹⁷. Free linoleic and linolenic acids rapidly oxidize into unstable hydroperoxides (LOOH), which subsequently split into volatile aldehydes (hexanals, decadienals) that lower smoke points and cause off-flavors.

The oil develops a distinct acrid sensation at the back of the throat (*Kratzigkeit*) and an odor of damp cardboard or oil paint¹⁷. The Acid Value (AV) can be tested by shaking 5 mL of oil with 15 mL of warm, neutral 95% ethanol and titrating with 0.1 M potassium hydroxide (KOH) lye, using red cabbage anthocyanin as a visual indicator. An Acid Value exceeding 4.0 mg KOH/g ($> 2\%$ FFA) renders the oil unfit for the table, requiring diversion to soap

making or lamp oil.

This failure is prevented by immediate post-harvest drying to below 8% moisture². Crushed meal must be pressed immediately after flaking, and fresh oil should be cellared below 12°C to allow sediment to settle cleanly².

3. Cake Permeability Blindness (Filter Blinding and Capillary Pore Collapse)

Terzaghi consolidation indicates that oil expression depends on the void ratio (e) and permeability (k) of the collapsing seed mass¹⁷. If meal is pressed too wet ($> 11\%$ moisture) or driven with fast, continuous stamper strikes without dwell periods, the seed particles deform plastically². Fine meal particles migrate into the pores of the canvas bags, forming an impermeable, compacted crust across the filter face².

The driving wedge refuses to advance despite heavy stamper blows, yet oil discharge stops completely. The canvas envelopes bulge under internal pressure, and the wooden spacer blocks make a sharp cracking sound under impact without expressing oil².

The operator must stop pressing and disengage the driving wedge using the release mechanism². The blinded canvas filters must be stripped and washed. To restore drainage, the flaking roller gap should be widened slightly to 0.35 – 0.45 mm, and 5% to 8% coarse, dry hulls (e.g., sunflower hulls) should be mixed into the meal to serve as porous drainage channels during compression¹⁷.

Plainly Stated Safety Hazards

1. Spontaneous Combustion of Drying Oil Residues

High-iodine drying oils—specifically Linseed (IV 175–200), Camelina (IV 135–160), Walnut (IV 140–150), and Hemp (IV 140–160)—undergo rapid, exothermic oxidation when exposed to atmospheric oxygen⁵⁹. Polyunsaturated fatty acids react with air, releasing steady thermal energy⁵⁹.

When these oils coat porous materials with high surface areas—such as linen cleaning rags, canvas filter envelopes, dry press cake flour, or fine sawdust—the material insulates the interior of the pile⁵⁹. Heat generated by autoxidation cannot dissipate by convection, causing internal

temperatures to rise above the ignition threshold of dry cellulose ($> 250^{\circ}\text{C}$), producing open flames without an external ignition source⁵⁹.

All textiles, canvas envelopes, and rags contaminated with linseed, camelina, walnut, or hemp oil must **never** be balled up or discarded in timber bins. All oily fabrics must be submerged in an iron water kettle immediately after use, or hung flat on exterior masonry walls to cure safely in the air. The oil mill must be swept daily; fine organic flour mixed with atomized drying oils creates severe dust-explosion hazards in confined timber rooms.

2. Kinetic Projectile and Structural Timber Rupture

The wedge press converts up to 1,765 Joules of vertical energy per drop from a 150 kg stamper into hundreds of kilonewtons of lateral force². If parts become misaligned, or if

wedges are struck while friction-locked at incorrect angles, timbers can split along the grain (f_v) or crush perpendicular to the fibers ($f_{c,90}$)². A misaligned or overly lubricated wedge can shoot upward out of the mortise like a projectile, presenting a fatal impact hazard². Operators must never stand directly over the open press trough or reach across driving wedges while stampers are running². The main base log must be examined weekly for checking or grain splitting flanking the mortise cavity; the beam must be bound externally with shrink-fitted wrought-iron hoops to counter transverse tension stresses². All moving wedges, spacer blocks, and stampers must be shaped exclusively from seasoned European Hornbeam (*Carpinus betulus*) to provide high impact fracture toughness².

3. Hydrogen Cyanide (HCN) Gas Inhalation During Meal Detoxification

During the soaking, mashing, and boiling of linseed press cake, native linamarase enzymes break down cyanogenic glycosides, liberating free hydrogen cyanide (HCN)⁴⁶. HCN has a boiling point of just 25.6°C (78°F); when mixed with warm water or boiled, it escapes as an invisible, toxic gas with an odor of bitter almonds⁴⁶. Inhaled hydrogen cyanide blocks cellular respiration by binding to cytochrome c oxidase, leading to dizziness, collapse, and asphyxiation. Mashing and boiling of linseed press cake must **never** take place in enclosed rooms, basements, or unventilated spaces⁴⁶. Detoxification must be carried out outdoors or under a high-draft masonry chimney⁴⁶. Workers boiling the mash must stand upwind of steam vents to avoid inhaling vapors during the first 20 minutes of heating⁴⁶.

Agronomic Integration and Culinary Autarky

On the sandy Arenosols of Berlin-Brandenburg, lipid autarky is achievable through an interconnected agroecological system rather than isolated single-crop farming. Cruciferous rapes and turnip rapes supply the baseline volumes of high-smoke-point, oleic-rich oils needed for sautéing and hot cookery, provided 6-year rotational gaps and spatial isolation protect crops against clubroot and erucic acid cross-pollination¹. Sunflowers, with deep taproots that mine water and potassium from deep sand horizons, furnish clean culinary oils for hot cooking and stable emulsions while returning valuable organic matter to the soil¹. Linseed sustains regional culinary heritage, providing the cold finishing oil that elevates boiled potatoes and curd cheese to civilized gastronomic status, while requiring minimal processing heat to protect its sensitive omega-3 fatty acids¹¹.

Camelina sativa serves as the agrotechnological cornerstone, yielding stable, omega-3-rich table oil on marginal 20-point sands that resist flea beetles and spring droughts without requiring agrochemicals⁸.

As these annual cropping systems mature, permanent agroforestry borders of late-leafing walnuts and cold-hardy hazelnuts expand along field perimeters, providing concentrated tree lipids that require no annual field preparation while buffering open cropland against wind erosion¹.

Expressed through the applied physics of a hornbeam wedge press, these crops supply clear

oils for the civilized table, while their residual high-protein press cakes nourish the dairy goats, swine, and poultry that complete the farm's closed nutrient cycle². By linking historic North German husbandry with mechanical physics and lipid biochemistry, the self-sufficient household and expanding community achieve complete independence in food oils, securing both caloric survival and an enduring culinary culture.

Works cited

1. 24 Berlin Crop Rotation Schedule.pdf
2. 15 - Wooden Wedge Oil Press Design.pdf
3. herren kein Interesse daran gehabt haben, dass Landarme bei,
<https://d-nb.info/1224688309/34>
4. Über Albrecht D. Thaer - Oderbruchpavillion,
<https://archiv.oderbruchmuseum.de/thaer/>
5. Möglinsche Annalen der Landwirthschaft - Google Books,
https://books.google.com/books/about/M%C3%B6glinsche_Annalen_der_Landwirthschaft.html?id=E806AAAaAAJ
6. ADEL VERBINDET - Brill, <https://brill.com/downloadpdf/edcollbook/title/48675.pdf>
7. Ein Vergleich der weltweit wichtigsten Anbauregionen für Ölsaaten,
https://literatur.thuenen.de/digbib_extern/bitv/zi022724.pdf
8. Leindotter – Comeback einer unterschätzten Ölpflanze - Flur & Furche,
<https://flurundfurche.de/leindotter-comeback-einer-unterschaetzten-oelpflanze/>
9. Garten und Ackerbau. Thaer - Möglin. Mark Brandenburg - textlog.de,
<https://www.textlog.de/41242.html>
10. Der Landwirt 1877 Jg. 13,
https://bibliotekacyfrowa.pl/Content/119800/PDF/GSL_P_31377_IV_1877_57250_0_46.pdf
11. ÖLMÜHLE STRAUPITZ - Pension Willischza,
<https://www.pension-willischza.de/oelmuehle-straupitz>
12. Im Spreewald gibt es gesundes Leinöl aus der Straupitzer Mühle,
<http://leeves.de/2016/12/18/leinoel-aus-der-hollaender-windmuehle-straupitz-im-spreewald/>
13. Die Geschichte - Windmuehle Straupitz,
<https://www.windmuehle-straupitz.de/geschichte>
14. Holländerwindmühle Straupitz - Spreewald,
https://willkommen.spreewald.de/de/spreewald/streaming/detail/POI/DAMASSpreewaldPoi_11597/hollaenderwindmuehle-straupitz
15. Holländerwindmühle Straupitz - Kräutermühlenhof Burg (Spreewald),
<https://www.kraeutermuehlenhof.de/orte/hollaenderwindmuehle-straupitz/>
16. Die Ölmühle - Windmuehle Straupitz,
<https://www.windmuehle-straupitz.de/geschichte/oelmuehle>
17. 22 Wedge Press Oil Extraction.pdf
18. 117 Minimum Acreage For Family Sustenance.pdf
19. Quantifizierung aktueller und zukünftiger Nährstoffeinträge und,
https://literatur.thuenen.de/digbib_extern/dn066338.pdf

20. Biographisches Handbuch zur Geschichte des Pflanzenbaus,
<https://dokumen.pub/biographisches-handbuch-zur-geschichte-des-pflanzenbaus-reprint-2013nbsped-9783110967104-9783598113246.html>
21. LfL-Information - Bayerische Landesanstalt für Landwirtschaft - LfL,
https://www.lfl.bayern.de/mam/cms07/publikationen/daten/informationen/endbericht-forschungsplan-oekologischer-landbau-2013-2017_lfl-information.pdf
22. Anbauratgeber Sonnenblumen - LG Seeds,
https://www.lgseeds.de/fileadmin/user_upload/_Mediathek/UFOP_Praxisinfo_Sonnenblumen_2024.pdf
23. Öko-Saatgut 2025 - Marktgesellschaft der Naturland Bauern AG,
https://marktgesellschaft.de/wp-content/uploads/2025/05/MGN_Saatgutkatalog_2025_final_Web.pdf
24. Welche Walnuss, wenn die Eisheiligen immer Minusgrade bringen?,
<https://forum.garten-pur.de/viewtopic.php?t=67924>
25. Rund um die Walnuss - BUND Bodensee-Oberschwaben,
https://www.bund-bodensee-oberschwaben.net/fileadmin/bodenseeoberschwaben/Rund_um_die_Walnuss.pdf
26. Haselnuss 'Wunder aus Bollweiler' kaufen - Garten Schlüter,
<https://www.garten-schluter.de/haselnuss-wunder-aus-bollweiler/68653>
27. Haselnuss 'Hallesche Riesennuss' - Baumschule Horstmann,
<https://www.baumschule-horstmann.de/shop/exec/product/64/2514/haselnuss-hallesche-riesennuss.html>
28. Haselnuss 'Wunder aus Bollweiler' - Fruchtwelt,
<https://fruchtwelt.de/products/haselnuss-wunder-aus-bollweiler>
29. Albrecht Daniel Thaer - Zeno.org,
<http://www.zeno.org/Literatur/M/Fontane,+Theodor/Reisebilder/Wanderungen+durch+die+Mark+Brandenburg/Das+Oderland/Das+Oderbruch+und+seine+Umgebungen/M%C3%B6glin/Albrecht+Daniel+Thaer>
30. Aussaatempfehlungen Herbst 2021 + Frühjahr 2022,
<https://altmarksaaten.de/wp-content/uploads/2021/07/Aussaatempfehlungen-2021-2022.pdf>
31. Bliesgau Ölmühle Leindotter Anbauinformationen | Pflanzenöl kaufen,
<https://bliesgauoele.de/produkte/leindotter-1/leindotter-anbauinformationen/>
32. Leindotter-Erbсен-Mischfruchtanbau Aufbau einer nachhaltigen,
<https://mil.brandenburg.de/sixcms/media.php/9/Forum%20-%20Leindotter%20als%20nachhaltige%20Lasur.pdf>
33. Öko Linsen Anbau in Deutschland,
https://marktgesellschaft.de/wp-content/uploads/2026/03/MG_Anbautelegramm_10_Linsen_2024_Druck.pdf
34. 2. Ertragsstruktur des Mischfruchtanbaus von Lein (Linum usitat,
https://literatur.thuenen.de/digbib_extern/bitv/dk041193.pdf
35. NABU-Streuobst- Rundbrief 2/2024,
<https://www.nabu.de/imperia/md/content/nabude/streuobst/rundbriefe/so-2-2024-10-6-2024-sm-bk-farbig.pdf>
36. Obstbäume kaufen Potsdam — alte Sorten & Hochstamm | KANIA,

- <https://www.kania-galabau.de/obstbaeume-kaufen-potsdam>
37. Wildobst | Vielfältig, wild und lecker - Baumschule Bradfisch,
<https://bradfisch.de/pflanzenverwendung/wildobst.html>
 38. Ländliche Entwicklung in Bayern - Gemeinde Mühlhausen,
<https://www.muehlhausen-sulz.de/assets/Uploads/shop/Gehoelzpfad-Muehlhausen-Auflage-2-2012.compressed.pdf>
 39. Der Anbau von Haselnuss Sorten in Deutschland,
<https://weinterrasse.ch/gerichte/haselnuss-sobe-sub/>
 40. Haselnuss 'Rotblättrige Zellernuss' kaufen - Garten Schlüter,
<https://www.garten-schluter.de/haselnuss-rotblaettrige-zellernuss/63405>
 41. SURVEY AND ANALYSIS OF POST-HARVEST LOSSES,
<https://openknowledge.fao.org/server/api/core/bitstreams/c14e56ad-40d0-4909-87b3-1d1001414e1f/content/AC301e04.htm>
 42. Post Harvest Technology of Rice | PDF - Scribd,
<https://www.scribd.com/document/312087369/Rice-Postharvest>
 43. Theory and application of crop threshing technologies,
<https://archive.org/download/theoryapplicatio0000devn/theoryapplicatio0000devn.pdf>
 44. Camelina (Camelina sativa (L.) Crantz) as Feedstuffs in Meat Type,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8833380/>
 45. Cold-Pressed Oilseed Cakes as Alternative and Sustainable Feed,
https://air.unimi.it/retrieve/8574fdd7-67bf-4171-9d89-8fc93b7ba939/foods_doi.org_10.3390_foods12030432.pdf
 46. Bioprocessing of Functional Ingredients from Flaxseed - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6222892/>
 47. Metabolism of the Cyanogenic Glucosides in Developing Flax - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7407305/>
 48. Evaluation of the health risks related to the presence of cyanogenic,
<https://www.efsa.europa.eu/sites/default/files/consultation/consultation/181127-ag.pdf>
 49. Cyanogenic Glucosides in Plant Foods | PDF | Cyanide - Scribd,
<https://www.scribd.com/document/856354146/cyanogenic-plant>
 50. Development of a method to remove cyanogen glycosides from,
https://www.researchgate.net/publication/227803805_Development_of_a_method_to_remove_cyanogen_glycosides_from_flaxseed_meal
 51. Camelina (Camelina sativa (L.) Crantz) - Encyclopedia.pub,
<https://encyclopedia.pub/entry/19602>
 52. The influence of compound feed enrichment with rapeseed and,
https://www.researchgate.net/publication/282454159_The_influence_of_compound_feed_enrichment_with_rapeseed_and_linseed_cake_on_the_meat_characteristics_and_fatty_acids_composition_of_beef_bulls
 53. Digestibility of amino acids in organically cultivated white-flowering,
https://www.researchgate.net/publication/50937941_Digestibility_of_amino_acids_in_organically_cultivated_white-flowering_faba_bean_and_cake_from_cold-pressed_rapeseed_linseed_and_hemp_seed_in_growing_pigs

54. Schlussbericht zum Thema "Modellhaftes Demonstrationsnetzwerk,"
https://lupinenverein.de/wp-content/uploads/2023/11/Lupinen-Netzwerk-Schlussbericht-gesamt_web.pdf
55. Chemical Composition and Functional Effects of Oilseed Cakes on,
<https://www.mdpi.com/2076-2615/16/14/2266>
56. (PDF) Linseed Cake Supplementation Increases Tissue n-3 PUFA,
https://www.researchgate.net/publication/400080266_Linseed_Cake_Supplementation_Increases_Tissue_n-3_PUFA_Levels_but_Reduces_Growth_Performance_in_Broiler_Chickens_at_Higher_Inclusion_Levels
57. Understanding rancidity of nutritional lipids,
<https://www.supplysidesj.com/supplement-regulations/understanding-rancidity-of-nutritional-lipids>
58. Drying oil, <https://en-academic.com/dic.nsf/enwiki/392892>
59. ASTM D6801 Spontaneous Heating Temperature Testing | Fire Safety,
<https://infinitalab.com/blog/astm-d6801-spontaneous-heating-temperature/>
60. Primer on Spontaneous Heating and Pyrophoricity,
<https://www.energy.gov/documents/doe-hdbk-1081-2014>