

Systemic Integration of *Cichorium intybus* and *Taraxacum officinale* as Locally Cultivated Coffee Surrogates in the Berlin-Brandenburg Region: Agronomic, Phytochemical, and Culinary Dynamics

The increasing global demand for caffeine-free, locally sourced beverage alternatives has catalyzed renewed commercial and gastronomic interest in the taproots of *Cichorium intybus* var. *sativum* (industrial chicory) and *Taraxacum officinale* (common dandelion)¹. Both species, belonging to the Asteraceae family, produce massive taproots rich in the complex carbohydrate inulin and a highly specific matrix of phenolic acids and bitter sesquiterpene lactones⁴. When subjected to controlled thermal degradation through roasting, these roots undergo profound physicochemical transformations, developing a volatile metabolomic profile that closely mimics the organoleptic properties of roasted *Coffea arabica*⁷.

The geographic transposition of this production system to the Berlin-Brandenburg region of Germany presents a unique set of pedoclimatic challenges and advanced agroecological opportunities¹⁰. To establish a viable, sustainable local supply chain for high-end culinary applications—specifically caffeine-free espresso surrogates and structural flavor components for complex desserts such as Tiramisu—every phase of production must be rigorously engineered¹². This requires a systemic understanding of crop rotational architecture, heavy metal phytoindication, post-harvest enzymatic kinetics, roasting thermodynamics, solid-liquid extraction rheology, and human gastrointestinal physiology.

Pedoclimatic Constraints and Agronomic System Design in Berlin-Brandenburg

Climatic Boundaries and Operational Frost Windows

The architectural design of an agronomic system for root cultivation in the Berlin-Brandenburg region must be inextricably linked to local climatic boundaries. Situated within USDA Hardiness Zone 7b, the region is characterized by significant seasonal temperature variations and an operational growing season strictly dictated by absolute frost dates¹⁰. Historical meteorological data indicates that the last spring air frost (0°C/32°F) typically occurs around May 8, while the first autumn air frost arrives near September 22, delineating a reliable frost-free window of approximately 136 to 137 days¹⁰.

However, chicory and dandelion exhibit robust cold tolerance, establishing themselves as

highly resilient cool-season crops. They thrive in moderate temperatures ranging from 10°C to 24°C, with an optimal photosynthetic range between 15°C and 20°C¹⁵. By observing light ground frost thresholds (-1°C), the operational growing season in Berlin expands to up to 163 days¹⁰. Furthermore, by deploying protective horticultural fleece or utilizing cold-hardy cultivars, this functional window can be pushed to roughly 204 days, allowing ample time for the massive accumulation of root biomass¹⁰. Extreme summer heat waves, which are increasingly prevalent in the Brandenburg region, present a significant physiological stressor. Temperatures exceeding 25°C (77°F) can induce premature bolting (floral initiation), decrease overall root quality, and rapidly elevate the concentration of bitter sesquiterpene lactones in the tissues¹⁰.

Soil Pedology, Hydrology, and Nutrient Management

The defining limitation of the Brandenburg agricultural landscape is its pedology. The region is dominated by Pleistocene-derived glacial sands and sandy loams (Arenosols), which are frequently characterized by critically low agricultural yield indices (rated between 25 and 40 Bodenpunkte)¹⁰. These coarse-textured soils possess an abundance of macropores, ensuring excellent aeration and unhindered taproot elongation, which is essential for chicory roots that can penetrate 30 to 50 cm into the soil profile¹⁰. However, this porosity simultaneously results in an exceptionally low Available Water Capacity (AWC) and a high susceptibility to the downward leaching of water-soluble macronutrients, particularly nitrate and potassium¹⁰. To cultivate deep-rooting Asteraceae species successfully in this fragile environment, the soil structure and organic carbon pools must be systematically optimized. The inherently low humus content of sandy Arenosols necessitates a rigorously structured crop rotation. The integration of nitrogen-fixing leguminous species, such as the deep-rooting Blue Lupin (*Lupinus angustifolius*), into a multi-year sequence preceding chicory cultivation guarantees a closed-loop supply of macronutrients¹⁰. Blue Lupin, through symbiotic infection by *Bradyrhizobium* species, biologically fixes massive volumes of atmospheric nitrogen (up to 180 kg N/ha) and develops taproots capable of penetrating up to 3.0 meters to fracture deep compaction layers¹⁰.

Soil/Nutrient Parameter	Pedological Impact on Asteraceae Taproots	Mitigation Strategy in Brandenburg Arenosols
Texture and Porosity	Coarse sands (Arenosols) permit unhindered vertical taproot elongation but cause acute summer drought stress ¹⁰ .	Maintenance of consistent irrigation; incorporation of engineered biochar to exponentially increase AWC ¹⁰ .
Nitrogen (N) Availability	Excessive synthetic N leads to rampant foliar growth at	Utilize residual biological N from preceding leguminous

	the expense of root expansion, and elevates tissue bitterness ¹¹ .	cover crops (e.g., Hairy Vetch, Blue Lupin); limit active N top-dressing ¹⁰ .
Potassium (K) Availability	Essential for carbohydrate translocation, water regulation, and massive taproot expansion ¹⁰ .	Soil inoculation with Potassium Solubilizing Bacteria (KSB) such as <i>Bacillus mucilaginosus</i> to weather highly insoluble soil silicates ¹⁰ .

Rotational Architecture and Phytosanitary Gaps

Chicory and dandelion cultivation must be integrated into the broader 5-year crop rotation characteristic of advanced organic systems in Berlin-Brandenburg¹⁰. A highly optimized sequence typically utilizes a heavy feeder phase (e.g., late winter cabbage or broccoli) followed by moderate feeders and soil-restoring species¹⁰. The primary pathological threats in this region dictate strict rotational intervals. *Plasmodiophora brassicae*, an obligate biotrophic pathogen causing clubroot in Brassicaceae, requires a strict 4-to-6-year cultivation pause between susceptible crops¹⁰. Similarly, *Rhizoctonia solani* poses severe risks to Solanaceae crops like potatoes¹⁰.

Because *Cichorium intybus* and *Taraxacum officinale* belong to the Asteraceae family, they act as highly effective non-host rotational breaks. They do not harbor *Plasmodiophora brassicae* or *Rhizoctonia solani*, thereby starving these soil-borne pathogens of their primary hosts¹⁰. When slotted into the third or fifth year of a rotation—following the termination of cover crops like "Landsberger Gemenge" or winter rye via roller crimping and ultra-shallow tillage—chicory and dandelion benefit from the highly friable, aerated, and pathogen-suppressed seedbed without requiring deep moldboard plowing that would rapidly oxidize organic matter¹⁰.

Cultivation Mechanics and Stand Establishment

Cichorium intybus var. *sativum* (Root Chicory)

Optimal root yield and morphology are highly sensitive to planting density and spatial geometry. For industrial root chicory, seeds are sown directly into the ground approximately 1.0 to 1.5 cm deep in early spring (April to May), timed precisely 3 to 4 weeks before the final frost-free date¹⁵. To maximize individual root fresh weight while maintaining a dense canopy to suppress competitive weed emergence, an intra-row spacing of 12 to 15 cm (4.7 to 6.0 inches) and an inter-row spacing of 30 to 50 cm is strongly recommended¹⁵.

Agronomic studies on plant density reveal that excessive crowding (e.g., extremely narrow intra-row distances) restricts individual root diameter, leading to weak, spindly taproots that lack the mass necessary for efficient roasting²². Conversely, excessive spacing promotes multiple head formations per root, which diminishes the concentration of deeply stored carbohydrates³⁰. The cultivar 'Pagana', for instance, exhibits optimal root fresh weight per plant

when cultivated at 15 to 20 cm spacings³⁰. During the vegetative phase, chicory requires consistent, moderate moisture, particularly during germination and initial taproot establishment, receiving 1 to 2 inches of water per week¹⁴. Drought stress during this critical window triggers premature bolting and forces the plant to synthesize higher concentrations of defensive, bitter secondary metabolites¹⁵.

Taraxacum officinale (Common Dandelion)

Dandelion cultivation mirrors the spatial and thermal parameters of chicory but requires specific adjustments for seed morphology. The seeds are exceptionally small and require photic stimulation (light) to break dormancy and sprout¹¹. Consequently, they must be sown ultra-shallowly, approximately 0.25 inches (0.6 cm) deep, and barely covered with fine soil¹¹. Germination kinetics are highly temperature-dependent, optimizing between 10°C and 21°C (50°F to 70°F); soil temperatures exceeding 25°C (77°F) induce thermal dormancy and stall germination, rendering midsummer sowings highly ineffective in Brandenburg¹¹.

Once true leaves establish, seedlings are thinned to stand 15 to 20 cm (6 to 8 inches) apart within rows spaced 30 to 45 cm apart¹¹. Dandelion is exceptionally resilient and low-maintenance, though maintaining steady moisture is critical to prevent the leaves and roots from developing overwhelming bitterness¹¹. In non-irrigated, pure crop setups, *Taraxacum officinale* yields approximately 1.5 to 2.4 t/ha of dry root mass³¹. However, under optimized agricultural conditions with targeted nitrogen application (e.g., 20 kg N/ha), yields can scale dramatically, approaching 10.5 t/ha³². Furthermore, advanced crossbreeding programs utilizing *Taraxacum kok-saghyz* (rubber dandelion) genetics are currently being explored to radically increase root biomass and diameter, potentially revolutionizing the commercial viability of dandelion as a primary cash crop³⁴.

Urban Soil Toxicity and Heavy Metal Phytoindication

A critical, often overlooked limiting factor for cultivating wild or semi-cultivated Asteraceae species in the Berlin metropolitan and peri-urban area is the inherent risk of heavy metal bioaccumulation. Both *Cichorium intybus* and *Taraxacum officinale* are exceptionally efficient phytoremediators and have been utilized globally as phytoindicators for severe soil pollution³⁶. Dandelion and chicory roots function as aggressive perennial storage organs, efficiently absorbing and concentrating elements from the soil solution via symplastic and apoplastic mechanisms³⁸. When grown in contaminated urban soils—which are frequently compromised by historical industrial runoff, tetraethyl lead residues from vehicular emissions, and atmospheric deposition—these plants accumulate exceptionally high levels of Cadmium (Cd), Lead (Pb), Zinc (Zn), and Copper (Cu)³⁸.

In controlled solution culture trials, chicory accumulated up to 890 µg Cd/g in its roots, while dandelion roots concentrated up to 1360 µg Cd/g³⁷. The transport factors indicate that while heavy metals like Ni and Cu may preferentially accumulate in the foliage, highly toxic elements like Cd and Pb frequently deposit in the root cortex and caudex—the exact tissues utilized for coffee surrogates³⁷.

Because the European Food Safety Authority (EFSA) and the World Health Organization (WHO)

set extremely strict maximum permissible concentrations for heavy metals in herbal and food materials (e.g., 0.3 mg/kg for Cadmium, 10.0 mg/kg for Lead, 0.2 mg/kg for Mercury), sourcing roots from unverified urban community gardens, vacant lots, or roadside foraging in Berlin poses a severe physiological hazard to consumers³⁸. Consumption of heavy metal-laced botanicals disrupts ionic balance, stimulates oxidative damage to cellular DNA, and causes irreversible systemic toxicity⁴⁰. Therefore, commercial cultivation must be strictly relegated to tested, pristine agricultural fringes in Brandenburg, ensuring that baseline EDTA-extractable soil concentrations of heavy metals result in a Target Hazard Quotient (THQ) well below 1.0³⁹.

Post-Harvest Physiology: Fructan Depolymerization and Enzymatic Kinetics

The primary carbohydrate stored in the taproots of both chicory and dandelion is inulin, a non-digestible, linear β -(2,1)-linked fructan terminating in an α -D-glucopyranose residue⁴. The extraction efficacy, roasting thermodynamics, and dietary prebiotic properties of the final beverage are fundamentally dictated by the Degree of Polymerization (DP) of this inulin matrix at the exact moment of thermal processing⁴³.

The Biosynthesis and Degradation Cycle

Inulin synthesis during the vegetative growing phase is driven by a complex interplay of enzymes. Sucrose:sucrose 1-fructosyltransferase (1-SST) initiates the process by producing the trisaccharide 1-kestose, which is subsequently elongated by fructan:fructan 1-fructosyltransferase (1-FFT) to build massive, long-chain fructan polymers⁴⁴. Under optimal autumn conditions, industrial chicory roots can consist of 44% to 68% inulin by dry weight, with a high proportion of molecules exceeding a DP of 30⁴⁷.

However, the approach of the Brandenburg winter triggers a catastrophic biological shift within the root cortex. As ambient temperatures drop, the roots initiate cold acclimation to prevent cellular freezing damage. This cold stress drastically downregulates 1-FFT and massively upregulates the expression of the fructan 1-exohydrolase (1-FEH) enzyme family⁶.

1-FEH functions as a highly aggressive biological scissor, systematically depolymerizing the long-chain inulin molecules by cleaving terminal fructosyl units⁴⁹. This endogenous degradation acts as an osmoregulatory defense mechanism, flooding the cells with soluble sugars to lower the freezing point of the plant tissue⁴⁹. While essential for plant survival, this rapid enzymatic shift destroys the industrial and culinary quality of the root⁶. An increased concentration of low-DP fructooligosaccharides (FOS) and free fructose leads to excessive, uncontrollable caramelization, extreme hygroscopicity (stickiness), and hazardous toxicological spikes during the roasting phase⁶.

Storage Thermodynamics

To arrest 1-FEH activity and preserve the high-molecular-weight inulin fraction, harvest timing and post-harvest storage must be exactly controlled. Harvesting should occur in late autumn (October to November) before soil temperatures drop precipitously and trigger the systemic expression of the 1-FEH genes⁵³.

Crucially, storing the harvested roots at standard commercial refrigeration temperatures (e.g., 4°C) actively accelerates inulin depolymerization by maintaining high 1-FEH hydrolytic activity without supporting concurrent 1-FFT synthesis⁶. In practical terms, roots left in cold storage will rapidly convert their stable inulin banks into highly reactive free fructose⁵³. Conversely, immediately processing the roots, or flash-freezing them at -18°C, effectively denatures or completely halts the 1-FEH enzymatic pathways, locking the carbohydrate profile in its optimal, high-DP state⁵³.

Thermodynamics of Roasting: Pyrolysis, Caramelization, and Toxin Mitigation

The roasting process is the thermodynamic crucible wherein the raw, astringent botanical root is transmuted into a complex, deeply aromatic coffee surrogate. This transformation relies on specific kinetic reactions occurring between 135°C and 220°C, balancing the degradation of carbohydrates, the destruction of bitter secondary metabolites, and the mitigation of neurotoxic byproducts⁵⁶.

Inulin Scission and Caramelization

As the internal temperature of the dehydrated root fragments surpasses 135°C, dry heating induces the rapid thermal scission of the fructan chains⁵⁷. By 195°C, the native inulin matrix is almost entirely shattered, experiencing up to 100% degradation⁵⁷. The thermal breakdown of the β -(2,1) bonds results in the formation of complex low-molecular-weight products, predominantly di-D-fructose dianhydrides (DFAs), alongside free monosaccharides⁴³. Simultaneously, these released sugars undergo intense non-enzymatic browning and caramelization. This process generates massive quantities of 5-hydroxymethylfurfural (HMF), a highly reactive platform chemical that imparts the deep, sweet, and characteristic caramel-like visual and olfactory traits to the roast⁸.

Thermal Degradation of Sesquiterpene Lactones

The inherent, aggressive bitterness of chicory and dandelion roots stems from a high concentration of sesquiterpene lactones (SLs), specifically those possessing a rigid guaiane skeleton: lactucin, lactucopicrin, and 8-deoxylactucin, along with novel sulfate conjugates like 11 β ,13-dihydrolactucin-8-O-sulfate (jaquinelin-8-O-sulfate)⁴. While these compounds provide the necessary bitter baseline required to mimic coffee, in their raw state, their astringency is overpowering and unpalatable⁴.

Thermal processing facilitates the necessary breakdown of these molecules. At sustained roasting temperatures, the rigid lactone rings undergo structural cleavage, isomerization, and partial volatilization²⁰. Exhaustive chromatographic analyses indicate that careful washing protocols followed by precise thermal application can significantly reduce the absolute concentration of lactucin and lactucopicrin⁶⁴. This thermal degradation effectively tempers the harsh bitterness into a smooth, resonant profile that perfectly complements the sweetness of the DFAs and HMF²⁷.

Phenolic Acid Transformations

Dandelion and chicory roots are exceptional sources of hydroxycinnamic acid derivatives, though they differ in their primary constituents. Dandelion root is heavily dominated by chicoric acid (dicaFFEoyltartaric acid), an ester of two caffeic acids with tartaric acid, which acts as a potent antioxidant and anti-inflammatory agent⁵⁵. Chicory root, conversely, is characterized by high levels of chlorogenic acid (5-caffeoylquinic acid)⁶⁷.

During roasting, the applied thermal energy causes the progressive destruction and transformation of these polyphenols. Chlorogenic and chicoric acid levels drop significantly as they are thermally cleaved into free caffeic acid, quinic acid, and subsequently converted into chlorogenic acid lactones (CGLs)⁶⁹. Despite the loss of the parent phenolic molecules, the overall antioxidant capacity of the final brew often increases post-roast. This paradox is explained by the synthesis of melanoidins—high-molecular-weight, highly antioxidative brown polymers formed during the late stages of the Maillard reaction⁵⁶.

Acrylamide Generation and Mitigation Strategies

A severe toxicological risk associated with roasting carbohydrate-rich roots is the generation of acrylamide⁶¹. Acrylamide forms via a specific Maillard reaction pathway when reducing sugars (which are highly abundant in roots where 1-FEH has prematurely broken down inulin) react with the amino acid asparagine at temperatures exceeding 120°C⁶¹.

Because acrylamide is a known neurotoxin and is metabolically converted in the human body into glycidamide (a DNA-damaging mutagen), the European Union has established strict benchmark levels under Commission Regulation (EU) 2017/2158 for coffee substitutes⁶¹.

To aggressively mitigate acrylamide formation without sacrificing the complex roast profile, advanced pre-treatments must be employed. Research demonstrates that soaking chicory or

dandelion root pieces in a Calcium Chloride ($CaCl_2$) solution prior to dehydration and roasting fundamentally alters the Maillard kinetics, reducing final acrylamide concentrations by up to 40%⁶¹. Alternatively, utilizing enzymatic pre-treatments, such as incubating the roots with L-asparaginase derived from *Penicillium crustosum*, biochemically cleaves the asparagine precursor before heat is applied, reducing acrylamide formation by 75% to 80% without negatively impacting the structural retention of beneficial bioactive compounds⁶¹.

Phytochemical Component	Pre-Roast State	Post-Roast Transformation	Sensory / Health Implication
Inulin (DP > 30)	Long-chain β -(2,1) fructan polymer ⁴ .	Massive thermal scission into di-D-fructose dianhydrides (DFAs) and free fructose ⁴³ .	Generates sweetness; provides structural viscosity (mPa·s) to the brew ⁵⁷ .
Sesquiterpene	High concentrations of	Thermal ring cleavage and partial	Tempers overwhelming

Lactones	lactucin and lactucopicrin ⁴ .	degradation ²⁰ .	harshness into a smooth, coffee-like bitter baseline ²⁷ .
Phenolic Acids	Chicoric acid (Dandelion), Chlorogenic acid (Chicory) ⁶⁵ .	Cleaved into caffeic/quinic acids, forming Chlorogenic Acid Lactones (CGLs) ⁶⁹ .	Synthesizes highly antioxidative melanoidins ⁵⁶ .
Asparagine + Reducing Sugars	Natural amino acids and free fructose ⁶¹ .	Maillard conversion into Acrylamide (>120°C) ⁶¹ .	High toxicity risk requiring mitigation via <i>CaCl</i> ₂ or L-asparaginase ⁶¹ .

Volatile Metabolomics and the Olfactory Profile

The ultimate objective of roasting chicory and dandelion is not merely to darken the root tissue, but to synthesize a highly specific matrix of volatile organic compounds (VOCs). Gas chromatography-mass spectrometry (GC-MS) and advanced olfactometry studies (utilizing Odor Activity Values, OAV) reveal that the aroma of roasted chicory is governed by complex pyrolysis and Maillard derivatives, specifically pyrazines and furans⁷.

The Paramount Role of Rotundone

The singular molecule overwhelmingly responsible for the characterizing, authentic "chicory coffee" aroma is rotundone⁹. Rotundone is an oxygenated bicyclic sesquiterpene that imparts a deeply woody, spicy, and resonant peppery olfactory note⁹. It possesses an extraordinarily low odor detection threshold in aqueous solutions—frequently calculated at a mere 8 to 16 parts-per-trillion (pptr)⁷⁵.

Odor Activity Value (OAV) analysis—which mathematically quantifies the sensory impact by dividing a compound's absolute concentration by its odor detection threshold—unequivocally identifies rotundone as the most potent impact odorant in roasted chicory extracts⁷⁴.

Supporting Odorants and Flavor Architecture

The complex aroma profile generated by the roasted roots is rounded out by a suite of high-OAV caramel- and sweet-smelling odorants synthesized during the intense heat of roasting:

- **Sotolon (3-hydroxy-4,5-dimethyl-2(5H)-furanone):** A powerful furanone that imparts intense burnt sugar, fenugreek, and maple syrup notes⁹.
- **Dihydromaltol & Cyclotene:** These cycloenolones provide soft, sweet, caramelized nuances that mimic the inherent sweetness of lightly roasted Arabica beans⁹.
- **Substituted Pyrazines (e.g., 2-ethyl-3,5-dimethylpyrazine):** Deliver the crucial roasted, nutty, and earthy aromas fundamental to the structural backbone of any

credible coffee surrogate⁹.

Dandelion roots mirror this chemical profile, generating a deeply earthy, roasted aroma punctuated by similar pyrazines and trace volatile phenolic compounds⁸⁰. The precise blending of these two roots yields a full-spectrum olfactory experience that masterfully replicates the acidity, roast complexity, and body of genuine coffee.

Solid-Liquid Extraction Kinetics and Rheology

Brewing the roasted, ground roots requires a precise understanding of solid-liquid extraction mechanics, which differ fundamentally from those of traditional coffee. In standard coffee brewing, the target Extraction Yield (EY)—the percentage of dry mass dissolved into the cup—is strictly bounded between 18% and 22%⁸². This EY results in Total Dissolved Solids (TDS) ranging from 1.15% to 1.45% for gravity-fed filter coffee, and 8% to 12% for high-pressure espresso⁸².

Because chicory and dandelion roots lack the highly recalcitrant, woody cellulose structure of coffee beans and are instead primarily composed of highly water-soluble carbohydrates (residual inulin and DFAs), their solubility kinetics are exceptionally rapid⁴⁷. The grounds are highly extractable, meaning that utilizing standard coffee brew ratios (e.g., a 1:15 ratio for filter, or prolonged immersion in a French Press) will rapidly force the extraction yield well past the 22% threshold⁸². Pushing the extraction yield too high dissolves the last, most recalcitrant phenolic compounds and residual sesquiterpene lactones, resulting in a hollow, overwhelmingly dry, and intensely bitter astringency⁸³.

To optimize the brew, a slightly coarser grind size and shorter contact times are mandatory. High-pressure, short-duration extraction via a Moka pot or a calibrated espresso machine is ideal, as it halts the extraction before the bitterest phenolic fractions overwhelm the cup, maximizing the sweet, viscous DFAs⁸².

Fluid Rheology and Viscosity

A unique physical characteristic of the resulting chicory and dandelion extract is its elevated rheological viscosity. Even after enduring thermal degradation, the residual short-chain inulin, fructooligosaccharides, and DFAs significantly increase the viscosity (measured in milliPascal-seconds, mPa·s) of the aqueous continuous phase⁷². This high concentration of complex carbohydrates acts as a natural emulsifier, dramatically reducing droplet mobility and stabilizing the formation of foam⁸⁶. This specific trait perfectly mimics the thick, luxurious mouthfeel of true espresso crema, an attribute highly prized in high-end culinary preparations⁸².

Gastrointestinal Implications: The Prebiotic and FODMAP Duality

The biological impact of the brewed root beverage on the human gastrointestinal tract is profound, exerting a powerful, dual-natured influence depending entirely on the consumer's baseline microbiome and gut health.

Prebiotic Health Claims and SCFA Synthesis

For individuals with a healthy gastrointestinal tract, the beverage acts as an exceptional functional food. Native chicory inulin and its complex FOS derivatives resist enzymatic

hydrolysis by human salivary and small intestinal enzymes due to the β -configuration of their anomeric C2 bonds⁸⁷. Consequently, these carbohydrates reach the large intestine virtually intact, where they are selectively fermented by beneficial gut microbiota, notably *Bifidobacterium*, *Lactobacillaceae*, and *Anaerostipes*⁸⁷.

This targeted bacterial fermentation yields high concentrations of Short-Chain Fatty Acids (SCFAs), primarily butyrate, acetate, and propionate⁹¹. Butyrate is particularly critical; it lowers colonic pH, acts as the primary energy source for colonocytes, stimulates the intestinal epithelial barrier by regulating tight junction proteins (TJPs), and aggressively promotes the clearance of bacterial pathogens⁸⁸. Based on exhaustive clinical trials demonstrating these physiological mechanics, the European Food Safety Authority (EFSA) authorized an exclusive Article 13.5 health claim confirming that a daily intake of 12 grams of native chicory inulin "contributes to normal bowel function by increasing stool frequency" and increasing overall stool softness⁹⁰.

The FODMAP Intolerance Paradox

Conversely, inulin is formally classified as a highly fermentable oligo-, di-, monosaccharide, and polyol (FODMAP)⁹³. In individuals suffering from Irritable Bowel Syndrome (IBS), Small Intestinal Bacterial Overgrowth (SIBO), or general fructose malabsorption, the rapid osmotic pull and aggressive fermentation of these fructans by dysbiotic bacteria generate excessive gaseous distension⁹³.

This manifests clinically as acute bloating, severe abdominal pain, and unpredictable, explosive motility⁹³. For this demographic, a low-FODMAP threshold strictly limits chicory or dandelion root consumption to ≤ 0.5 g per serving, rendering concentrated, high-TDS espresso-style brews highly problematic⁹⁵. Consequently, while the beverage acts as a potent, scientifically validated digestive tonic for a healthy gut, careful dose control and consumer awareness are mandatory to prevent severe GI distress in sensitive populations⁹⁵.

Culinary Application Engineering: Tiramisu Construction in Berlin

The integration of concentrated chicory and dandelion root extract into the classic Italian dessert Tiramisu requires specific structural, rheological, and sensory adaptations compared to traditional *Coffea arabica* preparations. A classic Tiramisu architecture relies on the precise capillary saturation of Savoiardi (ladyfingers)—a highly porous, low-moisture, egg-and-flour sponge biscuit—with a bitter liquid, which is subsequently layered with a lipid-dense, whipped mascarpone cream¹³.

Capillary Action and Saturation Mechanics

Because the dandelion/chicory brew exhibits a significantly higher dynamic viscosity than standard coffee due to the heavy suspension of polysaccharide hydrocolloids (inulin and DFAs), the fluid dynamics of the biscuit soak are fundamentally altered¹³. The highly viscous liquid

permeates the open cellular structure of the Savoiardi via capillary action at a heavily reduced velocity.

If the culinary technician utilizes the standard rapid-dip protocol used for fluid espresso, the high-viscosity surrogate will fail to penetrate the core of the biscuit, leaving a dry, unpalatable, chalky center¹³. Conversely, extending the soak time excessively will cause the hydrocolloids to over-saturate and chemically disintegrate the fragile starch matrix, resulting in structural collapse and a mushy texture¹³. Therefore, the brew must either be applied via controlled, repeated brushing to ensure even saturation, or slightly diluted with warm water to explicitly lower the viscosity coefficient prior to dipping¹³.

Sensory Balancing and Physiological Benefits

From a sensory and physiological perspective, the chicory-dandelion blend is exceptionally well-suited for high-end Tiramisu. The total absence of caffeine ensures the dessert can be consumed in the late evening without antagonizing adenosine receptors in the brain. Caffeine directly blocks adenosine—the chemical that signals the brain to initiate sleep—leading to disrupted circadian architectures and delayed sleep onset, a frequent clinical complaint associated with traditional espresso-soaked Tiramisu⁹⁸.

Flavor-wise, the aggressive, earthy bitterness provided by the thermally degraded sesquiterpene lactones acts as an essential palate cleanser, cutting through the heavy, lipid-dense profile of the mascarpone (or modern vegan cashew/aquafaba cream equivalents)¹³. Furthermore, the high concentration of the impact odorant rotundone imparts a complex, woody, and spicy depth that perfectly bridges the gap between the rich dairy fats and the final, bitter dusting of alkalized cocoa powder, elevating the dessert beyond a mere caffeine-free substitute into a highly sophisticated, regionally authentic culinary experience⁹.

Works cited

1. Chicory Market Size, Growth, Share, Trends and Forecast 2032, <https://www.credenceresearch.com/report/chicory-market>
2. Ground Chicory Market Size, Share | CAGR of 6.5%, <https://market.us/report/global-ground-chicory-market/>
3. Chicory Market Size, Share | CAGR of 6.8%, <https://market.us/report/global-chicory-market/>
4. (PDF) Sesquiterpene lactones and inulin from chicory roots, https://www.researchgate.net/publication/40209401_Sesquiterpene_lactones_and_inulin_from_chicory_roots_extraction_identification_enzymatic_release_and_sensory_analysis
5. Bioactive Compounds from Dandelion (*Taraxacum officinale*) - MDPI, <https://www.mdpi.com/2304-8158/15/4/782>
6. Differential expression of fructan 1-exohydrolase genes involved in, https://www.researchgate.net/publication/232755713_Differential_expression_of_fructan_1-exohydrolase_genes_involved_in_inulin_biodegradation_in_chicory_Cichorium_intybus_cultivars
7. Quality and Flavour Stability of Coffee Substitute Prepared by,

- https://www.researchgate.net/publication/6661348_Quality_and_Flavour_Stability_of_Coffee_Substitute_Prepared_by_Extrusion_of_Wheat_Germ_and_Chicory_Roots
8. Use of NMR-Based Metabolomics To Chemically Characterize the, <https://pubs.acs.org/doi/10.1021/acs.jafc.6b02423>
 9. Identification of Characterizing Aroma Components of Roasted, https://www.researchgate.net/publication/332339637_Identification_of_Characterizing_Aroma_Components_of_Roasted_Chicory_Coffee_Brews
 10. 24 Berlin Crop Rotation Schedule.pdf
 11. How to Grow Dandelion from Seed: Planting & Care, <https://organorepublic.com/blogs/growing-guides/dandelion>
 12. Tour Leader Treviso, <https://www.tourleadertreviso.com/section/home/>
 13. Vegan Tiramisu - Supper in the Suburbs, <https://supperinthesuburbs.com/2011/07/17/vegan-tiramisu/>
 14. From Seed to Harvest: A beginner's guide to growing chicory, <https://files.ufseeds.com/on/demandware.static/Sites-UrbanFarmer-Site/Sites-UrbanFarmer-Library/default/images/Instructions-for-Growing-Chicory.pdf>
 15. Chicory – Growing Guide - Deep Green Permaculture, <https://deepgreenpermaculture.com/2026/03/07/chicory-growing-guide/>
 16. Italian Dandelion Chicory | Key Growing Information, <https://www.johnnyseeds.com/growers-library/vegetables/chicory/italian-dandelion-chicory-key-growing-information.html>
 17. How to Grow Chicory in Your Garden - USU Extension, <https://extension.usu.edu/yardandgarden/research/chicory-in-the-garden>
 18. Innovative Applications and Research Methods in Agriculture, <https://img1.wsimg.com/blobby/go/188e75a3-9b73-49dc-8938-8655ce331291/downloads/819f77a7-4609-4658-adaf-70ac2be77161/Innovative%20Applications%20and%20Research%20Methods.pdf?ver=1766740858117>
 19. WOCMAP ABSTRACTS ISSN 0189-6016©2009 - ATHMSI Journals, <https://journals.athmsi.org/index.php/ajtcam/article/download/602/440/0>
 20. WO2002013625A2 - Non-bitter chicory extract - Google Patents, <https://patents.google.com/patent/WO2002013625A2/en>
 21. Home Herbal_ The Ultimate Guide to Cooking, Brewing, and, https://pubhtml5.com/zpek/qakw/Home_Herbal_The_Ultimate_Guide_to_Cooking%2C_Brewing%2C_and_Blending_Your_Own_Herbs/
 22. Growing chicory: from seed to fresh winter harvest - Plukkers.com, <https://plukkers.com/en/blogs/gewasinfo/witlof-kweken>
 23. roasted coffee beans: Topics by Science.gov, <https://www.science.gov/topicpages/r/roasted+coffee+beans.html>
 24. International Journal of Agriculture, Environment and Food Sciences, <https://dergipark.org.tr/tr/download/article-file/4115410>
 25. Bernadetta Lis - BIP, https://www.bip.uni.lodz.pl/fileadmin/user_upload/ROZPRAWA_DR_B.L_.pdf
 26. Asteraceae Family: Health Benefits & Uses | PDF | Lettuce - Scribd, <https://www.scribd.com/document/788223070/The-family-Asteraceae-or-Comp>

[ositae-docx-FINAL](#)

27. Relationship between Levels of Sesquiterpene Lactones in Chicory,
https://www.researchgate.net/publication/40152977_Relationship_between_Levels_of_Sesquiterpene_Lactones_in_Chicory_and_Sensory_Evaluation
28. inulin-type fructan fermentation: Topics by Science.gov,
<https://www.science.gov/topicpages/i/inulin-type+fructan+fermentation.html>
29. How to grow chicory - Gardeners' World,
<https://www.gardenersworld.com/how-to/grow-plants/how-to-grow-chicory/>
30. effects of plant density on root yield and leaf area in chicory,
https://www.researchgate.net/publication/252322400_Effects_of_Plant_Density_on_Root_Yield_and_Leaf_Area_in_Chicory_Cichorium_intybus_L
31. Taraxacum officinale in meadow fodder production,
<http://ir.polissiauniver.edu.ua/handle/123456789/15277>
32. Cultivation of dandelion (Taraxacum erythropodium) on coastal,
<https://scispace.com/pdf/cultivation-of-dandelion-taraxacum-erythropodium-on-coastal-3qrcp75vnm.pdf>
33. 36th Annual Conference & Meeting,
https://aaic.org/wp-content/uploads/2025/08/AbstractBook_2025AAICIOCC.pdf
34. Dandelion Handbook | Marburg - iGEM 2024,
<https://2024.igem.wiki/marburg/handbook/implementation>
35. Planting Density and Growth Cycle Affect Actual and Potential Latex,
https://www.researchgate.net/publication/334975672_Planting_Density_and_Growth_Cycle_Affect_Actual_and_Potential_Latex_and_Rubber_Yields_in_Taraxacum_kok-saghyz
36. Natural Lead Levels in Dandelions (Taraxacum officinale): A Weed,
<https://pubs.acs.org/doi/10.1021/acs.est.8b04191>
37. (PDF) Chicory (Cichorium intybus L.) and dandelion (Taraxacum,
https://www.researchgate.net/publication/226887516_Chicory_Cichorium_intybus_L_and_dandelion_Taraxacum_officinale_Web_as_phytoindicators_of_cadmium_contamination
38. Heavy Metals in Medicinal Plants - Encyclopedia.pub,
<https://encyclopedia.pub/entry/41334>
39. (PDF) Bioaccumulation of Heavy Metals by Herbaceous Species,
https://www.researchgate.net/publication/359887983_Bioaccumulation_of_Heavy_Metals_by_Herbaceous_Species_Grown_in_Urban_and_Rural_Sites
40. The Safety Assessment of Toxic Metals in Commonly Used Herbs,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8197804/>
41. Medicinal Plant Growth in Heavy Metals Contaminated Soils - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9504071/>
42. A simplified spectrophotometric method for the determination of,
https://www.researchgate.net/publication/225159679_A_simplified_spectrophotometric_method_for_the_determination_of_inulin_in_Jerusalem_artichoke_Helianthus_tuberosus_L_tubers
43. Determination of Inulin-Type Fructooligosaccharides in Edible,
https://www.researchgate.net/publication/264091099_Determination_of_Inulin-Ty

- [pe_Fructooligosaccharides_in_Edible_Plants_by_High_Performance_Liquid_Chromatography_with_Charged_Aerosol_Detector](#)
44. Water stress drastically reduces root growth and inulin yield in, <https://academic.oup.com/jxb/article/63/12/4359/640759>
 45. Properties of Fructan:Fructan 1-Fructosyltransferases from Chicory, https://www.researchgate.net/publication/277550259_Properties_of_FructanFructan_1-Fructosyltransferases_from_Chicory_and_Globe_Thistle_Two_Asteracean_Plants_Storing_Greatly_Different_Types_of_Inulin
 46. Fructan metabolism and changes in fructan composition during cold, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2015.00329/full>
 47. January 2018 - naturalherbssite, <https://naturalherbssite.wordpress.com/2018/01/>
 48. Metabolic Profiling of Sugarloaf Chicory Roots - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12942909/>
 49. (PDF) Fructan 1-exohydrolase is associated with flower opening in, https://www.researchgate.net/publication/228675766_Fructan_1-exohydrolase_is_associated_with_flower_opening_in_Campanula_rapunculoides
 50. Effect of defoliation on fructan pattern and fructan metabolizing, https://www.researchgate.net/publication/229466433_Effect_of_defoliation_on_fructan_pattern_and_fructan_metabolizing_enzymes_in_young_chicory_plants_Cichorium_intybus
 51. Fructan:fructan 1-fructosyltransferase and inulin hydrolase activities, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5334228/>
 52. Evolutionary origins and natural functions of fructans—a, https://www.researchgate.net/publication/230066738_Evolutionary_origins_and_natural_functions_of_fructans-a_climatological_biogeographic_and_mechanistic_appraisal
 53. Inulin and Sugar Contents in Helianthus tuberosus and Cichorium, https://www.researchgate.net/publication/230135225_Inulin_and_Sugar_Contents_in_Helianthus_tuberosus_and_Cichorium_intybus_Tubers_Effect_of_Postharvest_Storage_Temperature
 54. Influence of plant age on fructan content and fructosyltransferase, <https://academicjournals.org/journal/AJB/article-full-text-pdf/9282C8430466>
 55. A Study of Variation in the Major Phenolic Acid Components ... - MDPI, <https://www.mdpi.com/1467-3045/47/3/145>
 56. How do coffee substitutes compare to coffee? A comprehensive, https://www.researchgate.net/publication/353748594_How_do_coffee_substitutes_compare_to_coffee_A_comprehensive_review_of_its_quality_characteristics_sensory_characters_phytochemicals_health_benefits_and_safety
 57. Heat-induced degradation of inulin - ResearchGate, https://www.researchgate.net/publication/225574467_Heat-induced_degradation_of_inulin
 58. Thermal degradation and consequent fragmentation of widely, https://www.researchgate.net/publication/271574398_Thermal_degradation_and_consequent_fragmentation_of_widely_applied_oligosaccharides

59. Nut Free Cookies, <https://mariamindbodyhealth.com/nut-free-cookies/>
60. Application of in UI in in Bread Are View of Technological Properties, <https://www.scribd.com/document/624149953/Application-of-in-UI-in-in-Bread-A-re-View-of-Technological-Properties>
61. Coffee and chicory blend: properties, nutrition, and health implications, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12174007/>
62. Bioactive, Mineral and Antioxidative Properties of Gluten-Free ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9688964/>
63. bitter sesquiterpene lactones: Topics by Science.gov, <https://www.science.gov/topicpages/b/bitter+sesquiterpene+lactones>
64. Sesquiterpene Lactone Content and Overall Quality of Fresh-Cut, <https://pubs.acs.org/doi/10.1021/jf402189v>
65. Bioactive Phenolics and Antioxidant Capacity of Some Wild Edible, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7554971/>
66. Chicoric acid: chemistry, distribution, and production - Frontiers, <https://www.frontiersin.org/journals/chemistry/articles/10.3389/fchem.2013.00040/pdf>
67. Health Benefits of Key Constituents in Cichorium intybus L. - MDPI, <https://www.mdpi.com/2072-6643/15/6/1322>
68. Bioactive, Mineral and Antioxidative Properties of Gluten-Free, <https://www.mdpi.com/2304-8158/11/22/3692>
69. Showing report on Non-alcoholic beverages - Phenol-Explorer, <http://phenol-explorer.eu/reports/37>
70. Flavour formation in the coffee substitute 'lupin coffee' related ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12397916/>
71. acrylamide content in foods and factors that - EMU DSpace, <https://dspace.emu.ee/bitstreams/84da8949-a560-4d99-958a-d2d7a6aaff30/download>
72. Enzymatic High Fructose Syrup Production | PDF | Fructose | Glucose, <https://www.scribd.com/document/898478541/B-Chap-HFS-2011>
73. Identification of Aroma Active Compounds of Cereal Coffee Brew, https://www.researchgate.net/publication/235646737_Identification_of_Aroma_Active_Compounds_of_Cereal_Coffee_Brew_and_Its_Roasted_Ingredients
74. Identification and quantitation of predominant odorants in roasted, <https://www.ideals.illinois.edu/items/103497>
75. Effective analysis of rotundone at below-threshold levels in red and, https://www.researchgate.net/publication/49778439_Effective_analysis_of_rotundone_at_below-threshold_levels_in_red_and_white_wines_using_solid-phase_microextraction_gas_chromatographytandem_mass_spectrometry
76. (PDF) Identification of Rotundone as an Important Contributor to the, https://www.researchgate.net/publication/353358353_Identification_of_Rotundone_as_an_Important_Contributor_to_the_Flavor_of_Oak-Aged_Spirits
77. Identification of Characterizing Aroma Components of Roasted, <https://pubs.acs.org/doi/abs/10.1021/acs.jafc.9b00776>
78. Introduction to the Second International Flavor and Fragrance,

- <https://pubs.acs.org/doi/10.1021/acs.jafc.9b04188>
79. Identification of Dihydromaltol (2,3-Dihydro-5-hydroxy-6-methyl-4H,
https://www.researchgate.net/publication/26883292_Identification_of_Dihydromaltol_23-Dihydro-5-hydroxy-6-methyl-4H-pyran-4-one_in_Ryazhenka_Kefir_and_Comparative_Sensory_Impact_Assessment_of_Related_Cycloenolones
 80. NUTRITION AND HEALTH BENEFITS OF PURE MAPLE SYRUP,
<https://www.breedonsmaplesyrup.com/pdf/IMSI.pdf>
 81. Nutrition and Health Benefits of Pure Maple Syrup PDF - Scribd,
<https://www.scribd.com/document/407246371/nutrition-and-health-benefits-of-pure-maple-syrup-pdf>
 82. The Golden Ratios | Coffee Extraction Science, <https://decimalcoffee.com/ratios>
 83. Extraction Yield in Coffee: What It Is and How to Measure It,
<https://podiumcoffeeclub.com/blogs/blog/coffee-extraction-yield>
 84. Measuring Coffee Extraction (And Why It Makes People Mad) - Biocaf,
<https://www.biocaf.com/post/measuring-coffee-extraction-and-why-it-makes-people-mad>
 85. Coffee Extraction Calculator | TDS & Yield Estimator - ROUTES,
<https://routescoffee.co.uk/interactive-brewing-ratio/>
 86. Techno-Functional Properties and Applications of Inulin in Food,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12564896/>
 87. Inulin-Type Prebiotics - A Review: Part 1 - ResearchGate,
https://www.researchgate.net/publication/23800195_Inulin-Type_Prebiotics_-_A_Review_Part_1
 88. Introducing inulin-type fructans - ResearchGate,
https://www.researchgate.net/publication/7862802_Introducing_inulin-type_fructans
 89. Authorised EU health claim for chicory inulin | Request PDF,
https://www.researchgate.net/publication/322182489_Authorised_EU_health_claim_for_chicory_inulin
 90. Chicory inulin – what it is and how it affects digestion? - NutriDream,
<https://nutridream.eu/en/2024/10/13/chicory-inulin-what-it-is-and-how-it-affects-digestion/>
 91. digestive-health - CEFI,
<https://www.chicoryinulin.eu/research-topics/digestive-health>
 92. Dried chicory root improves bowel function, benefits intestinal,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11407914/>
 93. Like many, I am suspicious of chicory root fiber : r/FODMAPS - Reddit,
https://www.reddit.com/r/FODMAPS/comments/18x6qbt/like_many_i_am_suspicious_of_chicory_root_fiber/
 94. EFSA Fiber Health Claims: Which Are Real (and Why 9 of 10 Fail),
<https://goodfibercompany.com/blog/efsa-health-claims-fiber-explained/>
 95. Starbucks vs Dunkin': chicory benefits & drawbacks (2026),
<https://foodreveal.app/en/blog/starbucks-vs-dunkin-chicory-benefits-drawbacks>
 96. Cichorium Intybus in Ayurveda: Benefits, Uses & Healing Power,
<https://ask-ayurveda.com/wiki/article/4017-cichorium-intybus>

97. Cakes PDF | PDF | Cakes | Flour - Scribd,
<https://www.scribd.com/document/488778618/Cakes-pdf>
98. Tiramisu Caffeine Content: Will It Keep You Awake? - CAFELY,
<https://cafely.com/blogs/info/does-tiramisu-have-caffeine>