

A Comprehensive Botanical and Ethnobotanical Guide to Wild Edibles and Medicinal Roots in the Berlin-Brandenburg Region

1. The Legislative and Ecological Frameworks Governing Botanical Foraging

The extraction of wild plant materials in the Berlin and Brandenburg regions is mediated by a complex matrix of federal conservation laws, state-specific ecological directives, and precise seasonal restrictions. Understanding this legal architecture is an absolute prerequisite for any botanical, ethnobotanical, or culinary foraging endeavor, ensuring that human utilization does not compromise regional biodiversity or violate property and conservation statutes.

1.1 The Federal Hand Bouquet Rule (Handstraußregelung)

At the federal level, the fundamental legislation governing the interaction with wild flora is the Bundesnaturschutzgesetz (BNatSchG), particularly § 39. The baseline legal stance, as articulated in § 39 Abs. 1 BNatSchG, is that it is strictly prohibited to remove wild plants from their natural locations without reasonable cause or to devastate their populations¹. However, the law provides a vital exemption for non-commercial, personal foraging known as the "Handstraußregelung" (hand bouquet rule). Defined under § 39 Abs. 3 BNatSchG, this provision permits individuals to carefully harvest small quantities of wild flowers, grasses, ferns, mosses, lichens, fruits, fungi, and tea or medicinal herbs for personal use ("Eigenbedarf") from areas that are generally accessible to the public and not subject to specific protective designations¹. The concept of a "small quantity" is a qualitative legal standard, generally interpreted as an amount that can be reasonably held between the thumb and index finger, while for berries or mushrooms, it translates to a modest basket intended exclusively for the immediate personal consumption of the forager and their household¹. The legislation explicitly excludes any commercial harvesting of forest products. Individuals intending to harvest wild plants, fungi, or wood for commercial sale, or in volumes exceeding the personal use threshold, must obtain the explicit consent of the landowner and a collection permit (such as a "Holzsammelschein") from the lower nature conservation authority¹.

1.2 The Absolute Prohibition on Root Extraction and Woody Plants

A critical limitation of the Handstraußregelung is its strict boundary regarding the physical integrity of the plant organism and its surrounding soil matrix. The exemption applies solely to the aerial parts of herbaceous plants. It does not permit the uprooting of plants, as the harvesting of roots, bulbs, or tubers inevitably results in the death of the specimen and significantly disrupts the ecosystem⁶. Consequently, the extraction of valuable medicinal roots—such as chicory, dandelion, or horseradish—is explicitly excluded from the general

foraging exemption⁶. The extraction or severe pruning of verwooded plants, including trees and shrubs, is similarly prohibited⁶.

From a legal perspective, every plant, including its subterranean structures, is the property of the respective landowner. Removing a root without explicit authorization from the landowner (whether a private entity or the state forestry commission) constitutes theft and property damage under the Strafgesetzbuch (Criminal Code)⁵. Fines for such violations, especially within protected zones or involving protected species, can escalate dramatically, theoretically reaching up to €50,000 for severe infractions under the BNatSchG¹.

1.3 State Conservation Laws and Protected Species (BArtSchV)

Geographic zoning plays a definitive role in the legality of foraging. The Handstraußregelung is nullified on agricultural lands, which are heavily managed and frequently subject to the application of pesticides and herbicides, as well as on private, fenced properties¹. In the states of Berlin and Brandenburg, foraging is unequivocally banned within Naturschutzgebiete (Nature Reserves), National Parks, and the core zones of Biosphere Reserves, such as the Spreewald or Schorfheide-Chorin⁷. Furthermore, the Berliner Naturschutzgesetz (NatSchG Bln) emphasizes the protection of a "Biotopverbund" (habitat network), aiming to secure at least 10% of the state's surface area to ensure the regenerative capacity of natural goods and to prevent soil erosion¹⁰.

The protection status of specific plant species supersedes all general foraging rights. Species listed under the Bundesartenschutzverordnung (BArtSchV) or the European Habitats Directive (FFH) as specially or strictly protected may not be damaged, picked, or collected under any circumstances¹. Regional variations in these lists are of paramount importance. For example, while wild garlic (*Allium ursinum*) is relatively abundant and freely foraged in southern Germany, it is categorized as highly endangered on the Rote Liste (Red List) for the state of Brandenburg (Category 1)¹³. Consequently, harvesting *Allium ursinum* in the wild in Brandenburg is strictly forbidden to protect the fragile, fragmented populations¹⁶.

1.4 Seasonal Protections for Avian and Terrestrial Wildlife

Both state and federal laws impose stringent seasonal restrictions designed to protect the breeding and nesting habitats of wildlife. Under § 39 Abs. 5 Nr. 2 BNatSchG and corresponding sections of the Berliner Naturschutzgesetz, it is strictly prohibited to prune, fell, clear, or destroy hedges, living fences, bushes, and other woody vegetation between March 1st and September 30th⁶. While these statutes primarily regulate arboriculture and landscaping, they underscore the acute ecological sensitivity of the spring and summer months, mandating a respectful and non-disruptive approach from foragers navigating forest edges and understories during the peak breeding season⁶.

2. Environmental Toxicology and Zoonotic Risks in Urban Foraging

The geographical layout of Berlin and Brandenburg—where dense urban infrastructure intimately interfaces with extensive forests, lakes, and ruderal wastelands—offers unparalleled foraging access. However, this peri-urban environment presents specific toxicological and

parasitological vectors that must be systematically managed by anyone consuming wild flora.

2.1 Heavy Metal Bioaccumulation in Urban and Ruderal Soils

Urban soils frequently act as sinks for elevated levels of heavy metals (HMs) and metalloids, driven by historical industrial activities, vehicular emissions (including tire abrasion and fuel additives), and construction debris¹⁹. Common urban soil pollutants include lead (Pb), zinc (Zn), copper (Cu), cadmium (Cd), and chromium (Cr)²⁰.

Plants exhibit varying, species-specific capacities for phytoextraction and bioaccumulation. Ruderal species that are ubiquitous in Berlin, such as dandelion (*Taraxacum officinale*), stinging nettle (*Urtica dioica*), and ribwort plantain (*Plantago lanceolata*), possess highly adaptable root systems capable of absorbing heavy metals from contaminated matrices²¹. The distribution of these heavy metals within the plant anatomy is not uniform. For instance, elements like cadmium, copper, and nickel preferentially accumulate in the foliar tissues of herbaceous plants, whereas chromium often concentrates in the floral structures²¹. Conversely, lead tends to be sequestered in the root system and lower stalks due to its low mobility within the plant's vascular tissue²¹.

The chronic ingestion of heavy metals via the food web poses severe health risks to humans, inducing oxidative stress and contributing to neurological, cardiovascular, and renal pathologies¹⁹. Therefore, foraging in close proximity to high-traffic arteries, former industrial brownfields, or railway corridors is highly discouraged²⁰. Identifying uncontaminated rural or deeply forested collection sites is an essential prerequisite for the safe, long-term consumption of wild herbs and roots²².

2.2 Zoonotic Risks: *Echinococcus multilocularis* (Alveolar Echinococcosis)

A paramount biological risk in Central European foraging is the potential exposure to the microscopic eggs of the fox tapeworm (*Echinococcus multilocularis*), the causative agent of alveolar echinococcosis in humans²⁶. The adult tapeworm, measuring a mere two to six millimeters, resides in the small intestine of its definitive hosts—predominantly the red fox (*Vulpes vulpes*), which maintains a robust population density in Berlin's urban spaces, as well as raccoon dogs, and occasionally domestic dogs and felines²⁶.

These definitive hosts excrete tapeworm proglottids, each containing hundreds of eggs, into the environment via their feces. These eggs exhibit extreme environmental resilience, surviving for up to 190 days in the soil and remaining viable through extreme winter temperatures²⁶.

Humans act as aberrant intermediate hosts (Fehlzwischenwirt) if they accidentally ingest the eggs via contaminated soil, low-hanging wild berries, or low-growing herbs such as wild garlic or ground ivy²⁶.

Upon ingestion, the eggs hatch in the human gastrointestinal tract. The released larvae penetrate the intestinal mucosa and migrate via the portal vein primarily to the liver (affected in 98% of cases), where they proliferate slowly in a tumor-like, infiltrative manner²⁶. The incubation period is remarkably prolonged, often ranging from 10 to 15 years before the onset of clinical symptoms, which include cholestasis (manifesting as jaundice), abdominal pain,

portal hypertension, and hepatomegaly²⁷. Unbehandelt, the disease is progressive and highly lethal due to severe hepatic destruction²⁷.

According to the Robert Koch Institute (RKI), the absolute number of cases is relatively low—approximately 41 reported cases in Germany in 2022—but the severity of the disease demands rigorous prophylactic measures²⁷. Demographic analyses indicate that agricultural workers (representing 21.9% of cases), outdoor professionals, and pet owners (70% of cases) bear the highest risk burden²⁶.

To mitigate the risk of *E. multilocularis*, wild-foraged herbs and berries collected near the ground must be meticulously washed. However, washing alone is insufficient to guarantee the removal of the highly adhesive eggs²⁶. The only definitive method to destroy the eggs is thermal processing. Cooking or baking wild edibles at temperatures between 70°C and 80°C for at least five minutes ensures the complete denaturation of the parasite²⁶. Conversely, standard domestic freezing (-20°C) is entirely ineffective; the eggs maintain viability until subjected to sustained temperatures of -80°C for several days, a threshold unreachable by standard consumer appliances²⁶.

3. Invasive Neophytes: Foraging as Conservation

While native plants are subject to strict conservation laws, the Berlin and Brandenburg regions host several invasive neophytes. These non-native species often outcompete indigenous flora, altering soil chemistry and shading out native understories. Strategically foraging these invasive species presents a unique opportunity to harvest abundant culinary and medicinal resources while actively participating in local ecological management.

3.1 Garlic Mustard (*Alliaria petiolata*)

Alliaria petiolata, known locally as Knoblauchsrauke, is a widespread biennial herb of the Brassicaceae family³². Originating in Eurasia, it has become a dominant force in forest understories and edge habitats³². During its first year, the plant forms a basal rosette of dark green, reniform (kidney-shaped) leaves. In its second year, it bolts rapidly, producing an erect stem up to one meter in height, bearing alternate, deltoid to cordate leaves, and terminal racemes of small, white, tetramerous flowers³².

The distinctive ecological and culinary signature of *A. petiolata* relies on the glucosinolate-myrosinase defense system, a defining biochemical characteristic of the Brassicaceae order³². The plant synthesizes the glucosinolate sinigrin, a secondary metabolite derived from the amino acid methionine, which is stored safely within cellular vacuoles³⁶. In separate cellular compartments, the plant stores the hydrolytic enzyme myrosinase³⁵. When the plant tissue is mechanically damaged—whether by insect herbivory or by a human crushing the leaves—the enzyme and substrate mix³⁵. Myrosinase rapidly hydrolyzes sinigrin to produce volatile degradation products, primarily allyl isothiocyanate (AITC), which imparts the sharp, pungent, garlic-and-mustard aroma and flavor³².

From an ecological perspective, these isothiocyanates exhibit potent allelopathic properties. *A. petiolata* exudes these compounds into the surrounding soil, where they actively inhibit the growth and colonization of ectomycorrhizal fungi³². Because native tree seedlings and forest

flora depend heavily on these fungal symbioses for nutrient acquisition, the suppression of the fungi allows garlic mustard to become a highly invasive competitor, creating monocultures in the forest understory³².

Historically, garlic mustard was utilized extensively in folk medicine. The high concentrations of isothiocyanates provide robust antibacterial and antiseptic qualities, making it a traditional treatment for gangrene, putrid ulcers, and respiratory catarrh³⁹. Internally, it acts as a diaphoretic, expectorant, and mild vermifuge³⁹.

For modern culinary applications, the plant is best utilized fresh and raw, as it is rich in vitamins A, C, and E³⁴. The myrosinase enzyme is highly sensitive to thermal degradation. While its optimum catalytic activity occurs between 45°C and 55°C, the enzyme denatures rapidly at temperatures exceeding 75°C⁴¹. Cooking garlic mustard destroys the enzyme, preventing the formation of AITC and causing the desirable pungent flavor to dissipate entirely, often leaving behind a slightly bitter residual taste³⁴.

3.2 Himalayan Balsam (*Impatiens glandulifera*) and Jerusalem Artichoke (*Helianthus tuberosus*)

In the riparian zones and alluvial floodplains of Brandenburg, two other prominent neophytes have established dense populations: Himalayan Balsam (*Impatiens glandulifera*) and Jerusalem Artichoke or Topinambur (*Helianthus tuberosus*)⁴².

Impatiens glandulifera is an annual that grows rapidly to heights exceeding two meters, producing large, orchid-like pink flowers. It spreads aggressively along waterways, displacing native vegetation⁴⁴. The seeds, which are propelled explosively from mature pods, possess a nutty flavor and can be harvested in late summer.

Helianthus tuberosus, a member of the Asteraceae family, is a tall perennial that produces bright yellow, sunflower-like blooms late in the season⁴². It propagates vigorously through subterranean stoloniferous tubers⁴⁶. These tubers, rich in the carbohydrate inulin, are highly nutritious. While the extraction of roots is generally prohibited for native species, harvesting the tubers of *H. tuberosus* where it has become invasive along riverbanks is often encouraged as a method of population control, though foragers should still verify local regulations⁴³.

4. Botanical Profiles: Key Native Herbs and Safe Identification

Accurate botanical identification is the absolute cornerstone of safe foraging. The morphological divergence between edible species and their highly toxic sympatric counterparts can be remarkably subtle, necessitating a rigorous examination of leaf venation, stem architecture, and floral morphology across the entire growing season.

4.1 Wild Garlic (*Allium ursinum*) and Its Lethal Lookalikes

Wild garlic, or Bärlauch, is a bulbous herbaceous perennial of the Amaryllidaceae family, highly prized for its culinary versatility and mild, alliaceous pungency. It thrives in nutrient-rich, moist, calcareous deciduous woodlands and alluvial forests⁴⁷. The plant emerges in early spring, producing elliptical to broadly lanceolate leaves, followed in May by terminal umbels of stark

white, star-shaped flowers with six tepals⁴⁷. As noted previously, the collection of this plant is strictly prohibited in Brandenburg due to its Red List status, though it may be observed or legally foraged in regions where populations are secure¹³.

The distinguishing botanical features of *A. ursinum* are critical for differentiation from toxic species:

1. **Leaf Venation:** The leaves possess strictly parallel venation⁴⁷.
2. **Petiole Architecture:** Each leaf features a distinct, discernible petiole (leaf stalk) that connects directly to the subterranean bulb, meaning the leaves do not clasp a central stem⁴⁷.
3. **Abaxial Surface:** The underside (abaxial surface) of the leaf is definitively matte, distinguishing it from glossier counterparts⁴⁷.
4. **Aroma:** The foliage emits a potent garlic scent due to sulfur-containing compounds when bruised⁴⁸.

4.1.1 The Danger of Olfactory Fatigue

While the garlic scent is a primary diagnostic tool, it poses a severe heuristic trap. After a forager crushes a few *A. ursinum* leaves, their fingers become heavily coated in the essential oils. Consequently, any subsequent plant handled by the forager will smell of garlic⁴⁷. This phenomenon of olfactory fatigue renders the smell test dangerously unreliable during prolonged foraging sessions, making visual morphology the only secure method of continuous identification⁴⁹.

4.1.2 Toxic Sympatric Species

Wild garlic frequently shares its precise ecological niche with three highly toxic species, emerging concurrently in the spring understory⁴⁷. A rigorous comparative analysis is essential.

Botanical Feature	<i>Allium ursinum</i> (Wild Garlic - Edible)	<i>Convallaria majalis</i> (Lily of the Valley - Toxic)	<i>Colchicum autumnale</i> (Autumn Crocus - Lethal)	<i>Arum maculatum</i> (Lords-and-Ladies - Toxic)
Toxic Principle	Safe/Edible ⁴⁸	Cardiac glycosides (convallatoxin); induces severe cardiac arrhythmias ⁴⁷	Colchicine; a severe mitotic inhibitor, highly lethal even in small doses ⁴⁷	Calcium oxalate raphides; causes severe mucosal irritation, burning, and swelling ⁴⁷
Leaf	Individual leaves, each	Leaves typically	Leaves emerge directly from	Leaves emerge individually but

Emergence	with its own distinct stalk growing from the bulb ⁴⁷	emerge in pairs from a rhizome, tightly clasping a central pseudo-stem ⁴⁷	the soil in a tight rosette formation, lacking individual stalks ⁴⁷	are distinctly sagittate (arrow-shaped) ⁴⁷
Venation Pattern	Parallel, with veins widely spaced ⁴⁷	Parallel, with a very densely packed, tight nervature ⁴⁷	Parallel, forming a relatively rigid, stiff structure ⁴⁷	Reticulate (net-like) and highly irregular ⁴⁷
Texture & Underside	Soft and pliable; distinctly matte on the abaxial (under) side ⁴⁷	Firm and leathery; glossy on both the adaxial and abaxial surfaces ⁴⁷	Fleshy, stiff, and almost rigid; glossy on both surfaces ⁴⁷	Firm; frequently exhibits dark maculations (spots) on the adaxial surface, though young leaves may lack spots ⁴⁷
Floral Structure	White, star-shaped umbels blooming concurrently with late foliage ⁴⁷	White, pendulous, bell-shaped flowers arranged on a raceme ⁴⁷	Blooms in autumn (pinkish-purple goblet shape), not present during spring leaf emergence ⁴⁷	A distinct, complex spadix enclosed by a pale green, hood-like spathe ⁵⁰

4.2 Ground Ivy (*Glechoma hederacea*)

Glechoma hederacea, commonly known as ground ivy or Gundermann, is a pervasive, creeping perennial of the Lamiaceae (mint) family⁵¹. It occupies a wide range of habitats across the Berlin-Brandenburg area, thriving equally in shaded woodland margins, urban lawns, and dense hedgerows⁵¹. Morphologically, the plant is characterized by trailing, square stems that root aggressively at the nodes (stolons) and opposite, reniform (kidney-shaped) to cordate leaves with distinctively crenate (scalloped) margins⁵¹. In the spring, it produces axillary whorls of zygomorphic, purplish-blue flowers marked with darker purple spots on the lower lip⁵¹. Phytochemically, ground ivy possesses an incredibly complex profile. It is rich in volatile oils (including camphene, pinene, and pulegone), sesquiterpenes (such as glechomafuran),

flavonoids (quercetin, luteolin, apigenin), and significant concentrations of pentacyclic triterpenic acids, notably ursolic acid and oleanolic acid⁵³. It also contains robust levels of phenolic acids, particularly rosmarinic acid, which imparts potent antioxidant and anti-inflammatory properties⁵³. Recent biotechnological research has even utilized the high concentration of bioactive reducing agents in *G. hederacea* extracts for the green synthesis of Au/ZnO/Ag nanoparticles, highlighting its complex chemical utility⁵⁴.

Ethnobotanically, the synergy of astringent tannins and aromatic volatile oils makes *G. hederacea* a classic European remedy for respiratory and digestive ailments⁵². It is historically indicated for chronic bronchial catarrh, acting as a mild expectorant, decongestant, and antikatarrhal agent⁵¹. Furthermore, it has long-standing traditional applications for treating tinnitus and middle-ear congestion⁵¹. The bitter principles and volatile oils also confer carminative properties, stimulating gastrointestinal secretions to alleviate dyspepsia and mild gastritis⁵². Due to the presence of pulegone—an essential oil component with potential hepatic toxicity and abortifacient properties in massive doses—internal consumption is best kept to moderate, therapeutic levels, and it is generally contraindicated during pregnancy⁵⁶.

4.3 Yarrow (*Achillea millefolium*)

Achillea millefolium, or Schafgarbe, is a ubiquitous rhizomatous perennial of the Asteraceae family. It is immediately recognizable by its highly dissected, bipinnatifid, feathery leaves (the source of the specific epithet *millefolium*, meaning "thousand leaves") and dense terminal corymbs of small, white to pinkish capitula (flower heads)⁵⁷.

Yarrow possesses a highly robust pharmacological profile, featuring essential oils (rich in proazulenes which convert to the anti-inflammatory compound chamazulene upon distillation), sesquiterpene lactones, flavonoids, and astringent tannins⁵⁹. The European Medicines Agency (EMA) Committee on Herbal Medicinal Products (HMPC) has established a Community herbal monograph recognizing its established traditional use for several primary indications:

1. **Gastrointestinal Distress:** Used as a bitter tonic for temporary loss of appetite and to relieve mild, spasmodic gastrointestinal complaints, including bloating and flatulence⁵⁷.
2. **Gynecological Spasms:** Efficacious as an antispasmodic in mitigating minor pelvic cramps associated with menstrual periods⁵⁹.
3. **Dermatological Applications:** Applied topically as an aqueous infusion for the treatment of small, superficial wounds, utilizing its powerful astringent and mild antiseptic properties to promote tissue coagulation⁵⁷.

The standard posology for internal use typically involves infusing 1.5 to 4 grams of the comminuted (finely cut) herb in 150–250 ml of boiling water, consumed up to four times daily⁵⁹. Individuals with known hypersensitivities to other members of the Asteraceae family should exercise caution, as allergic cross-reactivity is well documented⁵⁹.

4.4 Mugwort (*Artemisia vulgaris*)

Artemisia vulgaris (Beifuß) is a tall, robust perennial of the Asteraceae family, forming dense stands in ruderal sites, railway embankments, and waste grounds throughout the urban landscape of Berlin⁶⁰. It is identified by its deeply pinnatifid leaves, which are dark green and glabrous on the adaxial surface but densely tomentose (covered in silvery-white hairs) on the

abaxial surface⁶¹.

Mugwort is rich in volatile oils, particularly 1,8-cineole, camphor, and the bicyclic monoterpene beta-thujone, as well as sesquiterpene lactones such as vulgarin⁶². Historically, it has been employed in European herbalism as a powerful bitter tonic to stimulate gastric juices, bile flow, and aid in the digestion of heavy, lipid-rich meals⁶⁴. Furthermore, it has a long-standing ethnobotanical reputation as an emmenagogue, used to stimulate menstrual flow⁶².

However, the presence of thujone dictates caution; in high concentrations, thujone acts as a neurotoxin and an abortifacient, meaning that extended internal use, or any use during pregnancy, is strictly contraindicated⁶². Additionally, mugwort pollen is a highly potent aeroallergen and exhibits significant cross-reactivity with other pollens (notably sunflower pollen) and certain food allergens, necessitating caution for atopically sensitive individuals when handling the flowering tops⁶².

5. The Apiaceae Family: Navigating Edibility and Lethality

The Apiaceae (formerly Umbelliferae) family represents one of the greatest challenges for the botanical forager. This family contains some of the most esteemed wild edibles and culinary herbs (such as wild carrot, caraway, and fennel), alongside some of the most virulently toxic plants indigenous to the Northern Hemisphere⁶⁶. Foraging within this family demands advanced botanical literacy, as the morphological boundaries between edible and lethal species are notoriously narrow⁶⁶.

5.1 Safe and Edible Apiaceae

Ground Elder (*Aegopodium podagraria*): Known locally as Giersch, this plant is a highly successful, often aggressive rhizomatous perennial that thrives in shaded gardens, parks, and forest margins⁶⁸. It is recognizable by its ternate leaves (divided into three leaflets, which are themselves often lobed or divided) and a distinctively triangular petiole cross-section⁶⁷. Ecologically, *A. podagraria* is highly adapted to understory light fluctuations. Interestingly, the plant can accumulate significant levels of nitrates in its foliage, a factor influenced heavily by leaf age and ambient shade⁶⁸. Culturally, it is an excellent, highly nutritious spinach substitute, and its specific epithet reflects its historical medicinal use as a poultice to treat the painful joint inflammation of gout (*podagra*)⁶⁷.

Wild Angelica (*Angelica sylvestris* and *A. archangelica*): These are monumental, architectural biennial or perennial herbs found in damp meadows and riparian zones⁶⁷. They feature large, bipinnate leaves with heavily inflated, sheathing leaf bases that wrap dramatically around a stout, hollow, purplish or green stem⁷¹. The entire plant is highly aromatic, and its roots and candied stems hold historical significance in herbal medicine and traditional confectionery⁶⁷.

5.2 Lethal Apiaceae Lookalikes

The identification of any edible Apiaceae species must invariably involve aggressively ruling out the presence of three specific neurotoxic species, all of which share the compound umbel

structure and deeply dissected foliage typical of the family⁶⁶.

Botanical Feature	Conium maculatum (Poison Hemlock)	Cicuta virosa (Water Hemlock)	Aethusa cynapium (Fool's Parsley)
Toxic Principle	Piperidine alkaloids (primarily Coniine) ⁶⁶	Polyacetylenes (primarily Cicutoxin) ⁶⁶	Polyacetylenes and various alkaloids ⁶⁶
Mechanism of Toxicity	Acts on nicotinic acetylcholine receptors, leading to ascending muscular paralysis and eventual respiratory failure. Famously utilized for the execution of Socrates ⁶⁶ .	Acts as a potent noncompetitive antagonist at the GABA receptor in the central nervous system, causing violent convulsions, seizures, and rapid death ⁶⁶ .	Causes severe gastrointestinal distress, intense burning of the mouth and throat, and neurological impairment ⁶⁶ .
Stem Morphology	Glabrous (hairless), hollow, and prominently marked with distinctive purple maculations (blotches) on the lower half ⁶⁶ .	Stout, hollow, typically green but may be suffused with a uniform purple tint; critically, it lacks the distinct spots of <i>C. maculatum</i> ⁶⁶ .	Glabrous, finely grooved, often exhibiting a slight bluish bloom or reddish tint at the extreme base ⁶⁶ .
Root/Rhizome Structure	Produces a long, white, fleshy taproot ⁶⁶ .	Features a highly distinctive vertical rhizome that is internally divided into hollow chambers by transverse septa; exudes a yellow toxic sap when cut ⁶⁶ .	Possesses a thin, white taproot strongly resembling a small radish ⁶⁶ .

Aromatic Profile	Emits a highly unpleasant, mousy odor when the foliage is crushed ⁶⁶ .	Emits a parsnip or carrot-like odor, which is deceptively pleasant and invites accidental ingestion ⁶⁶ .	Emits a nauseating, somewhat garlic-like odor when crushed ⁶⁶ .
Key Involucral Bracts	Features short, lanceolate bracts situated at the base of the main umbel ⁶⁶ .	Lacks main involucral bracts entirely, but possesses multi-leaved involucels located under the secondary umbellets ⁶⁶ .	Possesses 3 to 4 highly distinctive, elongated bracteoles that point sharply downward, hanging visibly below each secondary umbellet ⁶⁶ .

6. Medicinal Roots: Pharmacognosy, Biochemistry, and Clinical Application

While it is imperative to reiterate that the unpermitted excavation of wild roots is ecologically destructive and legally restricted by the Handstraußregelung⁶, understanding the deep pharmacognosy of regional roots is vital for botanical research. For individuals cultivating these species or possessing appropriate land-use permissions, the harvest of these subterranean structures yields powerful phytomedicines.

6.1 Horseradish (*Armoracia rusticana*)

Horseradish is a perennial crop of the Brassicaceae family, possessing a massive, fleshy, cylindrical white taproot. The root system is extraordinarily extensive; if undisturbed in loose soil, the primary and lateral roots can penetrate to depths of three to four meters, with a lateral spread of one meter⁷⁵. Native to Eastern Europe and the Mediterranean, it has naturalized heavily in damp, arable lands, waste grounds, and riparian zones across Germany⁷⁵.

The primary bioactive constituents of the intact root are the glucosinolates sinigrin (2-propenyl glucosinolate) and gluconasturtiin, which serve as the plant's primary defensive mechanism against herbivory⁷⁵. The intact root is virtually odorless. However, upon cellular disruption (via grating, cutting, or mastication), the endogenous enzyme myrosinase rapidly hydrolyzes these glucosinolates into highly volatile compounds: allyl isothiocyanate (AITC) from sinigrin, and 2-phenylethyl isothiocyanate (PEITC) from gluconasturtiin⁷⁵. These isothiocyanates are directly responsible for the lachrymatory (tear-inducing), intensely pungent characteristics of the root⁷⁵.

Pharmacologically, AITC and PEITC exhibit profound, broad-spectrum antibacterial and antifungal properties⁷⁶. The German Commission E has officially approved the internal use of A.

rusticana root (recommending an average daily dose of 20g of fresh root) as a supportive therapy for urinary tract infections and catarrhs of the respiratory tract⁷⁸. The volatile isothiocyanates are absorbed systemically and excreted partially through the lungs and kidneys, exerting direct, localized antimicrobial effects on those specific mucous membranes⁷⁶. Externally, galenical preparations containing a maximum of 2% mustard oil are utilized as a rubefacient to induce local hyperemia (increased blood flow), providing relief for minor muscle aches and arthritic pain⁷⁶. Due to the intense stimulatory effect on gastric mucosa, it is strictly contraindicated for individuals suffering from stomach or intestinal ulcers, or acute renal disorders⁷⁹.

6.2 Chicory (*Cichorium intybus*) and Dandelion (*Taraxacum officinale*)

Common chicory and dandelion share a similar biochemical profile, both being prominent, deep-rooted members of the Asteraceae family²¹. Chicory is an erect, branching perennial, easily identified in late summer by its striking azure-blue ligulate flowers that bloom on rigid, wiry stems along roadsides and dry embankments²¹. Dandelion is a ubiquitous rosette-forming perennial characterized by deeply lobed leaves and bright yellow capitula²¹.

The thick taproots of both species serve as primary commercial and botanical sources of inulin, a naturally occurring fructan-type polysaccharide that functions as the plant's primary carbohydrate energy reserve⁸¹. Inulin is stored as linear polymers of fructose linked by β -(2 $\rightarrow$ 1) glycosidic bonds, typically terminating with a glucose residue⁸¹.

The biosynthesis of inulin is enzymatically driven: sucrose:sucrose 1-fructosyltransferase (1-SST) initiates chain formation from sucrose, while fructan:fructan 1-fructosyltransferase (1-FFT) elongates the polymer⁸².

The structural and physiological properties of chicory inulin are dictated by its degree of polymerization (DP), which typically ranges between 7 and 15 fructose units⁸¹. The DP is highly sensitive to the seasonal harvest cycle. Inulin accumulation and chain length peak in early autumn as the plant maximizes its reserves for dormancy⁸². As winter approaches, the plant activates fructan 1-exohydrolase (1-FEH), a hydrolytic enzyme that degrades the long-chain fructans into shorter fructooligosaccharides (FOS)⁸². This degradation increases the osmotic concentration of the cell sap, thereby conferring vital frost resistance to the root tissue⁸².

Consequently, for the medicinal or dietary extraction of high-DP inulin—valued heavily in the food industry as a prebiotic dietary fiber that modulates the gut microbiome, acts as a fat replacer, and modifies texture—roots must be harvested in the late summer or early autumn⁸¹.

Beyond inulin, the roots contain potent sesquiterpene lactones, specifically lactucin and lactucopicrin⁸⁶. These bitter principles strongly stimulate gastric and biliary secretions. The EMA monograph and Commission E acknowledge both chicory and dandelion roots for their traditional use in optimizing digestion, stimulating appetite, and treating dyspepsia⁸⁰.

Furthermore, lactucin and lactucopicrin exhibit documented mild sedative and analgesic properties⁸⁷. Dandelion root specifically possesses a high concentration of potassium salts, contributing to a powerful diuretic effect; uniquely, it replaces the potassium lost through diuresis, avoiding the hypokalemia often associated with synthetic loop diuretics⁸⁰. Both roots are contraindicated in cases of biliary obstruction or severe gallbladder disease⁸⁰.

6.3 Stinging Nettle Root (*Urtica dioica*)

While the aerial parts of *Urtica dioica* are renowned for their nutritional density and topical counter-irritant properties via formic acid and histamine, the rhizome and root system (*Urticae radix*) hold a highly specialized and heavily researched pharmacological application⁹².

Nettle root is utilized extensively in the management of Benign Prostatic Hyperplasia (BPH) and associated lower urinary tract symptoms (LUTS)⁹³. The German Commission E, alongside numerous double-blind, placebo-controlled clinical trials, endorses its use to alleviate urinary difficulties (such as nocturia, decreased flow rate, and elevated residual urine volume) in stages I and II of BPH⁹².

The clinical efficacy of the root extract is attributed to a synergistic interplay of several phytochemical classes acting on the endocrine and proliferative pathways of the prostate:

1. **Lignans and SHBG Binding:** Specific lignans in the root, particularly (-)-3,4-divanillyltetrahydrofuran, bind with high affinity to Sex Hormone-Binding Globulin (SHBG)⁹³. By occupying these binding sites, the lignans inhibit SHBG from binding to the prostate membrane receptors, thereby disrupting a signal cascade that ordinarily promotes abnormal prostatic cellular proliferation⁹³.
2. **5-Alpha-Reductase Inhibition:** Botanical fractions within the root act to inhibit 5-alpha-reductase, the critical enzyme responsible for converting testosterone into the highly potent dihydrotestosterone (DHT)⁹⁶. Elevated levels of DHT are a primary hormonal driver of prostate glandular enlargement⁹⁶.
3. **Aromatase Inhibition:** Phytochemicals such as 9-hydroxy-10-trans-12-cis-octadecadienoic acid (HOA) inhibit aromatase, the enzyme that converts testosterone into estrogens⁹³. Age-related increases in estrogen signaling within the prostate are known to exacerbate cellular hyperplasia, making aromatase inhibition a key therapeutic target⁹³.
4. ***Urtica dioica* Agglutinin (UDA):** This heat- and acid-resistant lectin exhibits significant immunomodulatory effects and actively antagonizes the epidermal growth factor receptor, further preventing abnormal tissue growth and reducing localized inflammation⁹³.

6.4 Comfrey (*Symphytum officinale*) and the Pyrrolizidine Alkaloid Crisis

Comfrey (*Symphytum officinale*, Boraginaceae) possesses a thick, mucilaginous taproot that has been utilized for millennia to heal bone fractures, sprains, and severe contusions (hence the traditional common name "knitbone")⁹⁸. The primary beneficial compound in the root is allantoin, a highly potent stimulant of cell proliferation and connective tissue regeneration⁹⁹.

Despite its historical reverence, the internal consumption of comfrey root is now strictly obsolete and highly dangerous due to the presence of Pyrrolizidine Alkaloids (PAs)¹⁰⁰. PAs are secondary metabolites synthesized by the plant as a defense mechanism against herbivores¹⁰⁰. While the native PAs in the plant are relatively non-toxic, upon absorption into the human gastrointestinal tract, they are transported to the liver¹⁰⁰. Here, the hepatic cytochrome P450

enzymes metabolically activate the PAs into highly reactive dehydropyrrolizidine (pyrrolic) derivatives¹⁰⁰. These reactive electrophiles aggressively bind to cellular proteins and DNA in the hepatic endothelia, causing severe hepatotoxicity, specifically hepatic veno-occlusive disease (VOD), mutagenicity, and eventual liver failure¹⁰⁰.

Consequently, major regulatory bodies, including the European Medicines Agency (EMA) and the German Federal Institute for Drugs and Medical Devices (BfArM), have issued stringent public statements and regulations regarding PAs¹⁰¹. BfArM has established a maximum upper limit of 1.0 µg of PA relative to the daily dose for herbal medicinal products¹⁰³. The use of *S. officinale* must therefore be strictly limited to **external, topical applications on unbroken skin** to prevent the systemic absorption of the alkaloids into the bloodstream⁹⁹. Commercial comfrey ointments are heavily regulated, often utilizing PA-depleted extracts, and their application is generally restricted to a maximum of 4 to 6 weeks per year to mitigate any risk of chronic accumulation⁹⁹.

Works cited

1. Handstraußregelung: Pflanzen und Holz im Wald sammeln - Deurag, <https://www.deurag.de/blog/handstrausregelung/>
2. Vollzugshinweise zum Artenschutzrecht, https://www.la-na.de/media/vollzugshinweise_stand_19xi2010-2_1518592222.pdf
3. Wald-Knigge - Forstwirtschaft in Deutschland, <https://www.forstwirtschaft-in-deutschland.de/waelder-entdecken/wald-knigge/>
4. Darf ich Blumen pflücken? - Artenschutz - BMUKN, <https://www.bundesumweltministerium.de/themen/artenschutz/artenschutz-durch-den-buerger/darf-ich-blumen-pfluecken>
5. Darf man Wildkräuter sammeln? (Deutschland) - Waldsamkeit, <https://waldsamkeit.de/darf-man-wildkraeuter-sammeln/>
6. Yamadori und das Gesetz: Was ist beim Bäumessammeln erlaubt?, <https://bonsai-treff.com/yamadori-gesetz-baeume-sammeln-ohne-bussgeld/>
7. Handstraußregel - wie viel darf man in der Natur ernten, <https://www.samenhaus.de/gartenblog/handstrausregel-wie-viel-darf-man-in-der-natur-ernten>
8. Beeren sammeln in Brandenburg: Sommer-Mundraub | LICHTUNG, <https://lichtung.berlin/magazin/beeren-sammeln-brandenburg/>
9. Integratives Monitoring der deutschen Großschutzgebiete, https://nationale-naturlandschaften.de/wp-content/blogs.dir/62/files/2023_Skript_644_Integratives-Monitoring-der-deutschen-Grossschutzgebiete-Nationalparks-und-Biosphaerenreservate.pdf
10. Berliner NaturschutzGes. - NatSchGBln - Verband Wohneigentum, <https://www.verband-wohneigentum.de/berlin-brandenburg/downloadmime/18925/NatschGBln.pdf>
11. Heilpflanzen am Wegesrand - Deutscher Alpenverein, https://www.alpenverein.de/artikel/heilpflanzen-am-wegesrand_52c84ba3-09c8-4dc6-91b2-ed725ef7527d

12. BArtSchV - Verordnung zum Schutz wild lebender Tier,
https://www.gesetze-im-internet.de/bartschv_2005/BJNR025810005.html
13. Steht Bärlauch unter Naturschutz? | mundraub.org,
<https://mundraub.org/blog/steht-baerlauch-unter-naturschutz>
14. Befindet sich Bärlauch auf der Roten Liste? Folgen fürs Sammeln,
<https://www.gartenjournal.net/baerlauch-rote-liste>
15. Bärlauch Erkennen - Waldsamkeit, <https://waldsamkeit.de/baerlauch-erkennen/>
16. Rote Listen | Startseite - LfU Brandenburg,
<https://lfu.brandenburg.de/lfu/de/aufgaben/natur/artenschutz/rote-listen/>
17. Brandenburgisches Ausführungsgesetz zum ... - BRAVORS,
<https://bravors.brandenburg.de/gesetze/bbgngatschag>
18. Was tun bei Baumfällungen? - NABU Berlin,
<https://berlin.nabu.de/stadt-und-natur/naturschutz-berlin/baumschutz/26003.html>
19. The Use of Biosensors for Biomonitoring Environmental Metal,
<https://www.intechopen.com/chapters/66235>
20. Heavy Metals Bioindication Potential of the Common Weeds ... - MDPI,
<https://www.mdpi.com/1420-3049/24/15/2813>
21. (PDF) Bioaccumulation of Heavy Metals by Herbaceous Species,
https://www.researchgate.net/publication/359887983_Bioaccumulation_of_Heavy_Metals_by_Herbaceous_Species_Grown_in_Urban_and_Rural_Sites
22. Accumulation of heavy metals from soil in medicinal plants,
https://www.researchgate.net/publication/320205104_Accumulation_of_heavy_metals_from_soil_in_medicinal_plants
23. Assessment of Heavy Metal Accumulation in Soils and Dominant,
<https://www.mdpi.com/2223-7747/15/6/983>
24. Medicinal Plant Growth in Heavy Metals Contaminated Soils - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9504071/>
25. Potential Ecological and Human Health Risks of Heavy Metals in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7914753/>
26. Fuchsbandwurm – was man wissen sollte ... - Nimrods,
<https://www.nimrods.de/Fuchsbandwurm.pdf>
27. RKI-Ratgeber - Echinokokkose,
https://www.rki.de/DE/Aktuelles/Publikationen/RKI-Ratgeber/Ratgeber/Ratgeber_Echinokokkose.html
28. Alveoläre Echinokokkose - Wikipedia,
https://de.wikipedia.org/wiki/Alveol%C3%A4re_Echinokokkose
29. Fuchsbandwurm breitet sich aus: Ist er gefährlich?,
<https://www.apotheken-umschau.de/krankheiten-symptome/infektionskrankheiten/was-sie-ueber-den-fuchsbandwurm-wissen-sollten-966709.html>
30. Fuchsbandwurm | Symptome Menschen | Risikogebiete | Infektion,
<https://www.meine-gesundheit.de/krankheit/krankheiten/fuchsbandwurm>
31. Die Fuchsbandwurm-Furcht - Ärzte Zeitung,
<https://www.aerztezeitung.de/Medizin/Die-Fuchsbandwurm-Furcht-228672.html>
32. A review of garlic mustard (Alliaria petiolata, Brassicaceae) as an,

- <https://bioone.org/journals/the-journal-of-the-torrey-botanical-society/volume-143/issue-4/TORREY-D-15-00059/A-review-of-garlic-mustard-Alliaria-petiolata-Brassicaceae-as-an/10.3159/TORREY-D-15-00059.full>
33. A review of garlic mustard (*Alliaria petiolata* , Brassicaceae) as an,
https://www.researchgate.net/publication/305789047_A_review_of_garlic_mustard_Alliaria_petiolata_Brassicaceae_as_an_allelopathic_plant
 34. Garlic mustard (raw, organic) | Foundation G+E - diet-health.info,
<https://www.diet-health.info/en/recipes/ingredients/in/wf8893-garlic-mustard>
 35. Evolution of specifier proteins in glucosinolate-containing plants - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3482593/>
 36. Diversified glucosinolate metabolism: biosynthesis of ... - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4628127/>
 37. Free and Bound Volatiles of Garlic Mustard (*Alliaria petiolata*),
<https://hrcak.srce.hr/file/49441>
 38. Effects of Glucosinolate-Derived Isothiocyanates on Fungi - MDPI,
<https://www.mdpi.com/2309-608X/7/7/539>
 39. *Alliaria petiolata* (M. Bieb.) Cavara & Grande – Monograph,
<https://www.alleanzaverde.com/blog/alliaria-petiolata-m-bieb-cavara-grande-monograph/>
 40. (PDF) Mustard Seeds and Leaves: Exploring Nutritional Benefits,
https://www.researchgate.net/publication/393046037_Mustard_Seeds_and_Leaves_Exploring_Nutritional_Benefits_and_Therapeutic_Applications
 41. Studies regarding the myrosinase enzymatic activity from black,
https://www.researchgate.net/publication/269096428_Studies_regarding_the_myrosinase_enzymatic_activity_from_black_mustard_seeds_Brassica_nigra
 42. Pulsatilla | NABU,
https://www.nabu.de/imperia/md/content/nabude/botanik/pulsatilla/pulsatilla_3.pdf
 43. Monitoringprogramm für das UNESCO-Biosphärenreservat,
https://www.elbetal-mv.de/fileadmin/elbetal/downloads/01_Wissen_und_Versteuern/Forschung/2021_Monitoringprogramm_Biosphaerenreservat_Flusslandschaft_Elbe_MV.pdf
 44. Ökonomische Folgen der Ausbreitung von Neobiota,
<https://www.umweltbundesamt.de/system/files/medien/publikation/long/2433.pdf>
 45. biodiversitaet.pdf,
https://tlubn.thueringen.de/fileadmin/000_TLUBN/Service/Publikationen/Sonderpublikationen/inhalt/biodiversitaet.pdf
 46. Berichte des Botanischen Vereins zu Hamburg,
https://www.botanischerverein.de/wp-content/uploads/Ber_Bot_Verein_23.pdf
 47. So kannst Du Bärlauch sicher von Maiglöckchen, Herbst-Zeitlose,
<https://floraincognita.de/so-kannst-du-baerlauch-sicher-von-maigloeckchen-herbst-zeitlose-aronstab-und-salomonsiegel-unterscheiden/>
 48. Bärlauch richtig ernten und lagern: Bewahren Sie den Geschmack,
<https://samen.de/blog/baerlauch-richtig-ernten-und-lagern-so-bewahren-sie-den-vollen-geschmack.html>

49. Bärlauch und der Unterschied zu Aronstab, Maiglöckchen und,
<https://natur-tagebuch.ch/vom-baerlauch-und-den-unterschieden-zu-aronstab-maiglloeckchen-und-herbstzeitlosen/>
50. Aronstab-ähnliche Pflanzen: Wie Sie Verwechslungen vermeiden,
<https://www.gartenjournal.net/aronstab-aehnliche-pflanzen>
51. Glechoma hederacea, <https://eflora.neocities.org/Glechoma%20hederacea>
52. Ground Ivy (Glechoma hederacea): Benefits, Safety, Uses,
<https://www.herbalreality.com/herb/ground-ivy/>
53. Antiinflammatory Activity of Constituents from Glechoma hederacea,
https://www.researchgate.net/publication/51086572_ChemInform_Abstract_Anti_inflammatory_Activity_of_Constituents_from_Glechoma_hederacea_var_longituba
54. Facile synthesis of Au/ZnO/Ag nanoparticles using Glechoma ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7940322/>
55. A review of plant characteristics, phytochemistry and bioactivities of,
https://www.researchgate.net/publication/348571610_A_review_of_plant_characteristics_phytochemistry_and_bioactivities_of_the_genus_Glechoma
56. Ground Ivy, <https://studfile.net/preview/1903187/page:48/>
57. Millefolii herba - herbal medicinal product,
<https://www.ema.europa.eu/en/medicines/herbal/millefolii-herba>
58. Millefolii flos - herbal medicinal product,
<https://www.ema.europa.eu/en/medicines/herbal/millefolii-flos>
59. European Union herbal monograph on Achillea millefolium L., herba,
<https://fitoterapiabrasil.com.br/file/4014/download?token=DPaBn3Y>
60. Eatweeds Photo Guide | PDF | Perennial Plant | Leaf - Scribd,
<https://www.scribd.com/document/546218920/Eatweeds-Photo-Guide>
61. (PDF) Towards a Better Understanding of Artemisia vulgaris: Botany,
https://www.researchgate.net/publication/324006830_Towards_a_Better_Understanding_of_Artemisia_vulgaris_Botany_Phytochemistry_Pharmacological_and_Biotechnological_potential
62. Pelin (Artemisia absinthium L.) - Biljni preparati,
<https://www.biljni-preparati.com/preparati/pelin-artemisia-absinthium-l-2/>
63. MATERIA MEDICA BOTANICA - Mountain.Ash,
<https://www.drashleyromanchuk.com/materia-medica>
64. PROSEA - Plant Resources of South East Asia,
<https://prosea.prota4u.org/view.aspx?id=165>
65. Mugwort. The scientific evidence for health benefits - Wise Mind Herbs,
<https://wisemindh herbs.com/herbalhealth/sciencereview/mugwort.htm>
66. Gefleckter Schierling - Wikipedia,
https://de.wikipedia.org/wiki/Gefleckter_Schierling
67. Doldenblütler erkennen - Unterscheidungsmerkmale - NAOWA,
<https://naowa.de/doldenbluetler-erkennen/>
68. Environmental Accounting - DOKUMEN.PUB,
<https://dokumen.pub/environmental-accounting.html>
69. Archiv der Kategorie: Energie und Umwelt - Wilfried Probst,

- <https://www.wilfried-probst.de/category/energie-und-umwelt/>
70. Tag: Victor Resco de Dios - plant stomata encyclopedia,
<https://plantstomata.wordpress.com/tag/victor-resco-de-dios/>
71. Arznei-Engelwurz (Angelika) - Kostbare Natur,
<https://www.kostbarenatur.net/anwendungen-und-inhaltsstoffe/arznei-engelwurz-angelika/>
72. Liste der Doldenblütler (Apiaceae) Frankreichs - Monde Végétal,
<https://monde-vegetal.fr/de/blogs/blog/apiacees-de-france>
73. G. J. Spiridonova Das Blumen-ABC oder deutsch-russisches,
<http://elibrary.udsu.ru/xmlui/bitstream/handle/123456789/13419/2015215.pdf?sequence=1>
74. Apiaceae, Doldenblütler mit weißen und weißlichen Blüten,
https://botanik-bochum.de/pflanzenbilder_familien_apiaceae_weiss.htm
75. Back to the Roots—An Overview of the Chemical Composition and,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8197530/>
76. Armoracia rusticana Horseradish, Red Cole PFAF Plant Database,
<https://pfaf.org/user/Plant.aspx?LatinName=Armoracia%20rusticana>
77. Back to the Roots—An Overview of the Chemical Composition and,
<https://www.mdpi.com/1420-3049/26/11/3155>
78. Medicine Update:Horseradish - American Botanical Council,
<https://www.herbalgram.org/resources/herbalegram/volumes/volume-15/number-5-may/food-as-medicine-update-horseradish-armoracia-rusticana-brassicaceae/medicine-updatehorseradish/>
79. Horseradish - American Botanical Council,
<https://www.herbalgram.org/resources/commission-e-monographs/monograph-approved-herbs/horseradish/>
80. DANDELION - TARAXACUM OFFICINALE,
<https://webprod.hc-sc.gc.ca/nhp/nd-bdipns/atReq?atid=dandelion.pissenlit2&lang=eng>
81. Techno-Functional Properties and Applications of Inulin in Food,
<https://www.mdpi.com/2310-2861/11/10/829>
82. Inulin chain length modification using a transgenic approach ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6068068/>
83. 1-FFT amino acids involved in high DP inulin accumulation in,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2015.00616/full>
84. Inulin: functional ingredients in food industry, extraction, and health,
<https://www.explorationpub.com/Journals/eff/Article/1010188>
85. agronomy - IRIS - Unict,
<https://www.iris.unict.it/retrieve/dfe4d22e-dc8b-bb0a-e053-d805fe0a78d9/Inulin%20agronomy-12-00290-v2.pdf>
86. Bitter Foods are Sometimes Better - Science and Education Publishing,
<https://pubs.sciepub.com/jpm/7/1/1/index.html>
87. Lactucopicrin - Wikipedia, <https://en.wikipedia.org/wiki/Lactucopicrin>
88. Dandelion root with herb - American Botanical Council,

- <https://www.herbalgram.org/resources/commission-e-monographs/monograph-approved-herbs/dandelion-root-with-herb/>
89. Cichorium intybus: Traditional Uses, Phytochemistry, Pharmacology, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3860133/>
 90. Large Chicory aphid (Uroleucon cichorii (Koch ... - Semantic Scholar, <https://pdfs.semanticscholar.org/7658/67d0584e650bc5102ff938c73e6b065c93c6.pdf>
 91. Kasni (Cichorium intybus): A Unani Hepatoprotective Drug, <https://jddtonline.info/index.php/jddt/article/download/4162/3226>
 92. Stinging Nettle root - American Botanical Council, <https://www.herbalgram.org/resources/expanded-commission-e-monographs/stinging-nettle-root/>
 93. Urtica urens (Nettle) Monograph - Alternative Medicine Review, <https://altmedrev.com/wp-content/uploads/2019/02/v12-3-280.pdf>
 94. A comprehensive review on the stinging nettle effect and efficacy, <https://pubmed.ncbi.nlm.nih.gov/17509841/>
 95. A comprehensive review on the stinging nettle effect and efficacy, https://www.researchgate.net/publication/6322337_A_comprehensive_review_on_the_stinging_nettle_effect_and_efficacy_profiles_Part_II_Urticae_radix
 96. Stinging Nettle for Prostate Health - Supplements in Review, <https://supplementsinreview.com/prostate/stinging-nettle-prostate/>
 97. The Effects of Urtica Dioica L. Root Extract on Rat Testis, https://anatomyjournal.ir/browse.php?a_id=169&sid=1&slc_lang=en&html=1
 98. Phytopharmacy: An Evidence-Based Guide to Herbal Medicinal, <https://dokumen.pub/phytopharmacy-an-evidence-based-guide-to-herbal-medicinal-products-9781118543566-1118543564.html>
 99. Comfrey root: From tradition to modern clinical trials - ResearchGate, https://www.researchgate.net/publication/233887224_Comfrey_root_From_tradition_to_modern_clinical_trials
 100. Herbal hepatotoxicity in traditional and modern medicine - Frontiers, <https://www.frontiersin.org/journals/pharmacology/articles/10.3389/fphar.2015.00072/pdf>
 101. public-statement-use-herbal-medicinal-products-containing-toxic, https://www.ema.europa.eu/en/documents/public-statement/public-statement-use-herbal-medicinal-products-containing-toxic-unsaturated-pyrrolizidine-alkaloids-pas-including-recommendations-regarding-contamination-herbal-medicinal-products-pyrrolizidine_en.pdf
 102. Public statement on the use of herbal medicinal products containing, https://www.ema.europa.eu/en/documents/public-statement/draft-public-statement-use-herbal-medicinal-products-containing-toxic-unsaturated-pyrrolizidine-alkaloids-pas-including-recommendations-regarding-contamination-herbal-medicinal-products-pyrrolizidine_en.pdf
 103. Detailanzeige - ECV, https://www.ecv.de/suse_item.php?suseld=Z|pil8506&susePattern=
 104. BULLETIN ZUR ARZNEIMITTELSICHERHEIT - Paul-Ehrlich-Institut,

https://www.pei.de/SharedDocs/Downloads/DE/newsroom/bulletin-arzneimittelsicherheit/2022/4-2022.pdf?__blob=publicationFile&v=4