

Analytical and Biological Methodologies for the Evaluation of Soil pH and Macronutrient Profiles: From Chemical Indicators to Plant Bio-Diagnostics

1. Introduction

The accurate characterization of soil pH and macronutrient profiles—specifically nitrogen (N), phosphorus (P), and potassium (K)—is a foundational requirement for sustainable agricultural management, ecosystem monitoring, and the optimization of the global circular bioeconomy¹. While contemporary agronomy relies heavily on sophisticated analytical instrumentation such as inductively coupled plasma mass spectrometry (ICP-MS) and high-performance liquid chromatography (HPLC), these technologies are often inaccessible in resource-limited, remote, or small-scale farming contexts³. Consequently, there is a critical need for robust, rudimentary chemical testing methods and observational bio-diagnostics that can accurately approximate soil fertility parameters in the field.

The evaluation of soil profiles without advanced laboratory equipment relies on two primary paradigms. The first involves rudimentary chemical indicators—colorimetric and turbidimetric reagents that react with soil extracts to produce visually quantifiable changes. The second paradigm leverages plant bio-indicators, wherein the physiological, metabolic, and phenotypic responses of flora serve as living sensors for edaphic conditions⁵. Plants possess highly conserved, genetically regulated mechanisms for sensing and responding to nutrient deficiencies and toxicities. By interpreting these biological signatures—ranging from specific patterns of leaf chlorosis and necrosis to the accumulation of distinct secondary metabolites like anthocyanins—practitioners can infer the chemical status of the rhizosphere with remarkable precision⁷.

This report provides an exhaustive analysis of both rudimentary chemical testing methodologies and plant-based bio-indicators. It details the thermodynamic and kinetic principles governing field-level soil extractions, the molecular mechanics of colorimetric macronutrient assays, the evaluation protocols for organic soil amendments, and the transcriptomic and biochemical pathways that govern plant responses to variations in soil pH, nitrogen, phosphorus, and potassium.

2. Physicochemical Principles of Soil Nutrient Extraction

Before rudimentary chemical indicators can be applied, nutrients must be liberated from the solid soil matrix into an aqueous solution. Soil extraction is a critical variable; the efficiency of nutrient release is dictated by the chemical nature of the extractant, the volumetric ratio of the

soil-to-solution, and the kinetic duration of the extraction process¹⁰.

Developing a rapid soil test kit typically begins with the simultaneous extraction of multiple targeted nutrients using a universal soil extractant, or the sequential extraction using specialized reagents designed for specific chemical pools¹⁰. The objective is to mimic the natural exudates of plant roots, gently displacing exchangeable cations and dissolving precipitated mineral complexes without initiating the total destruction of the primary mineral lattice¹⁰.

Extractant Formulations and Buffering Dynamics

Several chemical formulations have been developed to target the "plant-available" fraction of soil nutrients. The Morgan extractant, introduced in 1941, remains one of the most viable

universal extractants for rudimentary testing¹². It utilizes a highly buffered solution of 0.73 M sodium acetate and 0.52 M acetic acid, maintained at a stable pH of 4.8¹⁴. This weak acid buffer effectively displaces exchangeable potassium, calcium, and magnesium, although it typically recovers a narrower range of phosphorus compared to stronger acid extractants¹⁰. For targeted phosphorus extraction, particularly in acidic to neutral soils, the Bray-1 extractant is frequently employed. Composed of 0.025 M hydrochloric acid and 0.03 M ammonium fluoride (NH_4F), the Bray-1 solution relies on the fluoride ion to form highly stable, soluble complexes with trivalent aluminum (Al^{3+}) and iron (Fe^{3+}) cations¹⁰. By sequestering these metals, the extractant forces the dissolution of previously insoluble aluminum and iron phosphates, thereby liberating orthophosphate into the solution. However, the Bray-1 method is fundamentally incompatible with highly calcareous soils, as the abundance of calcium carbonate rapidly neutralizes the dilute hydrochloric acid and precipitates the fluoride as insoluble calcium fluoride¹⁰.

To address highly alkaline and calcareous soils, the Olsen extraction method utilizes 0.5 M sodium bicarbonate ($NaHCO_3$) buffered to a pH of 8.5¹⁰. In this elevated pH environment, the high concentration of carbonate and bicarbonate anions promotes the precipitation of calcium carbonate, which thermodynamically drives the dissolution of calcium-bound phosphates, releasing bioavailable phosphorus into the aqueous extract¹⁰. Alternatively, simple distilled water or rainwater extractions are utilized in the most rudimentary testing kits, though these only capture the immediately soluble, highly mobile fractions, such as nitrates and unbound salts¹⁹.

Beyond the chemical composition of the extractant, the volumetric ratio of soil to extractant—typically ranging from 1:1 to 1:5—profoundly influences the resulting analytical concentration. Soils possess inherent buffering capacities, defined as their ability to resist changes in pH and ionic strength upon the addition of acidic or basic solutions¹¹. Soils with high cation exchange capacities (CEC), such as those rich in humified organic matter and smectite clays, require wider extraction ratios (e.g., 1:5) or extended mechanical shaking times to

overcome this buffering effect and achieve true chemical equilibrium in the extract¹¹.

3. Rudimentary Chemical Indicators for Soil pH and Alkalinity


Soil pH acts as the master variable in agricultural chemistry, governing the solubility, speciation, and bioavailability of virtually all essential nutrients². For instance, at highly acidic pH levels, aluminum and manganese become soluble to the point of acute phytotoxicity, while at alkaline pH levels, phosphorus precipitates as insoluble calcium phosphate, and critical micronutrients like zinc and iron become unavailable to the plant root².

The Effervescence Method: Acetic Acid and Sodium Bicarbonate

A highly accessible, binary diagnostic test for extreme soil pH relies on fundamental acid-base neutralization reactions utilizing readily available household reagents: white vinegar (dilute


acetic acid, CH_3COOH) and baking soda (sodium bicarbonate, $NaHCO_3$)²⁴. This mechanical fizz test rapidly triages soils into highly acidic, highly alkaline, or near-neutral categories without the need for electronic instrumentation or calibrated color charts²⁴.

To test for severe alkalinity, an aqueous soil slurry is treated with acetic acid. If the soil contains high levels of free carbonates (indicating a pH generally above 7.5), a vigorous effervescence occurs due to the generation of carbon dioxide gas, governed by the following neutralization reaction²⁴:



Conversely, to test for severe acidity, the soil is mixed with distilled water to form a paste, and solid sodium bicarbonate is added. Distilled water is strictly required for this step, as tap water possesses its own inherent pH and buffering capacity that can generate false negatives²⁴. In

soils with a pH significantly below 6.0, the abundance of free protons (H^+) in the soil solution reacts with the bicarbonate anion to release carbon dioxide²⁴:



While predominantly utilized as a visual qualitative test, the kinetics of this reaction can be monitored in a more structured laboratory setting by gravimetrically tracking the mass loss of the reaction vessel as carbon dioxide gas escapes into the atmosphere, providing a rough quantitative estimate of free protons or carbonates²⁸.

Anthocyanin-Based Colorimetric Indicators

For a more granular and continuous assessment of soil pH, natural halochromic compounds can be extracted from specific plant tissues, most notably the red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*). Red cabbage contains massive concentrations of water-soluble pigments known as anthocyanins, predominantly cyanidin and its glucoside derivatives²⁹.

Anthocyanins are flavonoid phytochemicals whose molecular structures—and consequently their light-absorption spectra—are exceptionally sensitive to the hydrogen ion concentration of their surrounding aqueous environment³⁰. By subjecting macerated red cabbage to

pressurized hot water extraction or simple boiling, these pigments are easily liberated to form a universal natural pH indicator solution¹⁹.

The chemical mechanics of this color transition depend on the sequential deprotonation and structural rearrangement of the anthocyanin molecule²². In highly acidic environments (pH 1–3), the pigment exists primarily as the fully protonated flavylium cation. In this structural state, the molecule absorbs light in the blue-green spectrum, transmitting a vibrant red to pink color to the observer²². As the pH rises into the weakly acidic to neutral range (pH 4–7), the loss of a proton yields the carbinol pseudobase and the quinoidal base, causing the solution to shift toward deep purple and violet hues³⁶. Upon entering alkaline conditions (pH 8–11), further deprotonation forms the anionic quinoidal base, leading to a distinct blue to blue-green color profile¹⁹. Finally, in highly alkaline environments (pH > 11), the molecule undergoes a structural ring-opening, forming a yellow chalcone structure, although at these extreme pH levels, the anthocyanin molecule often becomes chemically unstable and degrades rapidly³⁴.

When an aqueous soil extract is mixed with the cabbage indicator, the resulting color provides a reasonably accurate, semi-quantitative estimation of the soil pH¹⁹. While it lacks the exact precision of a calibrated glass-electrode pH meter, standardized color charts can be empirically developed by calibrating the visual readings against known buffer solutions, elevating this kitchen-chemistry reaction to a highly robust field technique³⁴.


4. Colorimetric and Turbidimetric Methodologies for Macronutrients

Once soil nutrients are successfully extracted into a liquid phase, rudimentary diagnostic kits utilize specific chemical complexation reactions that yield either a visible color change (colorimetry) or a quantifiable cloudiness (turbidimetry). These methods bypass the need for atomic emission spectroscopy, relying instead on fundamental principles of optical absorbance and light scattering.

Nitrogen: Nitrate Reduction and the Griess Reaction

Nitrogen is the most critical macronutrient driving vegetative growth. In aerobic agricultural soils, the dominant bioavailable form of nitrogen is the highly mobile nitrate anion (NO_3^-)³⁸. Direct colorimetric detection of nitrate is chemically challenging because the unreactive nitrate ion does not readily form colored coordination complexes. Therefore, field methods typically require a preliminary reduction step to convert nitrate into the highly reactive nitrite anion (NO_2^-)⁴⁰.

Traditionally, this reduction was achieved by passing the sample through a copper-plated cadmium column. However, due to the acute toxicity and carcinogenic nature of cadmium, modern field kits and green analytical protocols employ metallic zinc powder (Zn^0) as the reducing agent⁴⁰. This reduction occurs most efficiently in a buffered medium, typically an acetate/acetic acid solution formulated at a concentration of 0.75 mol/L and a pH of 6.0⁴⁰. Within this slightly acidic environment, the zinc acts as a powerful electron donor⁴⁰:




Following the reduction phase, the newly generated nitrite is quantified utilizing the classic Griess reaction. The sample is reacted with sulfanilamide under highly acidic conditions to form a transient, highly reactive diazonium salt. This intermediate is subsequently coupled with a chromogenic agent, specifically N-(1-naphthyl)-ethylenediamine (NED), to synthesize a highly stable, intensely colored pink-to-magenta azo dye⁴⁰. The intensity of the magenta color is directly proportional to the original nitrate concentration. Utilizing optimized zinc ratios (e.g., 300 mg of 20 MESH zinc), this method achieves an exceptional limit of detection (LoD) of 0.024 mg/L⁴⁰. The color intensity can be visually matched against a standardized printed gradient or measured using a portable, field-deployable spectrophotometer reading absorbance at approximately 540 nm⁴¹.

Phosphorus: The Molybdenum Blue Method

Phosphorus exists in the soil solution primarily as soluble orthophosphate anions ($H_2PO_4^-$ and HPO_4^{2-}). The quantification of these inorganic phosphates in soil extracts is universally achieved via the Molybdenum Blue colorimetric method⁴³.

When an acidic soil extract is combined with a reagent containing ammonium molybdate and potassium antimonyl tartrate, the orthophosphate reacts with the molybdate ions to form a complex heteropoly acid known as phosphomolybdic acid⁴³. The formation of this complex is highly dependent on a strictly maintained acidic pH:




In its fully oxidized, unreduced state, phosphomolybdic acid exhibits a pale yellow color that is difficult to quantify visually. However, upon the addition of a mild reducing agent—most commonly ascorbic acid—the complex is reduced to form "molybdenum blue," an intensely colored polyoxometalate compound⁴³. The inclusion of antimony (provided by the potassium antimonyl tartrate) acts as a critical sensitizing catalyst⁴⁷. While ascorbic acid alone reduces the complex slowly, the antimony dramatically accelerates the reduction kinetics and stabilizes the final blue color at ambient room temperature, eliminating the need for boiling water baths and making the test highly suitable for rapid field diagnostics⁴⁷. The darkness of the blue color corresponds directly to the concentration of plant-available phosphorus in the soil, measured optimally at an absorbance wavelength of 880 nm⁴³.

Potassium: Tetraphenylborate and Cobaltinitrite Turbidimetry


Unlike nitrogen and phosphorus, which readily form distinct chromophores, potassium (K^+) lacks easily accessible optical transitions in the visible spectrum. Therefore, rudimentary potassium testing cannot rely on colorimetry. Instead, it utilizes targeted precipitation reactions that induce measurable turbidity (cloudiness) in the soil extract⁴.

The Sodium Tetraphenylborate (NaTPB) Method: The most frequently deployed turbidimetric assay for potassium involves the addition of sodium tetraphenylborate (

$Na[B(C_6H_5)_4]$) to the soil extract. Potassium ions react stoichiometrically with the bulky, hydrophobic tetraphenylborate anion to form a sparsely soluble white colloidal precipitate known as potassium tetraphenylborate ($KTPB$)⁵⁰.




Because ammonium (NH_4^+) and certain heavy metals also form highly insoluble precipitates with this reagent, field kits must employ complex chemical masking agents to prevent false-positive turbidity readings⁵⁰. A typical advanced masking buffer contains precise concentrations of copper ions, tartaric acid, ethylenediaminetetraacetic acid (EDTA), and sodium hydroxide or formaldehyde⁵⁰. Formaldehyde, for instance, effectively sequesters ammonium via the following condensation reaction:



Once the interferences are masked, the resulting white turbidity can be visually matched against a standardized printed gradient, often assessed by measuring the visual obscuration of a black line drawn at the bottom of the test tube⁵².

The Sodium Cobaltinitrite Method: An alternative turbidimetric procedure utilizes sodium cobaltinitrite ($Na_3[Co(NO_2)_6]$). When introduced to a potassium-containing extract, a characteristic orange-yellow to brownish crystalline double salt, potassium sodium cobaltinitrite ($K_2Na[Co(NO_2)_6] \cdot H_2O$), rapidly precipitates out of solution⁵⁶.



This precipitate can be measured directly via turbidimetry⁴. Alternatively, the precipitate can be collected, washed, and dissolved. The nitrite groups contained within the dissolved complex can then be utilized to diazotize sulfanilic acid (mechanistically similar to the Griess reaction) to form an azo dye, indirectly quantifying the potassium through colorimetry⁵⁶. Despite the reagent's inherent instability in aqueous solutions—which typically decompose within two to three weeks due to the aquation of nitrite ligands unless stabilized with methanol—it provides a highly reliable metric for potassium⁵⁷. Remarkably, the cobaltinitrite method permits the quantification of exchangeable potassium in soils at concentrations as low as 0.005%, and functions entirely independently of interferences from alkaline earth sulfates or phosphates, which routinely compromise other methodologies⁵⁷.

5. Profiling Organic Soil Amendments: Mineralization, Maturation, and Phytotoxicity

The generation and application of organic fertilizers—encompassing composted humanure, poultry litter, and specialized botanical oil cakes—is central to closing the global nutrient loop¹. However, raw organic waste is highly putrescible and poses severe agronomic risks if applied directly to soil. Testing organic soil amendments requires profiling not just absolute NPK values,

but evaluating the kinetics of nutrient mineralization, the structural evolution of the carbon matrix, and the elimination of severe phytotoxins¹.

Nitrogen Mineralization Kinetics and Humification Spectroscopy

Nitrogen within raw organic waste is predominantly locked within highly polymerized organic forms, such as structural proteins, microbial cell walls, and nucleic acids, rendering it entirely unavailable for plant uptake¹. To evaluate the fertilizer quality of an organic amendment, agronomists must model its mineralization kinetics—the microbially mediated conversion of

organic nitrogen into bioavailable ammonium (NH_4^+) and nitrate (NO_3^-)³⁸.

The release of this residual organic nitrogen pool is a time-dependent process most accurately modeled using a first-order exponential decay equation:

$$N_{min}(t) = N_0(1 - e^{-kt})$$

where $N_{min}(t)$ is the cumulative amount of nitrogen mineralized at time t (days), N_0

represents the absolute size of the potentially mineralizable organic nitrogen pool, and k represents the first-order mineralization rate constant³⁹. Highly humified amendments like

vermicompost exhibit significantly slower, more stable mineralization rates (e.g., $k = 0.009$) compared to raw poultry manure, acting as premium slow-release fertilizers that prevent nitrate leaching and subsequent aquatic eutrophication².

The stabilization of the organic matrix, known as humification, involves the profound biochemical transformation of labile organic matter into highly stable, recalcitrant humic substances¹. This structural evolution cannot be fully evaluated through wet chemistry alone; it requires rigorous non-destructive spectroscopic testing. UV-Visible Spectroscopy is employed to evaluate the E4/E6 ratio (the absorbance at 465 nm versus 665 nm). A steadily declining E4/E6 ratio is the primary indicator of increasing molecular weight, higher degrees of aromatic condensation, and advanced maturation of the humic acids¹. Concurrently, Fourier-Transform Infrared (FTIR) Spectroscopy tracks the progressive destruction of aliphatic bands and the emergence of oxygenated functional groups (such as carboxyl and phenolic hydroxyl moieties) that are characteristic of mature, stable humus capable of high cation exchange¹.

Biological Indicators of Phytotoxicity: The Germination Index (GI)

Applying immature, unstabilized compost to agricultural soils induces severe phytotoxicity, characterized by the catastrophic inhibition of seed germination, severe root necrosis, and nutrient lock-up¹. This toxicity is chemically mediated by the presence of high concentrations of

un-ionized ammonia (NH_3), highly mobile heavy metals, and short-chain volatile fatty acids (VFAs)—particularly acetic, propionic, and butyric acids—generated in anaerobic micro-sites within the material⁴¹.

Instead of attempting to chemically quantify every potential phytotoxin, the safety and maturity of an organic amendment are empirically evaluated using a comprehensive plant bio-indicator: the Germination Index (GI) bioassay¹. This biological standard is most frequently performed

using garden cress (*Lepidium sativum*) or cucumber (*Cucumis sativus*) seeds due to their extreme sensitivity to aqueous compost extracts¹.

The GI integrates both the relative rate of seed germination and the vigor of radicle (root) elongation relative to a distilled water control, calculated using the formula:

$$GI = \left(\frac{\text{Germinated seeds}_{\text{treatment}}}{\text{Germinated seeds}_{\text{control}}} \right) \times \left(\frac{\text{Root length}_{\text{treatment}}}{\text{Root length}_{\text{control}}} \right) \times 100$$

As microbial succession progresses during composting, volatile organic acids are exhausted and biological nitrification successfully converts toxic ammonium into benign nitrate, causing phytotoxicity to steadily dissipate¹. An organic amendment extract yielding a GI > 60% indicates acceptable baseline maturity, while a GI > 80% signifies the complete disappearance of phytotoxins and confirms the absolute agronomic safety of the final product¹.

Pathogen and Contaminant Biomarkers

Organic amendments derived from municipal biosolids or animal manures harbor diverse arrays of enteric pathogens and emerging synthetic contaminants. The biological sanitization of these amendments relies on maintaining strict thermophilic regimes (e.g., maintaining internal pile temperatures above 55°C for extended periods)¹.

The ultimate biological indicator for confirming successful compost sanitization is the destruction of the thick-shelled ova of the human roundworm, *Ascaris lumbricoides*¹. *Ascaris* eggs possess extreme resistance to standard chemical disinfection and desiccation; thus, documenting a >99% reduction in viable *Ascaris* eggs guarantees the concurrent destruction of less resilient bacterial pathogens like *Salmonella* spp. and *Escherichia coli*¹.

A more insidious metric of contamination is the presence of antibiotic resistance genes (ARGs)¹. Animal manures are critical environmental reservoirs for ARGs propagated via mobile genetic elements like Class 1 integrons (*int1*)¹. While the thermophilic phase physically denatures resistant plasmids, advanced molecular testing frequently reveals an "ARG rebound" phenomenon during the cooling phase, driven by the rapid proliferation of surviving multi-drug resistant bacteria co-selected by the presence of residual heavy metals¹. The integration of advanced additives like biochar is actively tested to suppress this rebound by physically adsorbing heavy metals and disrupting horizontal gene transfer mechanisms¹.

6. Plant Bio-Indicators: Epigenetic, Metabolic, and Phenotypic Signatures

While chemical extractions and spectroscopic indices provide a highly accurate snapshot of soil reservoirs and amendment stability, the ultimate, integrated measure of nutrient bioavailability is the plant itself. Plants exhibit profound physiological, transcriptomic, and morphological plasticity in response to soil nutrient deficiencies, transforming them into highly calibrated, living environmental sensors.

Nitrogen Deficiency: Rubisco Degradation and Vacuolar Autophagy

Nitrogen is a primary constituent of amino acids, nucleic acids, and the porphyrin rings of

chlorophyll. In C_3 plants, over 75-80% of total leaf nitrogen is sequestered within the chloroplasts of mesophyll cells, predominantly as the enzyme ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) and the light-harvesting complex⁶⁷. Consequently, the most universal bio-indicator of nitrogen deficiency is progressive leaf yellowing (chlorosis), originating in older, lower leaves as the plant actively cannibalizes its own photosynthetic machinery to remobilize nitrogen to younger, actively growing tissues (sink organs)⁶⁹. The mechanism of this chlorosis represents a highly orchestrated form of programmed cell death and targeted proteolysis. Under nitrogen starvation, the breakdown of Rubisco and other stromal proteins is aggressively accelerated⁶⁷. The precise cellular mechanisms driving this degradation involve complex intracellular trafficking pathways that export chloroplast proteins to the central lytic vacuole for hydrolysis, as active Rubisco within the intact chloroplast is highly resistant to proteolytic attack⁷.

This piecemeal degradation and remobilization are executed via distinct vesicular pathways:

1. **Rubisco-Containing Bodies (RCBs):** These are small, double-membrane vesicles (0.5 to 1.5 μ m in diameter) that bud directly from the chloroplast, encapsulating stromal proteins like Rubisco and glutamine synthase⁷⁵. The formation and transport of RCBs to the vacuole are strictly dependent on the core cytoplasmic autophagy machinery (encoded by *ATG* genes such as *ATG5* and *ATG7*)⁷³.
2. **Senescence-Associated Vacuoles (SAVs):** Distinct from the central vacuole, SAVs are small, highly acidic, lytic compartments that emerge exclusively during senescence⁷⁵. They exhibit intense cysteine protease activity, particularly driven by the senescence-specific enzyme SAG12⁷⁵. Unlike RCBs, the formation of SAVs operates entirely independently of the autophagic (*ATG*) pathway and involves the direct sequestration of chloroplast proteins for rapid degradation⁷⁵.

Once delivered to the vacuole, proteins are degraded by specific proteases, such as the aspartic protease CND41, which strictly requires the Rubisco structure to be denatured prior to cleavage⁷. The liberated free amino acids—primarily synthesized into mobile transport forms like glutamine (Gln) and asparagine (Asn)—are actively loaded into the phloem for systemic translocation to the developing seeds or apical meristems⁷. Thus, the observable phenotypic chlorosis is not a passive decay, but the direct macroscopic expression of massive, vacuole-mediated organelle dismantling designed to salvage nitrogen⁷⁸.

Phosphorus Deficiency: The PHR1-miR399 Axis and Anthocyanin Signatures

Phosphorus (Pi) is integral to ATP synthesis, the structural integrity of nucleic acid backbones, and phospholipid membranes⁷⁹. Because orthophosphate is highly immobile in soil, deficiencies are common. Plants experiencing P deficiency exhibit hallmark diagnostic symptoms: a severe stunting of shoot growth, a dramatic reconfiguration of root system architecture (RSA), and the distinct accumulation of purple anthocyanin pigments in the leaves and stems⁸.

The regulatory hub orchestrating this systemic bio-indicator response is the MYB-CC transcription factor **PHOSPHATE STARVATION RESPONSE 1 (PHR1)**⁸. Under adequate P conditions, the expression and activity of PHR1 are post-transcriptionally suppressed by SPX domain proteins, which physically interact with PHR1 to prevent it from binding to DNA⁷⁹. However, upon detecting intracellular Pi starvation, this inhibition is lifted, and PHR1 binds specifically to cis-acting regulatory elements (P1BS) in the promoters of hundreds of Pi-responsive genes⁷⁹.

The activation of the PHR1 pathway executes several highly visible, systemic bio-indicator responses:

1. **Anthocyanin Accumulation and Photoprotection:** PHR1 directly upregulates another transcription factor, **PAP1 (MYB75)**, which drives the biosynthesis of anthocyanins in the shoot tissues⁸². This deep purple pigmentation serves a critical physiological role as a photoprotectant, shielding the vulnerable photosynthetic apparatus from severe oxidative damage caused by excess light energy that can no longer be processed due to the lack of ATP synthesis⁹.
2. **Herbivory Resistance via Jasmonic Acid:** Phosphorus deficiency indirectly induces the expression of *JAZ10*, activating the jasmonic acid (JA) signaling pathway⁸. This unexpected biochemical shift drastically increases the plant's resistance to generalist insect herbivores. For example, the survival rate and weight of *Spodoptera littoralis* caterpillars feeding on severely P-deficient mutants drop catastrophically due to the accumulation of JA/JA-isoleucine defense compounds⁸.
3. **Root Architecture and Systemic Signaling via miR399:** PHR1 initiates the transcription of microRNA399 (miR399) in the shoot⁸. Remarkably, miR399 acts as a highly mobile, systemic signal, traveling downward through the phloem to the root system⁸. In the roots, miR399 specifically targets and degrades the mRNA of *PHO2 (AtUBC24)*, which encodes a ubiquitin E2 conjugating enzyme⁹. The destruction of *PHO2* derepresses high-affinity Pi transporters (PHTs) in the root epidermis, maximizing Pi uptake⁸¹. Simultaneously, the plant fundamentally alters its morphology: primary root elongation is arrested via auxin accumulation at the root apex, while the proliferation of dense lateral roots and root hairs is heavily stimulated, optimizing the root system for shallow topsoil foraging where Pi is typically most concentrated⁹. Furthermore, to conserve phosphorus, the plant actively replaces structural membrane phospholipids with non-phosphorus sulfolipids and galactolipids⁸.

Potassium Deficiency: Transporter Dynamics and Putrescine-Induced Necrosis

Potassium operates as a fundamental osmolyte, regulating stomatal aperture, the activation of dozens of critical enzymes, and the maintenance of cellular turgor. The primary vectors for

K^{+} uptake from the soil solution into the root periphery cells are the voltage-gated inward-rectifying channel **AKT1** and the high-affinity K^{+}/H^{+} symporter **HAK5**⁸⁷. The activity of both transport systems is highly orchestrated by a calcium-sensing regulatory

network; upon detecting external potassium decline, the protein kinase CIPK23 interacts with and phosphorylates these transporters to maximize their acquisition efficiency⁹⁰.

When soil potassium levels drop completely below the operational affinity thresholds of AKT1 and HAK5, the plant enters severe metabolic deficiency⁸⁸. The overt visual bio-indicator of advanced potassium deficiency is marginal leaf necrosis—a distinct "scorched" appearance along the outer edges of mature leaves, often accompanied by severe curling and structural collapse⁹⁴.

Crucially, this necrotic phenotype is not merely a passive result of dehydration or stomatal failure, but a biochemically active toxicity mechanism involving massive alterations in polyamine metabolism⁹⁷. Under severe K deficiency, plants massively upregulate the expression of **arginine decarboxylase (ADC)**, while simultaneously downregulating enzymes like spermidine synthase⁹⁴. ADC is the primary enzyme driving the biosynthesis of the diamine **putrescine**⁹⁶. As potassium levels plummet, putrescine rapidly accumulates to exponential levels within the leaf tissues—often sequestered in the enlarged central vacuoles of mesophyll cells—functioning as a makeshift intracellular organic cation to temporarily maintain osmotic pressure and charge balance⁹⁷. While low baseline levels of putrescine provide stress tolerance (e.g., in *Panax ginseng* or hybrid poplars under salt stress), the hyper-accumulation induced by chronic potassium starvation results in severe cellular toxicity⁹⁹. This toxic accumulation

triggers the massive generation of hydrogen peroxide (H_2O_2) and subsequent lipid peroxidation, physically destroying the cell membranes and directly causing the hallmark necrotic tissue death (scorch) observed at the leaf margins⁹⁵.

7. Nutrient Antagonism and Physiological Competition

Accurately interpreting plant bio-indicators requires an understanding of soil chemistry beyond absolute elemental concentrations. Frequently, a plant will exhibit severe, phenotypic deficiency symptoms for a specific nutrient even when rudimentary chemical soil tests indicate that the nutrient is abundantly present in the soil. This paradox is driven by nutrient antagonism—the competitive interference and chemical precipitation between different elemental ions within the rhizosphere or during root epidermal uptake¹⁰³.

Mulder's Chart and Elemental Interplay

The complex, multi-directional web of synergistic and antagonistic relationships between soil elements is classically mapped on **Mulder's Chart**, a foundational diagram published in 1953 mapping nutrient interactions in fruit orchards¹⁰⁴. Two of the most agronomically significant antagonisms that confound simple chemical diagnostics involve phosphorus/zinc and potassium/magnesium dynamics.

Antagonistic Pair	Geochemical / Physiological Mechanism	Phenotypic Bio-Indicator Consequence
High Phosphorus vs. Zinc	Phosphorus-induced zinc	Even in zinc-sufficient soils,

	deficiency occurs when excessive orthophosphate reacts directly with soil zinc to form highly insoluble zinc phosphate complexes in the rhizosphere, physically removing zinc from the bioavailable pool. Furthermore, excess P downregulates systemic signaling for micronutrient uptake at the root surface ²³ .	heavy P fertilization induces severe zinc deficiency symptoms. Plants exhibit distinct interveinal chlorosis in newly emerging leaves, severely shortened internodes (a condition known as "rosetting"), and stunted overall growth ¹⁰⁹ .
High Potassium vs. Magnesium	Potassium (K^+) and Magnesium (Mg^{2+}) fiercely compete for binding sites on non-selective cation channels and dedicated transporter proteins at the root epidermis. Potassium, being highly mobile and polarizable, outcompetes magnesium for uptake ¹⁰³ .	Heavy K fertilization physically blocks Mg uptake. The plant exhibits classic Mg deficiency: distinct interveinal chlorosis on older leaves (as Mg is highly mobile and mobilized to new tissue), eventually leading to tissue necrosis and severely disrupting chlorophyll synthesis, as Mg is the central coordinating atom of the chlorophyll molecule ¹⁰³ .

Understanding these specific antagonistic mechanisms is absolutely vital for correct diagnosis; attempting to treat a visual zinc deficiency by simply broadcasting more zinc onto a high-phosphorus soil will fail, as the fundamental issue is stoichiometric precipitation, not absolute elemental absence²³.

8. Phytogeographical Bio-Indicators and Ecological Scoring Systems

Beyond diagnosing nutrient deficiencies in individual agricultural crops, the presence, abundance, and community structure of specific wild plant species can serve as highly reliable ecological bio-indicators for mapping soil pH and inherent nutrient fertility across broad landscapes. This approach relies on the fundamental ecological principle that species distribution is strictly constrained by abiotic physiological tolerance limits⁵.

Ellenberg and Landolt Indicator Values

The most robust theoretical frameworks for floristic bio-indication are the **Ellenberg Indicator**

Values (EIVs), originally developed for the vascular flora of Central Europe in 1974, and the **Landolt Indicator Values (LIVs)**, developed primarily for the highly diverse Alpine regions of Switzerland⁵.

These sophisticated systems assign ordinal numerical scores to thousands of plant species based on their realized ecological optima along specific, critical environmental gradients⁵. The most agronomically and chemically critical indices are:

- **R-value (Reaction):** Indicates a species' physiological preference for soil pH. A score of R1 signifies an extreme acidophyte (an organism restricted to highly acidic, low-pH soils with high aluminum availability), while R9 indicates an obligate calcicole (an organism restricted to highly alkaline, carbonate-rich soils)¹¹⁴.
- **N-value (Nitrogen/Nutrients):** Reflects the species' preference for overall soil fertility, biomass productivity, and nutrient saturation, rather than exclusively mineral nitrogen availability³⁸.

By conducting a simple floristic inventory of a specific site, an environmental assessor can calculate the mean indicator value of the entire plant community—often mathematically weighted by the relative abundance or physical cover of each individual species to improve accuracy¹¹³. High aggregate R-values reliably indicate calcareous, alkaline soil chemistry, while high N-values correlate strongly with gross primary productivity and nutrient saturation, often accurately predicting total herbage yield⁶¹.

While these phytogeographical systems represent an elegant, low-cost alternative to exhaustive chemical soil testing grids, they possess inherent structural limitations. First, EIVs and LIVs are based largely on historical expert consensus rather than strict empirical, physical calibration¹¹⁶. Second, the values reflect the *realized* ecological niche of the plant (which is heavily influenced by inter-species competition) rather than its absolute *fundamental* physiological niche¹¹⁴.

Furthermore, averaging ordinal variables (which lack equal class widths) is statistically problematic, and applying ANOVA tests to PCA scores derived from these indicators often violates assumptions of statistical independence⁶. Crucially, their accuracy diminishes significantly when transferred to biogeographical regions far outside their native calibration zones, often necessitating regional statistical recalibration to align the indicator scores with actual physical pH measurements¹¹³. Finally, while they show incredibly strong correlations for evaluating *spatial* differences across a landscape, recent studies utilizing Canopy Stability Indices (CSIs) demonstrate that they possess limited sensitivity for detecting subtle, long-term *temporal* changes in soil chemistry at a single site¹¹³. Nonetheless, they remain unparalleled tools for broad environmental monitoring, habitat assessment, and historical landscape reconstruction⁶.

***Hydrangea macrophylla* as an In Situ Aluminum and pH Sensor**

Perhaps the most famous, chemically complex, and visually striking single-species bio-indicator of soil chemistry is the bigleaf hydrangea (*Hydrangea macrophylla*). The sepals (the showy, petal-like structures of the inflorescence) of this plant exhibit a dramatic color polymorphism—ranging from deep crimson red to vibrant, striking blue—that is entirely

dictated by soil pH and aluminum bioavailability²².

Unlike standard genetic color variations found in most flowers, all colors in *H. macrophylla* arise from the exact same primary anthocyanin pigment: **delphinidin 3-O-glucoside**, existing alongside specific copigments, primarily 5-O-caffeoylquinic acid (neochlorogenic acid) and 3-O-caffeoylquinic acid (chlorogenic acid)¹²¹. The mechanism of the macroscopic color shift is a highly intricate function of rhizosphere biogeochemistry interacting directly with intracellular vacuolar coordination chemistry:

1. **Alkaline to Neutral Soils (pH > 6.5):** In non-acidic environments, aluminum in the soil exists as highly insoluble aluminum hydroxide ($Al(OH)_3$) precipitates, rendering it completely unavailable for root absorption¹²¹. In the absolute absence of aluminum, the delphinidin pigment within the sepal vacuoles remains unchelated, existing stably as the red flavylium cation. The vacuolar pH of these red cells is actively maintained at a highly acidic 3.3, and the total aluminum content is negligible (approximately 0.03 equivalents relative to anthocyanin)¹²¹. Under these conditions, the sepals remain pink or red¹²².
2. **Acidic Soils (pH < 5.5):** As soil pH drops, aluminum ions (Al^{3+}) become highly soluble and biologically mobile in the soil solution. The hydrangea aggressively absorbs these acidic ions, rapidly translocating them through the vascular system directly to the vacuoles of the sepal cells¹²¹. Within the vacuole, the Al^{3+} ion coordinates specifically with the 3',4'-dihydroxy groups on the B-ring of the delphinidin molecule. This powerful chelation event forces a deprotonation, converting the red delphinidin to its blue quinoidal base anion²². To stabilize this intensely blue metal-ligand complex, large quantities of the copigment (e.g., 5-O-caffeoylquinic acid) actively intercalate and stack around the complex, creating a massive, stable, supramolecular "hydrangea-blue" pigment structure²². This complexation process also induces a significant measurable shift in the vacuolar pH, raising it to 4.1 in the blue cells, which contain approximately 1.2 equivalents of Al^{3+} and an overwhelming 13 equivalents of 5-O-acylquinic acids relative to the anthocyanin¹²¹.

Because the presence of the blue pigment complex is strictly contingent upon the soil being acidic enough to physically solubilize aluminum, *H. macrophylla* serves as a highly accurate, living litmus test for both soil acidification and heavy metal mobility, offering a low-cost visual indicator for environmental monitoring¹²².

9. Instrumental Quantification of Bio-Indicators

While the visual assessment of phenotypic bio-indicators—such as the chlorosis associated with nitrogen or magnesium deficiencies—provides immediate, qualitative insight, it is inherently subjective and highly susceptible to observer bias¹²⁷. To bridge the gap between subjective visual scoring and destructive, time-consuming laboratory tissue extraction analysis (such as the Kjeldahl digestion method), field agronomists employ precision portable sensors, most notably the Soil-Plant Analysis Development (SPAD) meter¹²⁹.

A SPAD meter is a portable, non-destructive optical sensor that clamps directly onto a living, intact leaf. It evaluates the relative greenness and photosynthetic capacity of the tissue by measuring the physical transmittance of light emitted by two highly specific light-emitting diodes (LEDs)¹²⁹:

1. **650 nm (Red Light):** This specific wavelength represents the absolute peak absorbance for chlorophyll molecules. Light at this frequency is heavily absorbed by the photosynthetic pigments present in the mesophyll¹³⁰.
2. **940 nm (Infrared Light):** Chlorophyll does not absorb infrared light. This wavelength safely passes through the leaf and serves as an essential internal baseline reference, automatically adjusting for confounding variables in leaf morphology, such as cuticle thickness, specific leaf weight, cellular water status, and epidermal variations¹³⁰.

A microprocessor within the device calculates the ratio of these two transmittances to produce a unitless SPAD value (typically ranging from 0 to 100)¹²⁷. Because the absolute physical concentration of leaf chlorophyll is inextricably linked to the plant's overall nitrogen status (as N is a core structural component of the porphyrin ring of the chlorophyll molecule), SPAD readings provide a highly accurate, real-time proxy for leaf nitrogen content¹²⁹.

Extensive empirical evaluations across diverse species confirm that SPAD values exhibit a remarkably robust polynomial quadratic correlation ($R^2 \geq 0.84$) with total chemically extracted chlorophyll and actual foliar nitrogen percentages¹²⁷. For instance, readings dropping below species-specific critical thresholds (e.g., SPAD < 25 in species like *Quercus robur* or *Fagus sylvatica*) unequivocally indicate severe impairment of the photosynthetic apparatus and critical N deficiency (often correlating to <1.5% foliar nitrogen)¹²⁹. This allows for precise, on-demand fertigation adjustments in the field—maximizing yield while minimizing nitrogen runoff—without the days of delay associated with traditional laboratory analysis¹²⁸.

10. Conclusion

The evaluation of soil ecosystems and organic amendments in the absence of advanced analytical infrastructure is not a compromise of scientific rigor, but rather an exercise in applied biogeochemistry and precise physiological observation. Rudimentary chemical indicators, such as the zinc-reduction Griess assay for nitrates or the sodium cobaltinitrite turbidimetric test for potassium, leverage fundamental thermodynamic principles and transition-metal complexation to render invisible macronutrients visibly quantifiable in the field.

Concurrently, the utilization of plant bio-indicators elevates diagnostics from mere chemical concentration to actual biological availability. The manifestation of leaf chlorosis is not a passive decay, but a highly regulated autophagic dismantling of Rubisco via senescence-associated vacuoles to ensure the survival and remobilization of scarce nitrogen. The deep purple flush of a phosphorus-starved leaf is the systemic result of miR399 signaling mediated by the PHR1 transcription factor, drastically altering the plant's metabolic architecture. By integrating the immediate chemical insights provided by colorimetric soil extractions with the deep, time-integrated metabolic narratives told by the flora—whether through the pH-dependent chelation of aluminum in *Hydrangea macrophylla* or the SPAD-quantified chlorophyll density of an agricultural crop—practitioners can achieve a nuanced, highly accurate understanding of

rhizosphere dynamics and long-term soil health.

Works cited

1. 23 Biology Of Organic Fertilizers.pdf
2. Patterns and thresholds for soil pH across Europe in relation to soil, <https://nora.nerc.ac.uk/id/eprint/540384/1/N540384JA.pdf>
3. High-efficiency high performance liquid chromatographic analysis of, https://www.researchgate.net/publication/51211601_High-efficiency_high_performance_liquid_chromatographic_analysis_of_red_wine_anthocyanins
4. A Rapid Turbidimetric Potassium Test Modified for Use with the, https://www.researchgate.net/publication/244993530_A_Rapid_Turbidimetric_Potassium_Test_Modified_for_Use_with_the_Pressurized_Hot-Water_Extraction
5. (PDF) Ellenberg Indicator Values Disclose Complex Environmental, https://www.researchgate.net/publication/367323581_Ellenberg_Indicator_Values_Disclose_Complex_Environmental_Filtering_Processes_in_Plant_Communities_along_an_Elevational_Gradient
6. Landolt Indicator Values in Modern Research: A Review - MDPI, <https://www.mdpi.com/2071-1050/15/12/9618>
7. Nitrogen Recycling and Remobilization Are Differentially Controlled, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2442554/>
8. Phosphate Deficiency Induces the Jasmonate Pathway and, <https://academic.oup.com/plphys/article/171/1/632/6115091>
9. Overexpression of a Phosphate Starvation Response AP2/ERF, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2018.01186/full>
10. Evaluation of some Conventional methods for Estimating Available, <https://scispace.com/pdf/evaluation-of-some-conventional-methods-for-estimating-3dbnw0d8wl.pdf>
11. B.Tech. (Environmental Engineering) Laboratory Manual, <https://people.iitism.ac.in/~download/lab%20manuals/evs/UG%20Lab%20Manual%20-%20Dept%20of%20ESE.pdf>
12. tropical agricultural science - Pertanika UPM, [http://www.pertanika.upm.edu.my/resources/files/Pertanika%20PAPERS/JTAS%20Vol.%2047%20\(4\)%20Nov.%202024/21%20JTAS-3000-2023.pdf](http://www.pertanika.upm.edu.my/resources/files/Pertanika%20PAPERS/JTAS%20Vol.%2047%20(4)%20Nov.%202024/21%20JTAS-3000-2023.pdf)
13. an abstract of the thesis of - Oregon State University, <https://ir.library.oregonstate.edu/downloads/fj2365848>
14. Review of the Innovations and Challenges in Developing Rapid, https://www.researchgate.net/publication/387555500_Review_of_the_Innovations_and_Challenges_in_Developing_Rapid_Colorimetry_and_Turbidity_NPK_Soil_Test_Kits_for_Commercial_Soil_Nutrient_Analysis
15. Optimizing ammonium acetate procedure for determining available, https://dlsr.iut.ac.ir/article_3416_973df41790d68dbe89086784244fba21.pdf
16. Extractable Soil Phosphorus Concentrations and Creeping, <https://digitalcommons.lib.uconn.edu/cgi/viewcontent.cgi?article=1003&context=>

[plsc_articles](#)

17. A case study of crop planting zones in central China - ResearchGate,
https://www.researchgate.net/publication/337455584_Application_of_a_universal_soil_extractant_for_determining_the_available_NPK_A_case_study_of_crop_planting_zones_in_central_China
18. Soil Test Methods From the Southeastern United States - AESL,
<https://aesl.ces.uga.edu/sera6/PUB/Methodsmanualfinalsera6.pdf>
19. Acid and Alkaline Soil Practical | PDF - Scribd,
<https://www.scribd.com/document/1065642749/Acid-and-Alkaline-Soil-Practical>
20. Phosphorus loss (kg/ha) from site 1 and 3 and soil-test P results,
https://www.researchgate.net/figure/Phosphorus-loss-kg-ha-from-site-1-and-3-and-soil-test-P-results-mg-kg-and-loss_tbl3_228625915
21. Effect of Long-Term Paddy-Upland Yearly Rotations on Rice ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3417201/>
22. The chemical mechanism for Al³⁺ complexing with delphinidin,
https://www.researchgate.net/publication/43159523_The_chemical_mechanism_for_Al3_complexing_with_delphinidin_A_model_for_the_bluing_of_hydrangea_sepal
23. Phosphorus-zinc Interactions in the Soil Environment and their,
<https://journalijpss.com/index.php/IJPSS/article/download/5953/12848/9482>
24. Test Soil at Home: Easy DIY Tips for Accurate Results,
<https://homegrown-garden.com/blogs/blog/test-soil-at-home>
25. [Solved] Which chemical reaction occurs when you mix vinegar (acetic,
<https://testbook.com/question-answer/which-chemical-reaction-occurs-when-you-mix-vinegar--666d71ed052b77821b085e68>
26. Open Soil Atlas - mit:forschen!,
[https://www.mitforschen.org/sites/default/files/assets/projekte/user-4879/pdf/OSA%20Final%20Paper%20\(Englisch\).pdf](https://www.mitforschen.org/sites/default/files/assets/projekte/user-4879/pdf/OSA%20Final%20Paper%20(Englisch).pdf)
27. Baking Soda and Distilled Vinegar PH +- Levels : r/Hydroponics,
https://www.reddit.com/r/Hydroponics/comments/12ppmzw/baking_soda_and_distilled_vinegar_ph_levels/
28. Ask an Expert: Baking soda and vinegar - Page 2 - Science Buddies,
<https://www.sciencebuddies.org/science-fair-projects/ask-an-expert/viewtopic.php?t=2359&start=30>
29. Ph Science Fair Projects,
<https://enflomaplata.uep.edu.py/index.jsp/TNOV5L/705691/Ph%20Science%20Fair%20Projects.pdf>
30. The Medicinal Chemistry of Cyanidin and its Glycoside Derivatives,
https://www.researchgate.net/publication/392620853_The_Medicinal_Chemistry_of_Cyanidin_and_its_Glycoside_Derivatives_Focus_on_the_Antiproliferative_and_Potential_Anticancer_Activity
31. Flavonoid extracts from chrysanthemum with appropriate ... - SciOpen,
https://www.sciopen.com/local/article_pdf/10.1016/j.hpj.2023.10.002.pdf
32. Pressurized solvent extraction and monolithic column-HPLC/DAD,
https://www.researchgate.net/publication/5480385_Pressurized_solvent_extraction

- [on_and_monolithic_column-HPLCDAD_analysis_of_anthocyanins_in_red_cabbage](#)
33. Colonizing Taxa (Weeds) as Sources of Natural Pigments and Dyes,
https://www.researchgate.net/publication/362164688_Colonizing_Taxa_Weeds_as_Sources_of_Natural_Pigments_and_Dyes_Colonizing_Taxa_Weeds_as_Sources_of_Natural_Pigments_and_Dyes
 34. Red Cabbage pH indicator Question : r/chemistry - Reddit,
https://www.reddit.com/r/chemistry/comments/16n7xj8/red_cabbage_ph_indicator_question/
 35. Flavonoids: Chemistry, Biochemistry and Applications,
<https://dokumen.pub/flavonoids-chemistry-biochemistry-and-applications-9780849320217.html>
 36. 11.1 What is the pH value? | Acids, bases and the pH value | Siyavula,
<https://www.siyavula.com/read/za/natural-sciences/grade-9/acids,-bases-and-the-ph-value/11-acids,-bases-and-the-ph-value>
 37. Anthocyanin colour changes at different pH of the buffer solution,
https://www.researchgate.net/figure/Anthocyanin-colour-changes-at-different-pH-of-the-buffer-solution_fig1_351337488
 38. Assessing the habitat conservation status by soil parameters and,
<https://iforest.sisef.org/contents/?id=ifor0963-007>
 39. Indicator values suggest decreasing habitat quality in Swiss dry,
https://www.tuexenia.de/publications/tuexenia/Tuexenia_2019_NS_039_0315-0334.pdf
 40. A colorimetric method for use within portable test kits for nitrate,
https://www.researchgate.net/publication/312161186_A_colorimetric_method_for_use_within_portable_test_kits_for_nitrate_determination_in_various_water_matrices
 41. A Simple, Ecofriendly, and Fast Method for Nitrate Quantification in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11209534/>
 42. Photometric Determination of Nitrite in Soil Using Griess Method,
<https://www.sigmaaldrich.com/DE/en/technical-documents/protocol/analytical-chemistry/photometry-and-reflectometry/determination-of-nitrite-in-soils>
 43. Orthophosphate - Research Analytical Laboratory,
<https://ral.cfans.umn.edu/orthophosphate>
 44. A SIMPLE SPECTROPHOTOMETRIC METHOD FOR THE,
<https://ejournal.undiksha.ac.id/index.php/JST/article/view/13940>
 45. The Blue Molybdenum Reaction for the Determination of Phosphate,
https://www.researchgate.net/publication/374422525_The_Blue_Molybdenum_Reaction_for_the_Determination_of_Phosphate_in_Natural_Water_and_Detergent_Samples
 46. Phosphate Analysis,
<https://wateranalysis.christineingersoll.bergbuils.domains/phosphate-analysis/>
 47. (PDF) Determination of phosphorus by phosphorus molybdenum,
https://www.researchgate.net/publication/374279302_Determination_of_phosphorus_by_phosphorus_molybdenum_blue_spectrophotometry_by_bismuth_antimony

[ny_sensitization](#)

48. A Colorimetric Dip Strip Assay for Detection of Low Concentrations,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8125474/>
49. Standard Operating Procedure-Phosphate Analysis | PDF - Scribd,
<https://www.scribd.com/document/992123652/Standard-Operating-Procedure-phosphate-Analysis>
50. Exploring a low-cost turbidimetric sensor for available potassium,
<https://pubs.rsc.org/ay/article/16/46/8037/837219/Exploring-a-low-cost-turbidimetric-sensor-for>
51. Model STH Series,
<https://www.itm.com/pdfs/cache/www.itm.com/5010-01/manual/5010-01-manual.pdf>
52. A Low-Cost Portable Microcontrolled Nephelometer for Potassium,
<https://www.scielo.br/j/jbchs/a/MBv9SpnDsqwmXSRS76h96Nf/?lang=en>
53. A novel method for extracting potassium (K) from K-poor and sodium,
<https://par.nsf.gov/servlets/purl/10573752>
54. Method for measuring kalium in soil by tetraphenylboron sodium,
<https://patents.google.com/patent/CN101271072B/en>
55. SMART3 BLE Soil - LaMotte,
<https://lamotte.com/wp-content/uploads/2023/10/1985-06-BLE-MN.pdf>
56. Colorimetric Potassium Analysis in Soils | PDF - Scribd,
<https://www.scribd.com/document/614230759/Colorimetric-Determination-K>
57. Sodium Cobaltinitrite (CAS 13600-98-1) | ACS Reagent Grade,
<https://www.benchchem.com/product/b080926>
58. Use of Textile Fibres in Microscopic Qualitative Chemical Analysis-,
<http://lib3.dss.go.th/fulltext/Journal/analyst/Analyst1916-1920/1918/505vol43,1918p148-154.pdf>
59. A new spectrophotometric method for quantification of potassium,
https://www.academia.edu/32736218/A_new_spectrophotometric_method_for_quantification_of_potassium_solubilized_by_bacterial_cultures
60. Analytical Chemistry.,
<https://pubs.rsc.org/en/content/articlepdf/1910/ca/ca9109805340>
61. Untersuchungen zur Anwendung der Zeigerwerte nach Ellenberg,
https://www.researchgate.net/publication/272166264_Untersuchungen_zur_Anwendung_der_Zeigerwerte_nach_Ellenberg
62. onion epidermal cells: Topics by Science.gov,
<https://www.science.gov/topicpages/o/onion+epidermal+cells>
63. (PDF) Antiproliferative and Apoptotic Potential of Cyanidin-Based,
https://www.researchgate.net/publication/316638862_Antiproliferative_and_Apoptotic_Potential_of_Cyanidin-Based_Anthocyanins_on_Melanoma_Cells
64. Development of a Cost-Effective Method for Nitrate and Nitrite,
<https://pubs.acs.org/doi/abs/10.1021/jf904558c>
65. Improving Sustainability of the Griess Reaction by Reagent ... - MDPI,
<https://www.mdpi.com/2073-4360/14/3/464>
66. Part 1 (Mosser Lee Soil Test Kit -Sodium Tetraphenylborate Method),

- https://www.researchgate.net/publication/350610700_Testing_for_Potassium_in_Upper_Midwest_United_States_Soils_-Part_1_Mosser_Lee_Soil_Test_Kit_-Sodium_Tetraphenylborate_Method
67. Degradation of Rubisco and other chloroplast proteins under abiotic, https://www.researchgate.net/publication/251181329_Degradation_of_Rubisco_and_other_chloroplast_proteins_under_abiotic_stress
 68. Chloroplast Protein Degradation Pathways During Leaf Senescence, <https://www.mdpi.com/2223-7747/15/17/2576>
 69. Molecular basis of nitrogen starvation-induced leaf senescence, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2022.1013304/pdf>
 70. Autophagy, plant senescence, and nutrient recycling, <https://academic.oup.com/jxb/article/65/14/3799/2877429>
 71. Common bean (*Phaseolus vulgaris* L.) gas exchange capacity, <https://hrcak.srce.hr/file/429299>
 72. Senescence-associated proteins and nitrogen remobilization ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9276853/>
 73. Mobilization of Rubisco and Stroma-Localized Fluorescent Proteins, <https://academic.oup.com/plphys/article/148/1/142/6107379>
 74. Chloroplast Protein Degradation in Senescing Leaves - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6593067/>
 75. (PDF) Chloroplast Protein Degradation: Involvement of Senescence, https://www.researchgate.net/publication/249011433_Chloroplast_Protein_Degradation_Involvement_of_Senescence-Associated_Vacuoles
 76. Stromal protein degradation is incomplete in *Arabidopsis thaliana*, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3724497/>
 77. Mobilization of Rubisco and Stroma-Localized Fluorescent Proteins, <https://www.semanticscholar.org/paper/Mobilization-of-Rubisco-and-Stroma-Localized-of-to-Ishida-Yoshimoto/97e7712e538b739718c369186da56f4f906a9394>
 78. Chloroplast Recycling and Plant Stress Tolerance - IntechOpen, <https://www.intechopen.com/chapters/89287>
 79. Integration of P, S, Fe, and Zn nutrition signals in *Arabidopsis thaliana*, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4411997/>
 80. Sugar Signaling in Root Responses to Low Phosphorus Availability, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3135921/>
 81. Full article: A new insight into root responses to external cues, <https://www.tandfonline.com/doi/full/10.1080/15592324.2015.1049791>
 82. Molecular Regulation for Maintenance of Plant Phosphate, <https://www.preprints.org/manuscript/202306.0686/v1/download>
 83. MdSnRK1.1 interacts with MdJAZ18 to regulate sucrose-induced, <https://academic.oup.com/jxb/article/68/11/2977/3855003>
 84. Phosphate (Pi) Starvation Up-Regulated GmCSN5A/B Participates, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8623310/>
 85. Signaling Network in Sensing Phosphate Availability in Plants, <https://www.annualreviews.org/doi/pdf/10.1146/annurev-arplant-042110-103849>

86. Phosphate starvation: response mechanisms and solutions - NSF PAR,
<https://par.nsf.gov/servlets/purl/10516776>
87. Regulation of K⁺ Nutrition in Plants - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6435592/>
88. Physical and functional interactions of the potassium uptake protein,
<https://www.biorxiv.org/content/10.1101/2025.10.03.680249v1.full-text>
89. Molecular mechanisms involved in plant adaptation to low K⁺,
<https://www.ovid.com/journals/jebot/fulltext/10.1093/jxb/ert402-molecular-mechanisms-involved-in-plant-adaptation-to-low-k>
90. [PDF] Osmotic Stress Responses and Plant Growth Controlled by,
<https://www.semanticscholar.org/paper/Osmotic-Stress-Responses-and-Plant-Growth-by-in-Osakabe-Arinaga/5ba9787fc861d28cb421ae47d4b66550dd6a3d9e>
91. Natural variation in potassium deficiency responses among,
<https://academic.oup.com/pcp/article/66/6/956/8119714>
92. (PDF) High-Affinity K⁺ Transport in Arabidopsis: AtHAK5 and AKT1,
https://www.researchgate.net/publication/43299255_High-Affinity_K_Transport_in_Arabidopsis_AtHAK5_and_AKT1_Are_Vital_for_Seedling_Establishment_and_Postgermination_Growth_under_Low-Potassium_Conditions
93. Potassium transporter KUP9 participates in K⁺ distribution in roots,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10441886/>
94. Ethylene Evolution and Polyamine Accumulation by Tomato,
<https://journals.ashs.org/downloadpdf/view/journals/jashs/114/4/article-p651.pdf>
95. (PDF) Potassium deficiency inhibits leaf growth and promotes leaf,
https://www.researchgate.net/publication/357313411_Potassium_deficiency_inhibits_leaf_growth_and_promotes_leaf_necrotic_spots_in_Neolamarckia_cadamba_Roxb_Bosser
96. Current Advances in Plant Growth Promoting Bacteria Alleviating,
<https://www.mdpi.com/2076-3417/10/20/7025>
97. Polyamine Metabolism in Scots Pine Embryogenic Cells under,
<https://www.mdpi.com/2073-4409/10/5/1244>
98. (PDF) Putrescine: A Key Metabolite Involved in Plant Development,
https://www.researchgate.net/publication/359166475_Putrescine_A_Key_Metabolite_Involved_in_Plant_Development_Tolerance_and_Resistance_Responses_to_Stress
99. Plant Physiology and Biochemistry,
<https://repositorio.unesp.br/bitstreams/15e415ea-35fc-492c-84c7-3b3a021c50a2/download>
100. NASA space Medicine & Biology Aei 9 Medicine & Biology Aerospa,
<https://ntrs.nasa.gov/api/citations/19830021718/downloads/19830021718.pdf?attachment=true>
101. Exogenous Putrescine Enhances Salt Tolerance and Ginsenosides,
<https://www.mdpi.com/2223-7747/10/7/1313>
102. (PDF) Enhancing salt tolerance in a hybrid poplar (Populus nigra x,
https://www.researchgate.net/publication/392475756_Enhancing_salt_tolerance_in_a_hybrid_poplar_Populus_nigra_x_maximowiczii_through_exogenous_putrescine

- [e_application_Growth_physiological_and_biochemical_responses](#)
103. Nutritional antagonisms and deficiencies in CBD culture - CBD.fr,
<https://www.cbd.fr/blog/en/nutritional-antagonisms-when-an-excess-causes-a-d-efficiency-in-cbd-cultivation/>
 104. Mechanisms for nutrient interactions from organic amendments and,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10081454/>
 105. Nutrient Antagonism in Cannabis: Chart, Causes and the Fix,
<https://www.rxgreentechnologies.com/blog/nutrient-antagonism-cannabis/>
 106. How are nutrients taken up by the crop? - NutriAg,
<https://www.nutriag.com/nutrient-uptake/>
 107. (PDF) Phosphorus-induced zinc deficiency in wheat pot-grown on,
https://www.researchgate.net/publication/325955575_Phosphorus-induced_zinc_deficiency_in_wheat_pot-grown_on_noncalcareous_and_calcareous_soils_of_different_properties
 108. Phosphorus build up in soil affecting zinc availability to plants in rice,
<https://arrworyza.com/AdminPanel/download/Phosphorus%20build%20up%20in%20soil%20affecting%20zinc%20availability%20to%20plants.pdf>
 109. Sight Unseen: Zinc Deficiency in Soil | SoilBeat,
<https://www.soilbeat.com/blog/sight-unseen-zinc-deficiency-in-soil>
 110. agriculture - Semantic Scholar,
<https://pdfs.semanticscholar.org/2000/6eef031fde45d689c7aa2cb073b3a43a84aa.pdf>
 111. (PDF) Potassium induced suppression of magnesium uptake and,
https://www.researchgate.net/publication/382877178_Potassium_induced_suppression_of_magnesium_uptake_and_translocation_is_limited_in_oat_Avena_sativa_L
 112. Decoding Plant Metabolomic Response to Potassium and Nutrient,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12460989/>
 113. Validity of Ellenberg indicator values judged from physico-chemical,
https://www.researchgate.net/publication/40130330_Validity_of_Ellenberg_indicators_or_values_judged_from_physico-chemical_field_measurements
 114. Ellenberg Indicator Values Disclose Complex Environmental ... - MDPI,
<https://www.mdpi.com/2079-7737/12/2/161>
 115. Landscape patterns of indicator plants for soil acidity in the Bavarian,
http://www.bayceer.uni-bayreuth.de/bod/de/pub/html/20902Schmidtlein_Ewald2003.pdf
 116. Chapter 10: Plants as indicators of soil chemical properties,
https://literatur.thuenen.de/digbib_extern/dn061473.pdf
 117. Prediction of yield in the Rothamsted Park Grass Experiment by,
https://www.academia.edu/32462213/Prediction_of_yield_in_the_Rothamsted_Park_Grass_Experiment_by_Ellenberg_indicator_values
 118. Experience with using Ellenberg's R indicator values in Slovakia,
<https://d-nb.info/1342729404/34>
 119. How robust are community-based plant bioindicators? Empirical,
<https://d-nb.info/1218959258/34>
 120. (PDF) Vegetation Dynamics Studies Based on Ellenberg and,

- https://www.researchgate.net/publication/384783269_Vegetation_Dynamics_Studies_Based_on_Ellenberg_and_Landolt_Indicator_Values_A_Review
121. Identification of Seven Key Structural Genes in the Anthocyanin, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9497972/>
 122. Insight into chemical mechanisms of sepal color development and, https://www.researchgate.net/publication/349193235_Insight_into_chemical_mechanisms_of_sepal_color_development_and_variation_in_hydrangea
 123. Metal Complex Pigment Involved in the Blue Sepal Color, <https://pubs.acs.org/doi/10.1021/acs.jafc.5b02368>
 124. Thermodynamic and Kinetic Insights into Ternary Complexation, <https://colab.ws/articles/10.1016%2Fj.dyepig.2026.113876>
 125. and Red-Colored Sepal Cells of Hydrangea macrophylla, <https://www.tandfonline.com/doi/pdf/10.1271/bbb.80831>
 126. Multiple transcription factors and genes jointly regulate the light, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13059372/>
 127. Correlation of Visual Quality Grading and SPAD Reading of Green, https://www.researchgate.net/publication/232884093_Correlation_of_Visual_Quality_Grading_and_SPAD_Reading_of_Green-Leaved_Foliage_Plants
 128. Use of a chlorophyll meter and plant visual aspects for nitrogen, https://www.researchgate.net/publication/242762434_Use_of_a_chlorophyll_meter_and_plant_visual_aspects_for_nitrogen_management_in_tomato_fertigation
 129. The Potential of a Chlorophyll Content SPAD Meter to, <https://auf.isa-arbor.com/content/isa/34/2/89.full.pdf>
 130. A Computer-Vision-Based Approach for Nitrogen Content Estimation, <https://www.mdpi.com/2077-0472/11/8/766>
 131. Predictive capability of a leaf optical meter for determining leaf, <https://www.osti.gov/servlets/purl/1829084>
 132. The Potential of a Chlorophyll Content SPAD Meter to Quantify, <https://auf.isa-arbor.com/content/34/2/89>
 133. Use of a chlorophyll meter and plant visual aspect for nitrogen, https://horticultureresearch.net/jah/2006_8_1_8_11.PDF