

Comprehensive Swine Husbandry: Rearing Systems, Precision Nutrition, Behavioral Ecology, Waste Valorization, and Parasitic Biosecurity

Swine Rearing Systems and Environmental Engineering

Modern domestic swine (*Sus scrofa domesticus*) production requires precise environmental engineering, architectural design, and stocking management to optimize physiological throughput, herd biosecurity, and welfare¹. Production facilities are broadly categorized into intensively managed indoor confinement buildings—featuring slatted concrete or plastic flooring over deep pit slurries—and extensive or semi-intensive structures that utilize solid flooring bedded with straw, wood shavings, or sawdust². Allocation of usable floor space directly influences social stability, feed conversion efficiency, and pathogen pressure¹.

Overcrowding precipitates chronic physiological stress, elevates baseline circulating glucocorticoids, and exacerbates vices such as tail, flank, and ear biting⁴.

In nursery units housing weaned piglets from approximately 6 kg to 25–30 kg, standard intensive systems allocate 0.30 to 0.40 m² (3.2 to 4.3 sq ft) per piglet on fully slatted flooring¹. For grower-finisher cohorts transitioning from 30 kg to market weights exceeding 125–135 kg, space allocations increase nonlinearly¹. Fully slatted confinement pens provide 0.65 to 0.75 m² (7.0 to 8.0 sq ft) per animal, whereas solid-floored or bedded housing requires 0.85 to 1.00 m² (9.1 to 10.8 sq ft) to allow the distinct establishment of lying, feeding, and dunging territories³.

Breeding stock management has undergone significant structural transformation. Concerns over physical constraint in traditional individual gestation crates have driven regulatory shifts toward group-housing systems². Under statutory frameworks such as California Proposition 12 and European Union Council Directive 2008/120/EC, breeding sows and gilts must be housed in pens permitting unrestrained posture, including turning around without impediment⁶.

Proposition 12 mandates a minimum continuous floor allowance of 24 sq ft (2.23 m²) per breeding female⁶. Deep-bedded hoop barns recommended by the Midwest Plan Service (MWPS) similarly benchmark 24 sq ft per animal within bedded loafing packs, excluding feed stalls². Furthermore, crate confinement during the farrowing and lactating periods is increasingly constrained; standards require that sows are not moved into farrowing stalls earlier than five days prior to their projected farrowing date, encouraging loose or temporarily confined farrowing pen architectures⁷.

Porcine environmental engineering must accommodate the species' distinct thermoregulatory biology. Pigs possess non-functional apocrine sweat glands and have low

lung-to-body-volume ratios, limiting thermal dissipation through respiration and making mature stock highly vulnerable to heat stress⁸. In contrast, neonates enter extra-uterine life with high surface-area-to-mass ratios, thin subcutaneous fat layers, and limited glycogen stores, making them susceptible to lethal hypothermia¹. Consequently, farrowing rooms maintain a dual-climate regime: the sow's ambient space is maintained at 18°C to 20°C (64°F to 68°F), while localized microclimate zones (creep boxes or heated mats) supply 32°C to 35°C (90°F to 95°F) for piglets during their first postnatal week¹.

Mechanical ventilation operates via negative-pressure systems that balance convective heat loss with moisture, carbon dioxide (CO_2), hydrogen sulfide (H_2S), and ammonia (NH_3) removal⁸. Flow rate capacities are measured in cubic feet per minute (CFM) across cold, mild, and hot seasons⁸.

Production Phase	Body Weight / Status	Space Allowance: Slatted Floor (m ² / ft ²)	Space Allowance: Deep Bedding (m ² / ft ²)	Minimum Cold Ventilation (CFM/pig)	Maximum Hot Ventilation (CFM/pig)	Thermal Comfort Zone (°C / °F)
Neonatal Piglet	1.5–6 kg	Micro-zone only	Micro-zone only	Local creep control	Local creep control	32–35°C / 90–95°F ¹
Nursery Weaner	6–30 kg	0.30–0.40 / 3.2–4.3 ¹	0.50–0.60 / 5.4–6.5	2–3 ⁸	30–35 ⁸	24–28°C / 75–82°F ¹
Grower	30–70 kg	0.50–0.65 / 5.4–7.0 ³	0.70–0.85 / 7.5–9.1	5–7 ⁸	70–80 ⁸	18–22°C / 64–72°F ¹
Finisher	70–135 kg	0.65–0.75 / 7.0–8.0 ³	0.85–1.00 / 9.1–10.8 ³	10–12 ⁸	120–150 ⁸	15–20°C / 59–68°F ¹
Gestating Sow	Mature breeding	1.80–2.23 / 19.4–24.0 ⁶	2.23 / 24.0 ²	12–15 ⁸	150–180 ⁸	15–18°C / 59–64°F

Lactating Sow	Mature + Litter	4.0–5.0 / 43.0–53.8	6.0–7.0 / 64.5–75.3	20 ⁹	400–500	18–20°C / 64–68°F ¹
----------------------	-----------------	---------------------	---------------------	-----------------	---------	--------------------------------

Precision Nutritional and Dietary Requirements

Swine possess a monogastric gastrointestinal anatomy reliant on enzymatic breakdown in the stomach and small intestine, supplemented by limited microbial fermentation in the hindgut¹⁰. Modern feed formulation utilizes the Net Energy (NE) system rather than Metabolizable (ME) or Digestible Energy (DE)¹¹. The NE framework accounts for tissue energy retention by subtracting the heat increment associated with digestion and metabolic waste processing, which is particularly elevated during the catabolism of excess crude protein and structural fibers¹⁰. Precision amino acid nutrition is predicated on meeting requirements for Standardized Ileal Digestible (SID) essential amino acids rather than total crude protein¹⁰. Over-supplying crude protein increases urinary nitrogen excretion, degrades air quality through elevated ambient ammonia, and incurs metabolic energy costs for hepatic deamination and renal urea clearance¹². Lysine serves as the universal reference amino acid¹³. Daily requirements reflect the summation of basal endogenous maintenance costs and somatic protein accretion¹⁰:

$$\text{Lysine}_{\text{maintenance}} = 0.036 \times \text{BW}^{0.75} \quad (\text{g SID Lysine/day})$$

[cite: 14]

$$\text{Lysine}_{\text{accretion}} = 0.12 \times \Delta\text{P}_{\text{body}} \quad (\text{g SID Lysine/day})$$

[cite: 14]

In these models, metabolic body size ($\text{BW}^{0.75}$) governs somatic turnover, and daily whole-body protein accretion ($\Delta\text{P}_{\text{body}}$, in grams) is derived by dividing lean carcass tissue gain by 2.55¹⁴. The coefficient 0.12 reflects an average whole-body protein lysine concentration of 7.0% combined with a 58% partial efficiency of digestible lysine utilization¹⁴. Diets are balanced according to the Ideal Protein Concept, maintaining fixed ratios of secondary essential amino acids relative to SID Lysine (100%): SID Threonine at 60–65%, SID Methionine + Cysteine at 55–60%, SID Tryptophan at 18–22%, and SID Valine at 65–70%¹⁰.

Mineral nutrition centers on the balance between calcium (Ca) and phosphorus (P)⁵. Phosphorus requirements are expressed on a Standardized Total Tract Digestible (STTD) basis

rather than as total mineral phosphorus¹³. Because plant phosphorus is largely bound within phytate molecules (myo-inositol hexakisphosphate), endogenous porcine enzymes cannot hydrolyze it efficiently¹³. Diets are formulated with exogenous microbial 6-phytases (often derived from fungal or bacterial strains like *Aspergillus oryzae* or *Escherichia coli*) to liberate orthophosphate¹³. This addition reduces the need for supplemental inorganic dicalcium phosphate and lowers environmental phosphorus excretion¹³.

The STTD Ca:STTD P ratio must be tightly managed between 1.1:1 and 1.3:1⁵. Excessive dietary calcium forms insoluble calcium-phytate complexes in the duodenum, preventing both native and supplemental phytase activity and precipitating secondary phosphorus deficiencies⁵.

Water intake directly determines feed intake and nutrient utilization¹. Delivery infrastructure must ensure appropriate physical heights and flow rates¹⁶. Insufficient flow rates cause frustration, prolong feeder occupancy, and limit feed consumption, while excessive pressure induces water wastage that inflates manure pit volumes¹⁶. Physicochemical water quality requires monitoring: Total Dissolved Solids (TDS) must remain below 1,000 mg/L, nitrates below 100 mg/L, and sulfates below 500 mg/L to avoid osmotic diarrhea in weaners¹.

Production Stage	Body Weight (kg)	Net Energy (Mcal/kg)	Crude Protein (%)	SID Lysine (%)	STTD Phosphorus (%)	Total Calcium (%)	Drinker Flow Rate (L/min)	Water Intake Range (L/day)
Nursery (Phase 1)	6–11	2.50–2.55	20–22 ¹²	1.40–1.55 ¹²	0.45–0.50	0.80–0.85	0.5–1.0 ¹⁶	1.5–3.0
Nursery (Phase 2)	11–25	2.45–2.50	18–20	1.25–1.35	0.38–0.42	0.70–0.75	0.8–1.0 ¹⁶	3.0–5.0
Grower	25–50	2.40–2.45	16–18 ¹²	1.05–1.15 ¹²	0.31–0.34	0.65–0.70	1.0–1.5 ¹⁶	5.0–8.0
Finisher (Early)	50–80	2.40–2.45	14–16 ¹²	0.90–1.00 ¹²	0.26–0.29	0.55–0.60	1.5–2.0 ¹⁶	7.0–10.0

)								
Finisher (Late)	80–135	2.40–2.47	12–14	0.75–0.85	0.22–0.25	0.45–0.50	1.5–2.0 ¹⁶	8.0–12.0
Gestating Sow	Mature	2.15–2.25	11–13 ¹²	0.55–0.65 ¹²	0.24–0.28	0.65–0.75	1.5–2.0 ¹⁶	12.0–18.0
Lactating Sow	Mature + Litter	2.55–2.65	16–18 ¹²	0.95–1.10	0.38–0.42	0.75–0.85	2.0–3.0 ¹⁶	25.0–45.0

Behavioral Ecology, Foraging Dynamics, and Pasture Management

Domestic swine retain the fundamental behavioral repertory of *Sus scrofa*, exhibiting strong drives for rooting, foraging, and exploratory sniffing that occupy 50% to 70% of their active daylight hours¹⁸. The porcine snout is a specialized anatomical organ; the rostral bone (*os rostri*) provides rigid structural support to a cartilaginous disc supplied with tactile corpuscles and motor innervation. This adaptation enables pigs to locate and unearth buried tubers, rhizomes, earthworms, and subterranean fungi.

In intensive barren environments lacking exploratory substrates, this unsatisfied appetitive rooting drive frequently redirects toward pen hardware or other pigs, resulting in stereotypic snout rubbing, sham chewing, bar biting, or systemic tail biting¹⁸. Supplying manipulable, destructible rooting substrates—such as long-stemmed straw, pine shavings, or compost—is critical for satisfying this instinct and reducing aggression¹⁸.

Pastoral production systems leverage this behavioral drive by incorporating pasture foraging into the nutritional regimen¹⁸. The agronomic sward is typically established using mixtures of deep-rooted, drought-tolerant legumes (such as alfalfa, *Medicago sativa*, or white clover, *Trifolium repens*) and sod-forming perennial grasses (including perennial ryegrass, *Lolium perenne*, and orchardgrass, *Dactylis glomerata*)¹⁹. Mature gestating sows possess the functional large-intestinal volume and microbial diversity to digest fibrous biomass, obtaining 20% to 40% of their maintenance energy from pasture fermentation.

In contrast, growing and finishing pigs have smaller gastrointestinal capacities, rapid transit rates, and high nutrient requirements for somatic protein deposition, limiting their nutritional contribution from swards to between 5% and 15% of daily dry matter intake¹⁸. Consequently, growing pigs maintained on pasture require daily supplementation with balanced grain concentrates to prevent deficits in net energy and essential amino acids¹⁸.

Unmanaged pastoral swine often cause significant ecological disturbance¹⁸. Concentrated

rooting tears the root zone of forages, turning vegetation into bare, compacted soil within days¹⁸. This bare ground accelerates rainfall runoff, soil erosion, and localized leaching of urinary nitrogen and fecal phosphorus into surrounding watersheds. Paddock management strategies rely on rotational grazing or strip stocking. Swards are subdivided into discrete paddocks, grazing cohorts at 10 to 15 gestating sows per hectare (4 to 6 per acre) or 25 to 30 grow-finish pigs per hectare. Paddock transitions occur when vegetation height drops below 5 to 8 cm, followed by a minimum rest period of 4 to 8 weeks to allow vegetative root recovery and sward re-establishment.

A traditional intervention to suppress rooting damage was snout ringing, involving the insertion of wire rings through the dorsal rim of the snout cartilage to elicit discomfort whenever downward rooting pressure was applied. However, ethological evidence demonstrates that ringing frustrates innate exploratory behaviors, causing negative affective states and impairing welfare¹⁸. Organic production standards, notably European Union Regulation (EU) 2018/848, largely prohibit snout ringing¹⁸. Producers operating under strict welfare schemes instead use rotational grazing schedules, establish sacrifice paddocks during wet seasons, integrate root-resilient grass varieties, and offer supplemental roughage (such as silage or hay) to mitigate ground disturbance¹⁸.

Pasture exposure also re-establishes parasite transmission cycles that are largely disrupted in concrete indoor housing²⁰. Rearing swine on open ground exposes herds to several major helminths:

- **Ascaris suum (Large Roundworm):** Thick-coated eggs shed in swine feces can persist in pastoral soils for 5 to 10 years¹⁹. Ingested embryonated eggs hatch in the duodenum, after which larvae penetrate the intestinal mucosa, enter the mesenteric venules, and migrate through the hepatic parenchyma to create fibrous interstitial hepatitis, visible as "milk spot" lesions²². Larvae then migrate to pulmonary capillaries, break into alveolar spaces—triggering verminous pneumonia and coughing—and are carried up the bronchial mucociliary escalator to be swallowed, completing maturation to 30 cm adult worms in the jejunum²².
- **Metastrongylus apri, M. pudendotectus, and M. salmi (Lungworms):** These parasites depend on common earthworms (such as *Lumbricus terrestris* and *Eisenia fetida*) as obligatory intermediate hosts²¹. Swine ingest infected earthworms while rooting through topsoil²⁴. Third-stage (L_3) larvae penetrate the host's intestinal wall, migrate through mesenteric lymph nodes, pass through the right cardiac chambers, and settle in the bronchioles²¹. Clinical manifestations include chronic verminous bronchitis, dyspnea, and paroxysmal coughing fits termed "thumping," which predispose growing pigs to secondary bacterial infections from agents like *Pasteurella multocida* and *Mycoplasma hyopneumoniae*²¹.
- **Trichuris suis (Whipworm):** Larvae burrow into cecal and colonic mucosae, inducing chronic catarrhal, mucohemorrhagic typhlocolitis that mimics or exacerbates swine dysentery (*Brachyspira hyodysenteriae*)¹⁹.

Parasite control in pasture operations requires combining herd-level chemical anthelmintic

therapy with environmental pasture breaks²⁰. Broad-spectrum anthelmintics—primarily benzimidazoles (fenbendazole, flubendazole) or macrocyclic lactones (ivermectin, doramectin)—should be administered sequentially during movements to fresh ground to limit paddock egg shedding¹⁹.

Agro-Industrial Byproduct Valorization and Food Waste Upcycling

Feed comprises between 60% and 70% of variable operating costs in domestic swine enterprises¹³. Incorporating agricultural processing residues, distilling byproducts, and municipal food waste offers an avenue to lower feed expenses and divert organic materials from landfills, reducing greenhouse gas emissions²⁹. However, formulating these non-traditional resources requires accounting for high moisture variability, fiber composition, and dietary fatty acid profiles³².

Distillers Dried Grains with Solubles (DDGS), an abundant coproduct of the dry-grind ethanol industry, contains 26% to 30% crude protein, high digestible phosphorus, and concentrated residual corn oil³⁵. The lipid fraction of DDGS is high in polyunsaturated fatty acids,

predominantly linoleic acid ($C_{18:2\ n-6}$)³². Because monogastric animals lack the ruminal biohydrogenation mechanisms of cattle and sheep, ingested dietary fatty acids are deposited directly into porcine adipocytes with minimal structural alteration³².

Diets containing high levels of DDGS—or other polyunsaturated lipid sources—can compromise adipose tissue quality³². Elevating dietary linoleic acid intake above 1.6% to 1.8% of dry matter causes soft, oily belly fat that reduces commercial bacon slicing yields and accelerates oxidative rancidity³². Packers quantify this carcass quality metric using the Iodine Value (IV), which measures the degree of fatty acid unsaturation:

$$\text{Dietary IVP} = \text{Dietary Fat (\%)} \times \left(\frac{\text{Iodine Value of the Lipid Fraction}}{100} \right)$$

$$\text{Carcass Adipose IV} \approx 52.4 + (0.31 \times \text{Daily Linoleic Acid Intake})$$

Commercial abattoirs typically require carcass fat IV levels below 70 to 74 g/100 g³². To prevent excessive unsaturation, nutritionists restrict dietary DDGS inclusion to 15% to 20% during early-grower periods and often remove it entirely 30 to 45 days prior to slaughter, switching finishers to saturated lipid sources such as beef tallow³⁵.

Other agricultural processing streams present distinct physiological opportunities and

nutritional constraints:

- **Liquid Whey:** A dairy processing byproduct that supplies high-biological-value whey proteins (lactalbumin and lactoglobulin) alongside digestible lactose³³. However, its mineral ash content—particularly sodium and potassium—is elevated, requiring unrestricted freshwater access to avoid hyperosmotic salt toxicity³³.
- **Brewery Spent Grains (BSG):** High in neutral detergent fiber ($> 45\%$ NDF) and low in net energy, BSG is ill-suited for young pigs with limited gut volume³³. Conversely, it serves as an effective bulk-filling ingredient for gestating sows, promoting satiety and reducing stereotypic hunger behaviors during feed restriction³³.
- **Bakery Meal and Confectionery Residuals:** Consisting of discarded bread, doughs, cookies, and potato chips, this stream provides easily digestible, heat-gelatinized starches, free simple sugars, and supplemental fats that yield high net energy (> 2.7 Mcal/kg NE)³⁰. These qualities allow it to directly substitute for ground corn, provided sodium levels are monitored to avoid electrolyte imbalances⁴¹.

High-moisture agricultural discards spoil rapidly via wild yeasts and enteropathogenic bacteria, requiring stabilization before feeding³⁴. Solid-state ensiling utilizes homofermentative lactic acid bacteria (such as *Lactobacillus plantarum*) to convert water-soluble carbohydrates into lactic acid under anaerobic conditions³³. This reduces substrate pH below 4.2, suppressing the growth of pathogens like *Salmonella*, *Listeria monocytogenes*, and *Clostridium* species³⁴. In Japan and South Korea, food waste utilization has been formalized through centralized "Eco-Feed" systems³⁰. Japanese processors collect food waste from retailers, hotels, and food factories, process it using thermal dehydration or automated liquid fermentation, and distribute it to pig farms³³. Liquid Eco-Feed systems mince food wastes, blend them with supplementary amylase to reduce slurry viscosity, heat-treat the mixture, and inoculate it with characterized lactic acid cultures to stabilize storage⁴⁴. The finished liquid feed is then delivered directly through computer-controlled piping networks into farm feeding troughs⁴⁴.

Feeding animal-derived catering and food wastes carries disease transmission risks, particularly for high-consequence viral pathogens such as African Swine Fever (ASF), Classical Swine Fever (CSF), Foot-and-Mouth Disease (FMD), and vesicular diseases³³. Consequently, regulatory approaches vary substantially across international jurisdictions:

- **United States:** Regulated under the federal *Swine Health Protection Act* (7 U.S.C. 3801 et seq.) and 9 CFR Part 166⁴¹. The law prohibits feeding garbage—defined as any food waste containing meat, animal tissues, or refuse that contacted meat—unless processed at a licensed facility⁴¹. Processing requires heating the garbage throughout to a continuous rolling boil (100°C / 212°F at sea level) for at least 30 minutes with mechanical agitation⁴¹. Dry industrial products (like bakery waste, candy waste, and domestic dairy) are exempt from these boiling rules, provided they have not contacted animal meat tissues⁴¹. Several individual US states maintain bans that prohibit garbage feeding entirely within their borders⁴⁸.
- **European Union:** Governed by Regulation (EC) No 1069/2009 and Commission

Regulation (EU) No 142/2011⁴⁵. Following the 2001 Foot-and-Mouth Disease outbreak in the United Kingdom—which was traced to illegal feeding of untreated catering waste—the EU established a total ban under Article 11(1)(b) of Regulation 1069/2009⁴⁵. The feeding of catering waste, restaurant plate scrapings, domestic swill, and intra-species animal proteins to farm animals is prohibited, irrespective of thermal processing⁴⁵. Animal-derived byproducts may only be processed into feed ingredients via Category 3 rendering facilities with verified pathogen inactivation⁴⁹.

- **Japan:** Governed by the *Food Recycling Law* and the *Feed Safety Law*³³. The regulatory framework permits and subsidizes the processing of food wastes into Eco-Feed³³. To address ASF and CSF risks, raw materials containing meat must undergo certified thermal processing—at a minimum of 70°C for 30 minutes, or 80°C for 3 minutes—prior to drying or liquid fermentation, with the finished feed subject to biosecurity tracing³³.

Regulatory Domain	Governing Statute / Code	Swill / Food Waste Permissibility	Mandatory Thermal Regimes	Industrial Exemptions
United States	9 CFR Part 166; Swine Health Protection Act ⁴¹	Permitted only with state/federal facility licensure; prohibited in select states ⁴¹ .	Continuous boiling at 100°C (212°F) for 30 minutes under mechanical agitation ⁴¹ .	Industrial dry bakery waste, candy discards, eggs, domestic dairy ⁴¹ .
European Union	Regulation (EC) No 1069/2009; Reg (EU) No 142/2011 ⁴⁵	Strict Total Prohibition on feeding catering waste or swill to food animals ⁴⁵ .	None permitted for catering swill; industrial Category 3 rendering only ⁴⁵ .	Pure vegetable residues from commercial processing without animal contact ⁴⁹ .
Japan	Food Recycling Law; Feed Safety Regulations ³³	Permitted and Encouraged via formal Eco-Feed manufacturing certification ³³ .	Heat-treatment at 70°C for 30 min, or 80°C for 3 min (stricter standards for meat waste) ³³ .	Uncontaminated single-stream plant processing byproducts ³³ .

Epidemiology, Clinical Pathology, and Biosecurity

Eradication of Trichinosis

Trichinosis (trichinellosis) is a parasitic zoonosis caused by nematodes of the genus *Trichinella*⁵⁵. Within domestic and wild swine populations, the primary etiologic agents are *Trichinella spiralis*—an encapsulated species well adapted to domestic pigs—and sylvatic species, including *Trichinella britovi*, *Trichinella pseudospiralis* (non-encapsulated), and *Trichinella nativa* (freeze-tolerant)⁵⁶.

The life cycle is auto-heteroxenous, requiring a single individual host to act sequentially as both the definitive and intermediate host⁵⁷. Swine become infected by consuming muscle tissue

containing viable encysted first-stage (L_1) larvae⁵⁶. Following ingestion, gastric hydrochloric acid and pepsins dissolve the collagen capsule, releasing free larvae into the lumen⁵⁷.

Larvae penetrate the columnar epithelial lining of the small intestine (jejunum and upper ileum) and undergo four molts over 30 to 48 hours to reach sexual maturity⁵⁷. Following copulation, adult females release newborn larvae (approximately 100 μm in length) into the intestinal lymphatic and venous systems between days 5 and 7 post-infection⁵⁷. Migrating larvae disseminate systemically through the portal veins, cardiac chambers, and peripheral arterial vasculature to invade striated muscle fibers²⁴.

Once inside a skeletal muscle fiber, the larva alters host cell gene expression, inducing the loss of myofibrillar architecture, stimulating local angiogenesis, and triggering collagen production⁵⁷. This modified host cell is termed a *nurse cell*⁵⁷. The nurse cell-larva complex can remain viable within the host for years, completing transmission only when muscle tissue is consumed by another carnivorous or omnivorous host⁵⁶.

Porcine infections are typically subclinical, with clinical disease (pyrexia, periorbital edema, and stiff gait) occurring primarily during experimental challenges with high larval doses. Human consumption of undercooked, larva-bearing pork, however, can result in severe gastrointestinal illness followed by intense myalgia, myocarditis, facial edema, and life-threatening neurotrichinosis.

Swine infection on farms involves three primary transmission routes:

- **Ingestion of Raw or Inadequately Cooked Waste:** The historical and predominant domestic infection pathway, caused by feeding swill, abattoir scraps, or hunting offal containing viable larvae⁴¹.
- **Peridomestic Rodent Consumption:** Rats (*Rattus norvegicus* and *Rattus rattus*) serve as natural bridge hosts⁵⁹. Pigs readily capture and consume rats within feed passages; conversely, rats acquire infection by feeding on deceased swine or practicing cannibalism, maintaining a self-sustaining cycle within infested housing⁵⁹.
- **Tail-Biting and Herd Cannibalism:** Severe behavioral vices with tissue necrosis can transmit larvae between herd mates if infectious muscle tissue or microfilarial blood is ingested.
- **Pasture and Sylvatic Exposure:** Outdoor access exposes swine to infected native

wildlife—including red foxes (*Vulpes vulpes*), wild boars (*Sus scrofa*), and raccoons (*Procyon lotor*)—either through carcass scavenging or foraging on soils containing infected tissues²¹.

Preventing *Trichinella* transmission requires verifying on-farm operational controls⁵⁵. Under international frameworks (including EU Regulation 2015/1375 and USDA FSIS guidance), production facilities can achieve official recognition as **Controlled Housing Conditions** by maintaining strict biosecurity standards⁵⁵:

- **Perimeter Enclosures and Physical Construction:** All barns must be fully enclosed with intact siding, sealed entryways, concrete aprons, and mesh over ventilation louvers to prevent the entry of rodents, stray domestic pets, and wildlife⁴⁷.
- **Targeted Rodent Abatement Plans:** Holdings must implement an active pest management program featuring mapped, tamper-proof exterior bait stations containing anticoagulant rodenticides or mechanical kill-traps, inspected and replenished on a regular schedule⁴⁷.
- **Strict Feed Controls:** Herds must be maintained on grain- or certified concentrate-based rations, with an absolute prohibition on feeding swill, scrap meat, or unprocessed kitchen waste⁵⁵.
- **Carcass Handling:** Pig mortalities must be removed from pens immediately to prevent cannibalism, stored in sealed refrigerated storage areas, and disposed of via licensed rendering, on-site incineration, or deep-burial methods⁴⁷.

Under European Union controls, herds reared within officially verified Controlled Housing Conditions are subject to risk-based surveillance, requiring the post-mortem digestion testing of all breeding sows and boars, alongside at least 10% of market finishers⁵⁵. Carcasses originating from facilities without certified controlled housing must undergo 100% individual testing at slaughter⁵⁵.

The post-mortem surveillance standard is the **Magnetic Stirrer Method (MSM) for Pooled Sample Digestion**, codified under ISO 18743:2015 and Commission Implementing Regulation (EU) 2015/1375⁶¹. Compressive trichinostomy (flattening tissue between glass plates) has been

discontinued as a routine diagnostic standard because its detection threshold ($> 3\text{--}5 \text{ lpg}$) fails to reliably identify low-level infestations capable of causing human disease⁶¹.

The magnetic stirrer protocol is structured around defined sequential stages:

- **Tissue Collection:** Muscle samples are excised from primary anatomical predilection sites. The preferred site in domestic swine is the muscular crus of the diaphragm at the transition to the central tendon; secondary sites include the tongue and masseter muscles⁵⁷. Samples must be carefully trimmed of superficial fat, tendons, and fascia⁶⁵.
- **Enzymatic Digestion Preparation:** In a 3-liter glass beaker, 2.0 liters of tap water is heated to 46°C to 48°C and supplemented with $16 \pm 0.5 \text{ mL}$ of 25% hydrochloric acid (or equivalent volume adjusted for 37% stock to achieve $\sim 1\% \text{ HCl}$)⁶². Next, $10 \pm 0.2 \text{ g}$ of certified high-activity pepsin powder (1:10,000 NF activity) is stirred into

the warm acidic solution⁶².

- **Sample Digestion:** A pool of up to 100 individual tissue samples (typically 1 g per animal, or 3 to 5 g for breeding animals) is homogenized in a laboratory blender⁶³. The blended tissue is introduced into the acidic digestion fluid⁶². The beaker is covered with aluminum foil, set atop a heated magnetic stirrer calibrated to maintain fluid temperature between 44°C and 46°C, and agitated vigorously to generate a deep central vortex without splashing⁶². Stirring continues for approximately 30 minutes until visible myofibers dissolve⁶². Digestion is considered complete if less than 5% of the initial sample mass remains undigested⁶³.
- **Filtration and Initial Sedimentation:** The digest is poured through a 180 µm stainless steel wire mesh sieve into a glass separatory funnel to capture residual connective tissue⁶². The fluid is allowed to settle undisturbed for 30 minutes, allowing free larvae to descend by gravity to the stopcock⁶².
- **Secondary Clarification:** The technician opens the stopcock to draw off a 40 mL aliquot into a graduated cylinder, which is held undisturbed for 10 minutes⁶². The upper 30 mL of clear supernatant is carefully aspirated, leaving 10 mL of sediment⁶².
- **Microscopic Examination:** The 10 mL sediment is transferred to a gridded larval counting dish, rinsed with tap water, and examined under a stereomicroscope or trichinoscope at 15× to 20× magnification⁶². Any suspicious nematode structures are verified at 60× to 100× magnification to evaluate characteristic morphologic features, such as larval coiling, esophageal stichosome structure, and a rounded caudal terminus⁶³.

Untested pork carcasses or product lines derived from non-controlled production facilities must undergo post-harvest physical treatments to ensure the complete inactivation of *Trichinella* larvae prior to commercial release⁵⁸:

- **Thermal Inactivation (Cooking):** Pork products must reach an internal core temperature of at least 62.2°C (144°F) instantaneously, 60.0°C (140°F) for 1 minute, or 55.0°C (131°F) for 6 minutes, which denatures essential parasitic structural proteins⁶⁸.
- **Ionizing Irradiation:** Controlled industrial exposure to gamma radiation (via Cobalt-60 or Cesium-137) or electron beam accelerators with an absorbed dose of 0.3 to 0.75 kGy renders encysted larvae sexually sterile and non-infective⁵⁸.
- **Commercial Freezing:** Enforcing continuous sub-zero holding regimes based on cut thickness, categorized into Group 1 (meat pieces ≤ 15 cm thick) and Group 2 (pieces 15 to 50 cm thick or packed in large bulk containers)⁵⁸:

Freezer Temperature (°C / °F)	Group 1: Product Thickness ≤ 15 cm (Days Required)	Group 2: Product Thickness 15–50 cm (Days Required)

−15.0° C / +5.0° F	20 ⁵⁸	30 ⁵⁸
−23.3° C / −10.0° F	10 ⁵⁸	20 ⁵⁸
−28.9° C / −20.0° F	6 ⁵⁸	12 ⁵⁸

Commercial freezing parameters apply exclusively to pork derived from domestic swine infected with *Trichinella spiralis*⁵⁸. Freezing does not reliably inactivate cold-tolerant sylvatic genotypes, particularly *Trichinella nativa* and *Trichinella* genotype T-6, which are adapted to arctic and sub-arctic carnivores⁵⁶. These freeze-resistant larvae can survive extended commercial freezing schedules, meaning pork or wild game exposed to sylvatic cycles must be cooked thoroughly or verified by digestive testing prior to human consumption⁵⁶.

Synthesis and Strategic Industry Trajectory

Balancing domestic swine production requires reconciling animal welfare expectations, precision feed formulation, and biosecurity protocols. Modern facility design increasingly shifts toward loose group housing to satisfy behavioral welfare standards, such as California Proposition 12⁶. However, group pens require precise space allocations and controlled environmental ventilation to limit competitive aggression and respiratory pathogen accumulation¹.

Extending rearing into pastoral systems satisfies natural rooting and foraging behaviors, yet it creates distinct soil conservation challenges and re-establishes parasite cycles—including *Ascaris suum* and *Metastrongylus apri*—that require rotational pasture strategies and strategic deworming programs¹⁸.

In nutrition, formulating to Net Energy and Standardized Ileal Digestible amino acid patterns permits the inclusion of alternative agro-industrial byproducts, such as corn DDGS, bakery discards, and processed food wastes¹¹. Nonetheless, incorporating high-fat coproducts requires continuous monitoring of polyunsaturated fatty acid intakes and carcass Iodine Values to avoid uncohesive fat defects in market bellies³².

The utilization of food wastes remains governed by biosecurity trade-offs. While regions like Japan have established safe circular processing through thermal Eco-Feed systems, risks associated with African Swine Fever, Classical Swine Fever, and Foot-and-Mouth Disease justify total swill bans within the European Union and licensed cooking requirements in the United States³³.

Finally, controlling zoonotic pathogens like *Trichinella spiralis* relies on multiple verification points. The foundation rests on pre-harvest biosecurity: rodent exclusion, deadstock removal, and preventing access to unboiled food wastes⁴⁷. At processing, risk-based surveillance using the reference Magnetic Stirrer Method for pooled sample digestion ensures individual and herd safety⁵⁵. Integrating housing design, digestive biochemistry, behavioral management, and diagnostic monitoring forms an interconnected system that supports sustainable production

while protecting animal health and the human food chain¹.

Works cited

1. Pig Farming: Breeding, Farrowing, Nursery, Grow-Finish, Nutrition, <https://www.zubairkhalid.com/knowledge/animal-farming/swine/pig-farming-breeding-farrowing-nursery-grow-finish-nutrition-and-biosecurity>
2. Sow Housing Options for Gestation - Pork Information Gateway, <https://porkgateway.org/resource/sow-housing-options-for-gestation/>
3. Pig Housing and Farm Infrastructure: Barn Types & Space, <https://usapigs.com/pig-housing-farm-infrastructure/>
4. evaluation of dietary alterations that have potential to affect feed, https://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1646&context=gradschool_theses
5. Nutritional Requirements of Pigs - Management and Nutrition, <https://www.merckvetmanual.com/management-and-nutrition/nutrition-pigs/nutritional-requirements-of-pigs>
6. Prop 12: Gestation-sow Housing Considerations and Implications, <https://farmweld.com/prop-12-gestation-sow-housing-considerations-and-implications/>
7. Five burning questions on Prop 12 compliance - National Hog Farmer, <https://www.nationalhogfarmer.com/livestock-management/five-burning-questions-on-prop-12-compliance>
8. Minimizing Heat Stress in Pigs - Form-A-Feed, <https://formafeed.com/minimizing-heat-stress-pigs/>
9. Ventilation System Design for Pig Farrowing Rooms | Dr. Zubair Khalid, <https://www.zubairkhalid.com/knowledge/animal-farming/farm-management/ventilation-system-design-for-pig-farrowing-rooms>
10. Understanding the Nutrient Recommendations in the National, <https://porkgateway.org/resource/understanding-the-nutrient-recommendations-in-the-national-swine-nutrition-guide/>
11. Determination of optimal energy system and level for growing pigs, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11222115/>
12. Nutrient Requirements Of Swine Eleventh Revised Edition 2012, https://enflomaplata.uep.edu.py/ProductPdf/TFDpEZ/705655/nutrient_requirements-of_swine_eleventh_revised_edition_2012.pdf
13. National Swine Nutrition Guide tables on nutrient recommendations, <https://porkgateway.org/resource/national-swine-nutrition-guide-tables-on-nutrient-recommendations-ingredient-composition-and-use-rates/>
14. Nutrient requirements of swine - CABI Digital Library, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/19990404682>
15. Nutritional Value of Ingredients - K-State Animal Science, <https://www.asi.k-state.edu/extension/swine/swinenutritionguide/pdf/KSU%20Nutritional%20Value%20of%20Ingredients%20fact%20sheet.pdf>
16. Pig Water Delivery Systems: Drinkers, Flow Rates, and Quality,

- <https://www.zubairkhalid.com/knowledge/animal-farming/swine/pig-water-delivery-systems-drinkers-flow-rates-quality>
17. Free Swine Drinker & Feeder Stocking Calculator - ConductScience, <https://conductscience.com/tools/swine-drinker-feeder-stocking-calculator>
 18. (PDF) Welfare and environmental impact of organic pig production, https://www.academia.edu/99325664/Welfare_and_environmental_impact_of_organic_pig_production
 19. Broad-Spectrum Control of Worms | The Pig Site, <https://www.thepigsite.com/articles/broadspectrum-control-of-worms>
 20. Deworming recommendations for pasture-raised pigs in Michigan, <https://www.canr.msu.edu/news/deworming-recommendations-pigs-in-michigan>
 21. Lungworms (*Metastrongylus* spp.) demonstrated in domestic pigs, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9006407/>
 22. Anthelmintics in Pigs – RUMA, <https://www.ruma.org.uk/anthelminitics-in-pigs/>
 23. Worms - NADIS, <https://www.nadis.org.uk/disease-a-z/pigs/worms/>
 24. Understanding Worms in Swine: A Practical Guide | Elanco, <https://farmanimal.elanco.com/au/insights-centre/understanding-worms-in-swine>
 25. Protect your pigs from verminous pneumonia - Pork, <https://www.canr.msu.edu/news/protect-your-pigs-from-verminous-pneumonia>
 26. (PDF) An identification key for the five most common species of, https://www.researchgate.net/publication/264248313_An_identification_key_for_the_five_most_common_species_of_Metastrongylus
 27. Lungworm Infection in Animals - Respiratory System, <https://www.msdevetmanual.com/respiratory-system/lungworm-infection/lungworm-infection-in-animals>
 28. Lungworm in pigs | Langford Vets, <https://www.langfordvets.co.uk/animal-insights/pigs/lungworm-in-pigs/>
 29. The Piggy Solution: Harnessing Food Waste for Sustainable Hog, https://www.researchgate.net/publication/393752362_The_Piggy_Solution_Harnessing_Food_Waste_for_Sustainable_Hog_Farming
 30. Advancing resource mediated food waste conversion, <https://notulaeobotanicae.ro/index.php/nbha/article/download/14683/10056/73601>
 31. “What a Waste”—Can We Improve Sustainability of Food Animal, <https://www.mdpi.com/2071-1050/12/17/7071>
 32. Pork Fat Quality, Iodine Values and Feeding DDGS | The Pig Site, <https://www.thepigsite.com/articles/pork-fat-quality-iodine-values-and-feeding-ddgs>
 33. Eco-Feeds in Japan | FFTC Agricultural Policy Platform (FFTC-AP), <https://ap.ffmpeg.org.tw/article/2794>
 34. Ensiling of food waste for subsequent inclusion in ruminant diets, <https://cdnsiencepub.com/doi/10.1139/cjas-2025-0021>
 35. The effects of feeding diets containing corn distillers dried grains, <https://pubmed.ncbi.nlm.nih.gov/19966154/>
 36. (PDF) Pork fat quality of pigs fed distillers dried grains with solubles, https://www.researchgate.net/publication/293191027_Pork_fat_quality_of_pigs_fe

- [d_distillers_dried_grains_with_solubles_with_variable_oil_content_and_evaluation_of_iodine_value_prediction_equations1](#)
37. Dietary Linoleic Acid Content Can Predict Carcass Fat Quality,
<https://www.thepigsite.com/articles/dietary-linoleic-acid-content-can-predict-carcass-fat-quality>
 38. Solving the puzzle: How to meet pork carcass iodine value standard,
<https://www.nationalhogfarmer.com/hog-health/solving-the-puzzle-how-to-meet-your-pork-carcass-iodine-value-standard>
 39. Assessment of the effects of diets containing DDGS with,
<https://porkcheckoff.org/research/assessment-of-the-effects-of-diets-containing-ddgs-with-supplemental-tallow-on-fat-digestibility-growth-performance-carcass-and-fat-quality-in-growing-finishing-pigs/>
 40. Effects of dried distillers grains with solubles on carcass fat quality of,
<https://academic.oup.com/jas/article-pdf/88/11/3666/23653722/3666.pdf>
 41. Rules and Regulations - USDA-Aphis,
https://www.aphis.usda.gov/sites/default/files/interim_rule_pro-products.pdf
 42. Circular Eco-feed chain for responsible pork consumption and,
<https://edepot.wur.nl/515344>
 43. (PDF) Production Efficiency of Animal-Feed Obtained From Food,
https://www.researchgate.net/publication/356116213_Production_Efficiency_of_Animal-Feed_Obtained_From_Food_Waste_in_Japan
 44. Surplus food as animal feed - REFRESH,
https://eu-refresh.org/sites/default/files/06_Karen%20Luyckx%2C%20Jennifer%20David%2C%20Tomoyuki%20Kawashima_animal%20feed.pdf
 45. Animal Feedstuffs – Tuesday, 25 Nov 2014 - Oireachtas.ie,
<https://www.oireachtas.ie/en/debates/question/2014-11-25/222/>
 46. 9 CFR § 166.1 - Definitions in alphabetical order. - Law.Cornell.Edu,
<https://www.law.cornell.edu/cfr/text/9/166.1>
 47. 9 CFR Part 166 -- Swine Health Protection - eCFR,
<https://www.ecfr.gov/current/title-9/chapter-I/subchapter-L/part-166>
 48. PART 166—SWINE HEALTH PROTECTION - Maine.gov,
https://www.maine.gov/dacf/ahw/animal_health/documents/swine-health-protection-on-federal-laws-9-cfr-part-166-up-to-date-as-of-4-14-2026.pdf
 49. Animal by-products not intended for human consumption | EUR-Lex,
<https://eur-lex.europa.eu/EN/legal-content/summary/animal-by-products-not-intended-for-human-consumption.html>
 50. Animal By-Products Regulations - Wikipedia,
https://en.wikipedia.org/wiki/Animal_By-Products_Regulations
 51. Feeding food waste to pigs can reduce land use of EU pork production,
<https://www.cabidigitallibrary.org/do/10.5555/collection-news-24746/full/>
 52. Factsheet - 32003D0328 | European Free Trade Association - EFTA.int,
<https://www.efta.int/eea-lex/32003d0328>
 53. GLOBAL FOOD WASTE MANAGEMENT: - World Biogas Association,
<https://www.worldbiogasassociation.org/wp-content/uploads/2018/05/Global-Food-Waste-Management-Full-report-pdf.pdf>

54. (PDF) PS2 - (Novel) ingredients and processing PS2-138 Efficacy of,
https://www.researchgate.net/publication/393103925_PS2_-_Novel_ingredients_and_processing_PS2-138_Efficacy_of_Different_Form_of_Plantain_Plantago_Lanceolata_L_on_Bone_Mineralization_and_Physiochemical_Characteristics_of_Broiler_Meat
55. B COMMISSION IMPLEMENTING REGULATION (EU) 2015/1375 of,
<https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:02015R1375-20250409>
56. fmd with viaa test incl. - WOAHA,
<https://www.woah.org/app/uploads/2021/03/3-01-20-trichinellosis.pdf>
57. A review of testing and assurance methods for Trichinella,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8379475/>
58. Recommendations on post-harvest control of Trichinella in food,
https://www.researchgate.net/publication/331308569_International_Commission_on_Trichinellosis_Recommendations_on_post-harvest_control_of_Trichinella_in_food_animals
59. Occurrence of Trichinella spp. in rats on pig farms,
<https://www.aaem.pl/Occurrence-of-Trichinella-spp-in-rats-on-pig-farms,99555,0,2.html>
60. Commission Implementing Regulation (EU) 2015/1375 of 10 August,
<https://www.legislation.gov.uk/eur/2015/1375/contents>
61. Gelatin medium for preserving of Trichinella spp. for quality control,
<https://repozytorium.piwet.pulawy.pl/bitstream/handle/123456789/580/Gelatin%20medium%20for.pdf?sequence=2&isAllowed=y>
62. Magnetic Stirrer Method for the Detection of Trichinella Larvae in,
https://www.researchgate.net/publication/314223979_Magnetic_Stirrer_Method_for_the_Detection_of_Trichinella_Larvae_in_Muscle_Samples
63. Trichinella - Legislation.gov.uk,
<https://www.legislation.gov.uk/eur/2015/1375/annexes/data.xht?view=snippet&wrape=true>
64. B COMMISSION REGULATION (EC) No 2075/2005 of 5 ... - EUR-Lex,
<https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CONSLEG:2005R2075:20141111:EN:PDF>
65. Magnetic Stirrer Method for the Detection of Trichinella Larvae in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC5409328/>
66. Magnetrührers Verfahren zum Nachweis von Trichinella Larvae in,
<https://www.jove.com/de/t/55354/magnetic-stirrer-method-for-detection-trichinella-larvae-muscle>
67. Trichinosis - Control of,
[https://www1.agric.gov.ab.ca/\\$Department/deptdocs.nsf/all/rsb12821/\\$FILE/tipm_chapter09.pdf](https://www1.agric.gov.ab.ca/$Department/deptdocs.nsf/all/rsb12821/$FILE/tipm_chapter09.pdf)
68. REFRESH D6.7 Animal Feed Supplementary materials,
<https://eu-refresh.org/sites/default/files/REFRESH%20D6.7%20Animal%20Feed%20Supplementary%20materials.pdf>
69. (PDF) International Commission on Trichinellosis: recommendations,

https://www.researchgate.net/publication/12227797_International_Commission_on_Trichinellosis_recommendations_on_methods_for_the_control_of

70. Curing and Smoking Meat Techniques | Curing (Food Preservation),
<https://www.scribd.com/document/81528346/Curing-and-Smoking-Meats-for-Home-Food-Preservation>
71. Control recommendations for the inactivation of *Trichinella spiralis* in,
<https://inspection.canada.ca/en/preventive-controls/meat/trichinella-spiralis>