

Swine Abattoir Operations: Comprehensive Protocols for Skin Preservation, Safe Evisceration, and Carcass Splitting

The modern commercial processing of domestic swine (*Sus scrofa domesticus*) requires the meticulous synchronization of biological management, thermodynamic interventions, and mechanical engineering. Unlike bovine or ovine slaughter, where the hide is entirely removed prior to evisceration, the economic value of numerous pork cuts—including bone-in bellies, crackling, and cured hams—depends upon the preservation of the skin and its underlying subcutaneous fat layer¹. Achieving this requires specialized thermal scalding to release hair follicles, abrasive mechanical scraping and singeing to clean the epidermis, safe evisceration to prevent enteric pathogen contamination, and precise skeletal splitting to facilitate chilling and grading. This report provides an exhaustive, highly technical analysis of the physiological, mechanical, and microbiological mechanisms governing the complete swine slaughter cascade.

Pre-Slaughter Husbandry, Nutrition, and Lairage Dynamics

The preservation of pork skin and fat quality, as well as the physiological behavior of the carcass during slaughter, begins well before the animal reaches the abattoir. Swine rearing systems, precision nutrition, and pre-slaughter handling directly dictate the thermodynamic behavior of the fat during scalding and the biochemical decline of muscle pH post-mortem⁴.

Nutritional Influences on Skin and Fat Integrity

Swine possess a monogastric gastrointestinal anatomy that relies on enzymatic breakdown in the stomach and small intestine, meaning that ingested dietary fatty acids are deposited directly into porcine adipocytes with minimal structural alteration⁴. Modern precision nutrition utilizes the Net Energy (NE) system, which accounts for tissue energy retention by subtracting the heat increment associated with digestion⁴. Daily requirements are modeled using the Ideal Protein Concept, prioritizing Standardized Ileal Digestible (SID) amino acids, with lysine serving as the primary reference⁴.

The inclusion of agro-industrial byproducts in swine diets directly impacts the integrity of the subcutaneous fat and the skin. Distillers Dried Grains with Solubles (DDGS), a coproduct of the ethanol industry, contains a high lipid fraction rich in polyunsaturated fatty acids, predominantly linoleic acid ($C_{18:2\ n-6}$)⁴. Diets containing excessive polyunsaturated lipids produce soft, oily belly fat that accelerates oxidative rancidity, reduces commercial slicing yields, and compromises the structural integrity of the skin-fat matrix during high-temperature singeing⁴. Packers quantify this carcass quality metric using the Iodine Value (IV), which measures the

degree of fatty acid unsaturation. To maintain the structural rigidity of the skin and fat, commercial abattoirs require carcass fat IV levels to remain below 70 to $74 \text{ g}/100\text{g}$, often necessitating the removal of DDGS from finisher diets 30 to 45 days prior to slaughter⁴.

Production Phase	Body Weight (kg)	Net Energy (Mcal/kg)	Crude Protein (%)	SID Lysine (%)	STTD Phosphorus (%)	Total Calcium (%)
Nursery (Phase 1)	6–11	2.50–2.55	20–22	1.40–1.55	0.45–0.50	0.80–0.85 ⁴
Grower	25–50	2.40–2.45	16–18	1.05–1.15	0.31–0.34	0.65–0.70 ⁴
Finisher (Late)	80–135	2.40–2.47	12–14	0.75–0.85	0.22–0.25	0.45–0.50 ⁴

Transport Stress and the *RYR1* Gene

The physiological stress experienced by swine during transport and lairage directly influences post-mortem muscle glycolysis. Genetic lines carrying the *RYR1* (ryanodine receptor 1) point mutation (the halothane gene) are highly susceptible to malignant hyperthermia⁹. In these animals, acute stress prior to slaughter triggers rapid glycogen depletion and an accelerated drop in muscle pH while the carcass temperature remains high (above 37°C), leading to the denaturation of myofibrillar proteins and the development of pale, soft, and exudative (PSE) meat⁵. Conversely, prolonged transport or excessive lairage times (e.g., 15 hours) exhaust glycogen reserves entirely, resulting in dark, firm, and dry (DFD) meat and an increased incidence of skin damage from pen fighting¹¹. Optimal lairage duration is strictly calibrated between 2 and 3 hours to allow cortisol levels to return to baseline without inducing fatigue or severe glycogen depletion¹².

Stunning Modalities and Exsanguination

The initial stage of the slaughter floor process requires rendering the animal insensible to pain and distress before the initiation of exsanguination.

Electrical and Controlled Atmosphere Stunning

Under frameworks such as the European Union Council Regulation (EC) No 1099/2009, strict technical parameters dictate stunning efficacy¹⁴. Electrical stunning operates by passing an alternating current through the brain, causing massive neural depolarization and an epileptic seizure¹⁷. For pigs, the legislation mandates a minimum current of 1.3 Amperes, maintained for at least 3 seconds¹⁴. High-throughput facilities frequently utilize automated head-to-chest (or head-to-heart) systems; this applies a primary 1.3 A current across the brain followed

immediately by a secondary cycle across the thorax to induce cardiac fibrillation, ensuring the animal cannot regain consciousness prior to blood loss²⁰. However, poorly calibrated electrical parameters can trigger violent muscle contractions that snap capillary beds, causing blood splash (ecchymosis) and bone fractures¹¹.

To circumvent individual handling stress, large-scale operations predominantly employ carbon dioxide (CO_2) stunning in gondola or paternoster systems. Swine are lowered into pits

containing hypercapnic gas mixtures—typically exceeding 80% to 90% CO_2 ¹⁸. While CO_2 stunning facilitates low-stress group handling and reduces the incidence of PSE meat compared to electrical methods, it induces severe mucous membrane irritation and respiratory gasping prior to the loss of consciousness²⁴. Experimental controlled atmosphere stunning (CAS)

systems utilizing hypercapnic-hypoxia mixtures (e.g., 20% CO_2 with $< O_2$ and residual nitrogen or argon) have demonstrated reduced aversion behaviors, though they present higher rates of incomplete stuns and require longer exposure times to ensure death²¹.

Stunning Modality	Minimum Parameters	Physiological Mechanism	Welfare & Quality Implications
Electrical (Head-Only)	1.3 Amperes for $\geq$ 3 seconds ¹⁴	Disruption of neural function; induced epilepsy	High risk of PSE and ecchymosis; immediate unconsciousness ¹¹
Electrical (Head-to-Heart)	$\geq$ 1.3 A (head), $\geq$ 1.0 A (chest) ²⁰	Epilepsy followed by cardiac ventricular fibrillation	Eliminates recovery risk; prolonged insensibility ²⁰
Hypercapnic Gas (CO_2)	$> CO_2$ concentration ¹⁸	Intracellular acidosis and central nervous system depression	Highly aversive prior to unconsciousness; improves meat quality ²⁴
Inert Gas / Hypoxia	20% CO_2 , $< O_2$ (with N_2 /Argon) ²⁵	Anoxia and mild hypercapnia	Reduced initial aversion; higher risk of stun failure ²¹

Thoracic Exsanguination

To secure rapid cerebral ischemia and death, the stun-to-stick interval must be tightly

controlled—ideally under 15 seconds for head-only electrical stunning to prevent the return of brainstem reflexes¹⁸. Exsanguination in swine is performed via a thoracic stick¹⁹. The operator introduces a sterilized knife into the jugular furrow at the base of the neck. The blade is inserted behind the sternum, aimed toward the tail, and pushed deep into the thoracic cavity to sever the brachiocephalic trunk, the common carotid arteries, and the jugular veins near the base of the heart²⁶. Once the handle is flush with the skin, the operator pivots the blade at a 45-degree angle to open the vessels maximally, ensuring a massive and rapid volumetric blood loss²⁶.

Thermodynamics of Scalding and Skin Preservation

Because the skin is retained on the carcass, the thick, deeply embedded bristle hairs must be entirely removed. This is accomplished through scalding, a precision thermodynamic process designed to denature the proteinaceous anchors of the hair follicles without cooking the underlying dermal collagen¹.

Histological Architecture and Collagen Denaturation

The integument of the pig features a stratified squamous keratinized epidermis, a highly developed dermal layer rich in collagen and elastin, and a hypodermis consisting of subcutaneous adipose tissue²⁹. Hair follicles invaginate deeply into the dermis and are stabilized by an inner and outer root sheath composed of highly keratinized cells³¹.

The primary structural component of the dermis is collagen, representing 75% to 90% of its dry weight³⁴. Mammalian collagen exists as a triple-helix structure stabilized by hydrogen bonding³⁴. When subjected to elevated temperatures, the thermal energy disrupts these hydrogen bonds, causing the triple helix to unfold into a disordered, random coil structure—a phase transition known as thermal denaturation³⁴. In porcine skin, this massive structural deterioration and loss of mechanical integrity begins precisely between 60°C and 65°C³⁷.

The objective of scalding is to transmit sufficient thermal energy to soften the keratin within the stratum corneum and the follicular root sheaths, expanding the gap between the feather/hair calamus and the surrounding follicle, thereby loosening the hair³¹. If the scalding temperature is too high (exceeding 62°C to 63°C), the dermal collagen undergoes severe denaturation and coagulates; it physically shrinks around the base of the bristles, permanently trapping them in the skin and rendering mechanical dehairing impossible¹. Conversely, under-scalding (below 58°C) fails to disrupt the keratin network, forcing operators to use excessive mechanical scraping that damages the epidermis¹.

Time-Temperature Matrices and Scalding Modalities

To navigate this narrow thermodynamic window, commercial swine scalding adheres strictly to a time-temperature matrix of 58°C to 62°C (136°F to 143°F) for an exposure duration of 4 to 6 minutes¹. Adjustments are made based on seasonal variations in hair growth. During the "hard-hair" season (autumn), exposure times may be extended to 7.5 or 9.0 minutes at 60°C, whereas the "easy-hair" season (late winter) permits shorter scalding times of 4.5 minutes at 58°C¹.

Target Species / Age	Scalding Designation	Target Water Temperature	Required Immersion Time	Primary Objective
Poultry (Broilers)	Soft/Semi-scald	50°C - 53°C ¹	60 - 180 seconds ¹	Leaves epidermal layer completely intact for display
Poultry (Mature)	Sub/Medium-scald	54°C - 58°C ¹	60 - 120 seconds ¹	Breaks down epidermis; eases coarse feather removal
Swine (Easy-Hair)	Standard Vat/Tunnel	58°C - 59°C ¹	4.5 - 5.0 minutes ¹	Softens follicular keratin; protects dermal collagen
Swine (Hard-Hair)	Extended Scald	60°C - 62°C ¹	6.0 - 9.0 minutes ³⁹	Overcomes dense autumn coat anchoring

Despite the prolonged exposure to 60°C water, the deep muscle tissues remain thermally protected by the subcutaneous fat layer. Thermocouple measurements recorded during commercial scalding demonstrate that while the immediate subdermal interface peaks at 53°C, it rapidly dissipates to 46°C within 1.5 minutes post-scald³⁹. Consequently, deep muscle temperatures remain stable at approximately 40.6°C, ensuring that the process does not artificially accelerate post-mortem glycolysis or negatively impact ultimate meat pH³⁹.

Scalding is executed using two primary industrial architectures:

1. **Vat (Immersion) Scalding:** The carcass is fully submerged in a communal tank. While economically efficient, vat scalding presents significant microbiological hazards. Scald water accumulates feces, dirt, and organic matter, acting as a cross-contamination vector if contaminated water enters the lungs through the sticking wound². Pathogen survival is mitigated by maintaining continuous temperatures above 60°C and occasionally alkalizing the water to a pH of 11.0 using sodium carbonate, which suppresses the survival of *Enterobacteriaceae*².

2. **Tunnel (Condensation) Scalding:** Carcasses are conveyed on overhead rails through enclosed chambers where hot water sprays or heated steam cascade over the skin¹. This isolates individual carcasses, prevents the aspiration of communal water, and allows for superior hygienic control².

Mechanical Dehairing, Singeing, and Polishing

Because the thermal disruption of the follicular sheath is reversible upon cooling, the carcass must transition to mechanical dehairing immediately following the scald².

Scraping and Mechanical Extraction

In automated high-throughput abattoirs, the scalded carcass is deposited into a continuous dehairing machine. These units consist of heavy steel U-shaped frames housing dual rotating shafts equipped with flexible rubber paddles or flails, often tipped with metal scrapers¹. As the carcass tumbles through the machine, the abrasive rotational force of the paddles flexes against the skin, mechanically dragging the loosened bristles out of the follicles². Overhead manifolds continuously spray warm water across the tumbling carcass to flush away the extracted hair and cellular debris into perforated collection trays below¹. In smaller operations, operators utilize bell scrapers—blunt, curved metal tools—to manually shave the carcass following the directional grain of the hair growth².

While highly effective at hair removal, the mechanical dehairing machine is a recognized vector for microbial cross-contamination. The tumbling action vigorously redistributes any residual fecal matter and environmental flora across the entire epidermis, frequently increasing the total aerobic plate count (APC) and spreading indicator organisms like *E. coli*⁴³.

Singeing and Thermal Shock

To eliminate the remaining fine hairs and stubble—particularly around the ears, hooves, and skin folds—the carcass passes through a singeing furnace². The singeing chamber envelops the carcass in direct gas flames, elevating the ambient temperature to approximately 1200°C for a brief 10 to 15-second exposure². This intense burst of thermal energy incinerates the residual keratin fibers and induces a rapid surface dehydration that shrinks and sets the skin, granting the carcass a clean, tight appearance².

Furthermore, the singeing furnace acts as a highly effective surface decontamination step. The extreme, instantaneous heat destroys the vegetative bacteria smeared across the carcass during the dehairing phase. Microbiological baseline studies consistently demonstrate that singeing delivers a substantial log reduction in APC, *Enterobacteriaceae*, and *Salmonella*, effectively resetting the superficial hygiene of the carcass⁴³.

Polishing and Recontamination

Immediately exiting the singe, the carcass is covered in carbonized deposits and singed hair remnants. The carcass enters a polishing machine containing rotating nylon brushes or specialized rubber whips accompanied by high-pressure cold water². The abrasive friction scours the carbon away, yielding the characteristic pale, clean finish of commercial pork. However, polishing presents a paradox in slaughter hygiene. The mechanical scrubbing forces bacteria from the machinery deeply into the now-empty, dilated hair follicles. Studies repeatedly

highlight polishing as a primary mode of carcass recontamination, often causing a massive rebound in total bacterial counts and coliforms⁴³. Consequently, meticulous sanitation of the polishing whips and immediate downstream chemical interventions are required to restore microbiological safety⁴³.

Safe Evisceration and Enteric Contamination Control

The evisceration process marks the transition from external hide management to the extraction of the internal organs. Because the gastrointestinal tract contains billions of colony-forming units of enteric pathogens—including *Salmonella enterica*, *Campylobacter spp.*, and *Yersinia enterocolitica*—any laceration of the viscera will catastrophically contaminate the sterile muscle tissue⁴⁴.

The Bunging Procedure

Evisceration begins at the posterior of the suspended carcass with a critical operation known as bunging. The objective is to physically isolate and seal the rectum to prevent the leakage of fecal material during internal manipulation⁵². The operator utilizes a specialized knife to execute a circular, circum-anal incision around the bung, cutting through the perianal skin and pelvic connective tissue while taking extreme care not to puncture the colon wall⁵².

Once the rectum is mobilized, it is physically drawn outward. Operators then implement a containment strategy: the rectum is tightly tied off with surgical twine, or the entire anal sphincter is inverted into a plastic bag and secured with an elastomeric band or ring-and-tie deployment mechanism⁵². This physical barrier guarantees that as the viscera are later pulled downward through the abdominal cavity, no stomach contents or fecal fluids can backflow onto the exposed pelvic muscles⁵².

Ventral Midline Incision and Extraction

With the bung secured, the abdomen must be opened. The operator executes a shallow vertical incision down the ventral aspect of the carcass, cutting exclusively through the skin and subcutaneous fat to expose the *linea alba*⁵⁶. The *linea alba*—a fibrous, avascular band of connective tissue uniting the abdominal muscles—serves as the anatomical guideline for cavity entry⁵⁷.

To prevent accidental perforation of the intestines, the operator utilizes a "finger guide" technique. The knife is inserted with the blade facing outward (toward the operator), while the non-cutting hand or a specialized shielded guard is inserted behind the blade into the abdominal cavity⁵³. The operator pushes the knife downward toward the sternum, unzipping the abdomen while the hand physically deflects the viscera away from the sharp edge⁵³.

Following the opening of the abdomen, the pelvic girdle (aitch bone) is bisected using a heavy meat cleaver or bone saw⁵³. The operator reaches into the cavity and systematically tears the mesenteric connective tissues that anchor the intestines to the dorsal wall⁵³. The entire gastrointestinal tract, liver, spleen, and the sealed bung are pulled out as a single, intact mass. Finally, the diaphragm is severed, allowing for the extraction of the pluck (the trachea, lungs, and heart)⁵³.

Zero Tolerance Policies and Final Trimming

Regulatory agencies globally recognize evisceration as a paramount Critical Control Point (CCP). In the United States, the Food Safety and Inspection Service (FSIS) enforces a strict "zero tolerance" policy for any visible fecal, ingesta, or milk contamination on the carcass⁵⁰. If visible contamination occurs due to an evisceration error, operators are strictly prohibited from washing the carcass with water, as high-pressure hosing merely dilutes and drives the pathogenic bacteria deeper into the meat fibers⁵⁰. Instead, the contaminated tissue must be physically excised using a sterilized knife in a dedicated trimming step⁵⁰. This zero-tolerance examination is a mandatory precursor to any subsequent chilling or chemical antimicrobial interventions⁵⁰.

Carcass Splitting and Neurological Tissue Management

To facilitate rapid heat dissipation during chilling, simplify grading, and prepare the carcass for primal fabrication, the intact eviscerated carcass is split symmetrically down the sagittal plane into two halves⁶³.

Mechanical Splitting Operations

Splitting requires immense mechanical torque and extreme precision. It is accomplished using either counter-balanced manual bandsaws or advanced robotic systems⁵³. In highly automated facilities (processing upwards of 750 to 1,500 hogs per hour), machines such as the Frontmatec Autoline APS7 or the Buster VI utilize optical sensors and internal back-rollers to map the exact dimensions of the carcass⁶³.

The saw blade—typically designed with 4 teeth per inch (TPI) and a sharp, pointed profile—is optimized to sever heavy bone without wandering⁶³. The blade must travel perpendicularly to the skin surface, bisecting the vertebral bodies and cleanly opening the neural canal⁶⁸. Off-center splits destroy the integrity of the *longissimus dorsi* (loin muscle), significantly reducing the yield of high-value retail cuts and invalidating the standardized grading formulas (such as the 10th rib fat depth equation) used to determine the standardized fat-free lean (SFFL) weight⁶³.

Carcass Measurement Variable	Measurement Location	Utility in Carcass Valuation	Standard SFFL Equation Impact
Hot Carcass Weight	Pre-chill scale	Baseline mass for yield calculation	Positive coefficient (e.g., $+0.465 \times$) ⁷⁰
10th Rib Fat Depth	$3/4$ distance over loin eye ⁷⁰	Predicts total subcutaneous	Heavy negative coefficient (e.g.,

		adiposity	$-21.896 \times$ $)^{70}$
Loin Muscle Area (LMA)	Cross-section at 10th/11th rib ⁷⁰	Direct proxy for total somatic protein	Positive coefficient (e.g., $+3.005 \times$ $)^{70}$

Spinal Cord Extraction and Cross-Contamination

As the splitting saw drives through the vertebral column, it inevitably fragments the spinal cord, generating a highly potent slurry of cerebrospinal fluid, nervous tissue, bone dust, and saw dust⁶⁹. While the risk of Transmissible Spongiform Encephalopathies (TSEs) like BSE is predominantly an issue in cattle, the contamination of pork meat with central nervous system (CNS) tissue remains a strict quality and hygiene violation⁶⁹. Furthermore, Advanced Meat Recovery (AMR) systems require absolute certification that no spinal cord material enters the supply chain, often validated via Glial Fibrillary Acidic Protein (GFAP) rapid testing⁷³.

To eliminate this hazard, slaughter lines deploy specialized vacuum removal systems immediately following the splitting saw. Pneumatic or hydraulic tools equipped with contoured suction nozzles track the open spinal canal⁷⁴. Utilizing high-pressure vacuums, these devices aspirate the fragmented spinal cord, protective sheaths, spinal ganglia, and residual bone dust out of the carcass, leaving a clean, sterile cavity⁶⁵. Simultaneously, the splitting saws undergo continuous hot water or chemical sanitation within their housing between each carcass to prevent the mechanical transfer of infectious material⁶⁹.

Antimicrobial Interventions and Pathogen Reduction

Despite rigorous hygienic controls during evisceration and splitting, total sterility is impossible. Therefore, the pre-chilled carcass is subjected to a comprehensive multi-hurdle decontamination strategy to achieve statutory log reductions of *Salmonella* and *E. coli*.

Organic Acid Applications

The deployment of organic acids—specifically lactic acid and peroxyacetic acid (PAA)—is the primary chemical intervention utilized in modern abattoirs⁷⁸.

Lactic Acid ($C_3H_6O_3$): Naturally occurring and approved by both the FSIS and the European Food Safety Authority (EFSA), lactic acid is typically applied as a 2% to 5% heated spray (42°C to 55°C) across the entire carcass⁷⁸. Its mechanism of action relies on the un-dissociated acid molecule permeating the bacterial cell membrane. Once within the neutral cytoplasm, the acid dissociates, releasing protons that disrupt the proton motive force (PMF) and drastically lower the internal pH to a lethal, unrecoverable level⁷⁸. Extensive validation demonstrates that 2% to 5% lactic acid sprays consistently achieve a 1.5 to >4.0 log reduction in *Salmonella*, *E. coli* O157:H7, and Enterobacteriaceae on carcass surfaces⁶¹.

Peroxyacetic Acid (PAA): PAA is a potent oxidizer synthesized by combining acetic acid and

hydrogen peroxide⁸⁰. Applied via low-pressure sprays at concentrations of 200 to 400 ppm, PAA indiscriminately attacks the bacterial cell wall and disrupts critical enzymatic protein structures⁸⁰. While highly effective—and environmentally favorable due to its rapid breakdown into harmless acetic acid and water—comparative trials on porous meat surfaces often indicate that lactic acid provides slightly superior and more sustained inhibition of *Salmonella* compared to PAA⁶¹.

Antimicrobial Intervention	Standard Concentration	Physiological Mechanism	Typical Log Reduction	Regulatory Stance
Lactic Acid	2.0% - 5.0% ⁷⁹	Disrupts PMF; intracellular acidification ⁸²	1.5 to 4.0 Log CFU/ <i>cm</i> ² [cite: 61, 81]	FSIS & EFSA Approved ⁷⁹
Peroxyacetic Acid (PAA)	200 - 400 ppm ⁸⁵	Severe oxidative damage to cell wall ⁸⁰	1.0 to 2.5 Log CFU/ <i>cm</i> ² [cite: 61, 88]	FSIS Approved; breaks down to water/acetic acid ⁸⁴
Steam Vacuuming	> Steam ⁸⁹	Instantaneous thermal protein denaturation ⁹⁰	1.0 to 2.0 Log CFU/ <i>cm</i> ² [cite: 55, 90]	Approved for localized fecal excision substitute ⁹¹

In addition to chemical sprays, establishments heavily rely on localized steam vacuuming. Steam vacuum wands blast pressurized steam (exceeding 82°C) onto minor, localized contamination spots, instantly denaturing bacterial proteins. The wand immediately vacuums away the resulting condensation and debris, providing a powerful thermal kill without causing the underlying muscle tissue to cook or discolor⁹⁰.

Trichinella spiralis Surveillance

Beyond bacterial pathogens, securing the safety of the pork supply requires the mitigation of parasitic hazards, most notably the zoonotic nematode *Trichinella spiralis*⁴. Animals reared in fully integrated indoor confinement facilities undergo strict biosecurity protocols (rodent exclusion, strict feed controls prohibiting swill feeding) and may be granted "Controlled Housing" status, subjecting them only to statistical risk-based testing⁴.

However, carcasses originating from pastoral, outdoor, or non-certified facilities must undergo mandatory 100% individual post-mortem testing before commercial release⁴. The global standard for this surveillance is the Magnetic Stirrer Method (MSM) for Pooled Sample Digestion (ISO 18743:2015)⁴. Technicians excise samples from the muscular crus of the diaphragm—the anatomical predilection site for encysted larvae⁴. The tissue is mechanically blended and introduced into a heated 44°C to 46°C fluid containing hydrochloric acid and high-activity

pepsin⁴. This enzymatic bath dissolves the host's muscular and collagenous tissues over 30 minutes, simulating gastric digestion and releasing any viable larvae, which are subsequently collected via sedimentation and identified under a stereomicroscope⁴. Un-tested carcasses must undergo statutory thermal inactivation or strict commercial freezing (e.g., -15.0°C for 20 days) to destroy the larvae⁴.

Post-Mortem Chilling and Pork Belly Quality

The final critical phase of the slaughter process involves ultra-rapid temperature reduction. Prompt chilling arrests remaining microbial proliferation, limits post-mortem drip loss (shrinkage), and firmly sets the physical properties of the skin and subcutaneous fat⁹⁸.

Thermodynamics of Ultra-Rapid Chilling

Because porcine carcasses retain an insulating layer of dermal tissue and thick subcutaneous fat, extracting core thermal energy is substantially more difficult than in skinned beef carcasses. Modern facilities execute a two-phased ultra-rapid (blast) chilling protocol⁹⁸. Initially, the split halves are conveyed into high-velocity blast tunnels where they are bombarded with frigid air (-4°C to -6°C) moving at 2 to 4 meters per second for one to three hours⁹⁸.

This extreme thermal shock rapidly drops the surface temperature, freezing residual moisture and instantly solidifying the liquid lipids within the adipocytes. This firming action is vital for preserving the macroscopic structure of the carcass, facilitating clean, automated butchery of the primal joints⁹⁸. Following the blast phase, carcasses transition into equilibration coolers maintained at -1°C to 2°C with low air circulation for an additional 14 to 16 hours, allowing the deep tissue temperatures to stabilize⁹⁸.

However, thermodynamic velocity must be balanced against the biochemical state of the muscle. If the core muscle temperature is forced below 10°C before the ultimate pH has fallen below 6.0—while cellular ATP is still abundant—the sarcoplasmic reticulum loses its capability to sequester calcium ions⁹⁸. This triggers a massive, irreversible contraction of the myofibrils known as cold shortening, yielding severely toughened meat⁹⁸.

Implications for Skin and Bacon Fabrication

The interplay of precise nutritional management (controlling Iodine Values), optimal scalding, and blast chilling culminates in the structural quality of the pork belly. High-quality bacon fabrication demands a firm, structurally cohesive belly with a distinct lean-to-fat ratio⁸. The ultra-rapid dehydration of the skin during the blast chill sets the denatured dermal collagen matrix, preserving the tensile strength of the stratum corneum³⁴.

This structural firmness is paramount; bellies exhibiting soft fat (driven by high dietary polyunsaturated fatty acids) or damaged skin (from improper scalding temperatures) suffer from layer separation, poor slicing yields, and inferior sensory qualities (e.g., uneven crackling expansion during high-temperature pan-frying)³. Consequently, every parameter—from the feed trough to the chilling tunnel—must operate within strict tolerances to produce a commercially viable, skin-on pork product.

Conclusion

The successful conversion of a live porcine animal into a biologically secure, high-yield carcass

is a triumph of applied biochemistry and mechanical engineering. The requirement to preserve the skin dictates the entirety of the slaughter timeline. Thermal scalding must be perfectly calibrated to an exact time-temperature matrix to denature follicular keratin without destroying the tensile strength of the dermal collagen. Because the subsequent mechanical dehairing and polishing operations unavoidably redistribute superficial flora, the ensuing evisceration process relies on absolute anatomical precision—securing the bung and navigating the linea alba—to prevent catastrophic enteric contamination.

When reinforced by automated skeletal splitting, immediate vacuum extraction of neurological tissue, and targeted organic acid decontamination, these rigorous protocols guarantee microbiological safety. Ultimately, it is the seamless integration of pre-slaughter physiological control, precision thermodynamic processing, and uncompromising HACCP-driven hygiene that sustains the safety, yield, and global economic viability of the commercial pork industry.

Works cited

1. Scalding and Its Significance in Livestock Slaughter ... - ResearchGate, https://www.researchgate.net/publication/279195003_Scalding_and_Its_Significance_in_Livestock_Slaughter_and_Wholesome_Meat_Production
2. Dressing Techniques for Pigs: From Scalding to Evisceration, <https://agriculture.institute/meat-animals-abattoir-practices/dressing-techniques-for-pigs/>
3. The Quality Traits of Pork Belly and Impact Factors of ... - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11907425/>
4. 50 Domestic Pig Rearing and Nutrition.pdf
5. Critical Points Affecting Fresh Pork Quality within the Packing Plant, <https://porkgateway.org/resource/critical-points-affecting-fresh-pork-quality-within-the-packing-plant/>
6. Fitness Check of the EU Animal Welfare legislation - {SWD(2022, https://food.ec.europa.eu/system/files/2022-10/aw_eval_revision_swd_2022-328_en.pdf
7. Scalding Technique in Poultry & Pig Slaughtering - Dalvoy, <https://www.dalvoy.com/en/upsc/mains/previous-years/2017/ani-husb-veter-scienc-e-paper-ii/scalding-technique-poultry-pig>
8. BELLY THICKNESS EFFECTS ON THE PROXIMATE, https://www.researchgate.net/publication/230067850_BELLY_THICKNESS_EFFECTS_ON_THE_PROXIMATE_COMPOSITION_PROCESSING_AND_SENSORIAL_CHARACTERISTICS_OF_BACON
9. Effects of different stunning methods on the welfare and carcass, <https://repository.up.ac.za/bitstreams/0a80279b-5961-4f44-8b99-ee69d6775849/download>
10. Effect of unloading, lairage, pig handling, stunning and season on, https://www.researchgate.net/publication/45820849_Effect_of_unloading_lairage_pig_handling_stunning_and_season_on_pH_of_pork
11. The influence of transport on pig meat quality - ResearchGate, https://www.researchgate.net/publication/322859823_The_influence_of_transport

- [_on_pig_meat_quality](#)
12. Effects of lairage and slaughter conditions on animal welfare and, <https://cdnsiencepub.com/doi/pdf/10.4141/cjas10020>
 13. Biosecurity and Lairage Time versus Pork Meat Quality Traits in a, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20230148253>
 14. Quantification of the Effects of Electrical and CO2 Stunning on ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9312002/>
 15. Animal welfare infringements at slaughter in Sweden, <https://research.slu.se/files/19135404/fulltext.pdf>
 16. Welfare of animals at slaughter and killing: A new regulation on the, https://www.researchgate.net/publication/225136367_Welfare_of_animals_at_slaughter_and_killing_A_new_regulation_on_the_protection_of_animals_at_the_time_of_killing
 17. Verification of the technical parameters of head-only electrical, https://www.researchgate.net/publication/44589943_Verification_of_the_technical_parameters_of_head-only_electrical_stunning_of_pigs_under_commercial_conditions
 18. Humane Slaughter: Methods and Importance in Meat Production, <https://agriculture.institute/meat-animals-abattoir-practices/humane-slaughter-methods-importance/>
 19. Slaughter and Dressing | Veterian Key, <https://veteriankey.com/slaughter-and-dressing/>
 20. (PDF) Assessing pig welfare at stunning in Northern Italy, https://www.researchgate.net/publication/266030420_Assessing_pig_welfare_at_stunning_in_Northern_Italy_commercial_abattoirs_using_electrical_method
 21. A comparative literature review of current and potential commercial, https://australianpork.com.au/sites/default/files/2024-08/Final_report_Stunning%20review.pdf
 22. Effects of slaughter method on carcass and meat characteristics in, https://projectblue.blob.core.windows.net/media/Default/Research%20Papers/Bee%20&%20Lamb/slaughter_and_meat_quality_feb_2012-final-report.pdf
 23. Unnecessary suffering during the slaughter of cattle and pigs, <https://www.frontiersin.org/journals/animal-science/articles/10.3389/fanim.2025.1633616/full>
 24. The selection of alternatives to high CO2 stunning to be tested in the, <https://edepot.wur.nl/691217>
 25. Animal Welfare and Meat Quality Assessment in Gas Stunning, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7766013/>
 26. If we can't ban halal meat, we should at least let people know when, https://www.reddit.com/r/unitedkingdom/comments/4ww50y/if_we_cant_ban_halal_meat_we_should_at_least_let/
 27. independent-scientific-literature-review-livestock-welfare-processing, <https://www.agriculture.gov.au/sites/default/files/documents/independent-scientific-literature-review-livestock-welfare-processing-establishments.pdf>
 28. At-Home Hog Slaughter - SDSU Extension,

- <https://extension.sdstate.edu/home-hog-slaughter>
29. (PDF) DISTINCTIVE FEATURES OF THE PIGLET SKIN REVEALED,
https://www.researchgate.net/publication/357646900_DISTINCTIVE_FEATURES_OF_THE_PIGLET_SKIN_REVEALED_BY_LIGHT_AND_TRANSMISSION_ELECTRON_MICROSCOPY
 30. Challenges of Porcine Wound Models: A Review - MDPI,
<https://www.mdpi.com/2673-8937/5/1/4>
 31. The Defeathering Effect by Scalding in Chickens Follows Their,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10451932/>
 32. Skin tissue regeneration for burn injury,
<https://www.huasan.net/wp-content/uploads/2025/12/Skin-tissue-regeneration-for-burn-injury.pdf>
 33. Wound healing compositions containing keratin biomaterials,
<https://patents.google.com/patent/US8258093B2/en>
 34. Burn-related Collagen Conformational Changes in ex vivo Porcine,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6915721/>
 35. Mechanical Properties of Porcine and Fish Skin-Based Collagen,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8271417/>
 36. Differential scanning calorimetry and temperature dependence of,
https://www.researchgate.net/publication/243301733_Differential_scanning_calorimetry_and_temperature_dependence_of_electric_conductivity_in_studies_on_denaturation_process_of_bone_collagen
 37. Mechanical behavior of full-thickness burn human skin is rate ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11096406/>
 38. Effective enzymatic debridement of burn wounds depends on the,
https://www.researchgate.net/publication/342379527_Effective_enzymatic_debridement_of_burn_wounds_depends_on_the_denaturation_status_of_collagen
 39. The effect of scalding on subcutaneous and ham temperatures and,
<https://pubmed.ncbi.nlm.nih.gov/22060746/>
 40. The effects of duration of processing on pork quality,
<https://dr.lib.iastate.edu/server/api/core/bitstreams/fe2a9a29-1d04-41cb-86f8-6c1fc4558d89/content>
 41. Early postmortem harvest practices influence pork color - VTechWorks,
<https://vtechworks.lib.vt.edu/server/api/core/bitstreams/f22ae594-d928-4b0a-8c1e-4ab9c2f4b58f/content>
 42. Modeling nonlinear inactivation of hygiene indicator bacteria in pig,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11711692/>
 43. Risk-based Determination of Critical Control Points for Pork Slaughter,
<https://t-stor.teagasc.ie/server/api/core/bitstreams/f61c6e35-b335-4eac-a9c4-d6a71463e1ff/content>
 44. Print version ISSN 1315-2556,
http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S1315-25562004000100015
 45. Effect of dehairing operations on microbiological quality of swine,
<https://pubmed.ncbi.nlm.nih.gov/10606156/>
 46. Singeing to improve visual aspect and microbiological quality of pig,

- <https://www.scielo.br/j/rbspa/a/7N4D9psKwjv8VZLbyNYbCzS/?format=pdf&lang=en>
47. Microbiological contamination of pig carcasses at different stages of, <https://pubmed.ncbi.nlm.nih.gov/17133797/>
 48. Pig scalding by pushing, discover how it helps your facility - mecanova, <https://mecanova.eu/en/pig-scalding-by-pushing>
 49. Compliance Guideline for Controlling Salmonella in Market Hogs, <https://www.fsis.usda.gov/sites/default/files/import/Controlling-Salmonella-in-Market-Hogs.pdf>
 50. HACCP Plan for Pork Slaughterhouse | PDF | Animal Slaughter, <https://www.scribd.com/document/712785968/HACCP-PLAN-PORK-SLAUGHTERING>
 51. Introducing a New Digestibility Assay Procedure - CABI Digital Library, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20123111739>
 52. Bung Sealing in Slaughtering Process | PDF - Scribd, <https://www.scribd.com/document/554372986/RING-AND-TIE-BUNG-SLAUGHTERING-RING-PROCESS>
 53. Slaughter Day: Part 2, Eviscerating and Splitting Pigs, <https://lowtechinstitute.org/2018/02/07/slaughter-day-part-2-eviscerating-and-splitting-pigs-from-contributor-matt-miles/>
 54. Evisceration - Surv-live'l, <https://survivel.cultu.be/evisceration>
 55. Agenda Item 4 CX/FH 15/47/5 add.1 October 2015 JOINT FAO/WHO, https://www.fao.org/fao-who-codexalimentarius/sh-proxy/es/?Ink=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252FMeetings%252FCX-712-47%252FWDs%252Ffh47_05_add1e.pdf
 56. Soft Tissue Surgery, An Issue of Veterinary Clinics of North America, <https://dokumen.pub/soft-tissue-surgery-an-issue-of-veterinary-clinics-of-north-america-exotic-animal-practice-0323414753-9780323414753.html>
 57. Peritoneal Procedures | Clinical Gate, <https://clinicalgate.com/2015/03/26/peritoneal-procedures/>
 58. Atlas of Large animal surgery Veterinary | PDF - Slideshare, <https://www.slideshare.net/slideshow/atlas-of-large-animal-surgery-veterinary/266545106>
 59. Herniorrhaphy (Proceedings) - DVM360, <https://www.dvm360.com/view/herniorrhaphy-proceedings>
 60. Verification of Procedures for Controlling Fecal Material, Ingesta, <https://www.fsis.usda.gov/policy/fsis-directives/6420.2>
 61. Effects of antimicrobial interventions on food safety and an, <https://mountainscholar.org/bitstreams/08c888c9-f2c3-49ee-a05d-f0562b1e9bcd/download>
 62. HACCP Model for Traditional Swine Slaughter - Maine.gov, https://www.maine.gov/dacf/qar/inspection_programs/documents/mmpi/HACCP-Model-Swine-Slaughter-Traditional.pdf
 63. Autoline split saw for pigs - Automatic hog splitting saw - Frontmatec, <https://www.frontmatec.com/en/pork-solutions/clean-line-chill-room/autoline-machi>

- [nes/autoline-split-saw-aps7](#)
64. Procedures for Estimating Pork Carcass Composition,
<https://porkgateway.org/resource/procedures-for-estimating-pork-carcass-composition/>
 65. BUSTER VI | Electric Splitting Saw | Jarvis UK & Ireland,
<https://jarvisuk.com/products/buster-vi/>
 66. Carcass Bandsaw - MESI - Saws & Blades,
<https://mesisaws.com/en/carcass-bandsaw/>
 67. Meat - Burton Saw International,
<https://burtonsaw.co.uk/bandsaw-blades/food-industry/meat/>
 68. Swine Extension Program - Animal and Range Sciences,
https://animalrangeextension.montana.edu/swine/4_rib.html
 69. Overview of current and alternative slaughter practices - PoPuPS,
<https://popups.uliege.be/1780-4507/index.php?id=17415&file=1&pid=14187>
 70. Procedures for Estimating Pork Carcass Composition,
<https://porkgateway.org/wp-content/uploads/2015/07/procedures-for-estimating-pork-carcass-composition.pdf>
 71. Details and results of controlled trials with a carcass-splitting saw,
https://www.researchgate.net/figure/Details-and-results-of-controlled-trials-with-a-carcass-splitting-saw_tbl2_8261002
 72. Meat Technology 61-02 Scientific Journal,
http://www.journalmeattechnology.com/index.php/meat_technology/issue/download/15/15
 73. Removal of Spinal Cord during Slaughter - BIFSCo,
https://www.bifsc.org/Media/BIFSCO/Docs/spinal_cord_removal2002.pdf
 74. Standard Practice for Splitting Saws - Kentmaster China,
<http://en.kentmaster-china.com/news/76.html>
 75. method for removing the spinal cord from slaughter animals,
<https://patentscope.wipo.int/search/en/WO2003086084>
 76. Pork Processing Archives - Jarvis Anz,
<https://jarvisanz.com.au/product-category/pork-processing/>
 77. (PDF) Automation for the Modern Slaughterhouse - ResearchGate,
https://www.researchgate.net/publication/329277230_Automation_for_the_modern_slaughterhouse
 78. The Use of Organic Acids (Lactic and Acetic) as a Microbial ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8534660/>
 79. Evaluation of the safety and efficacy of the organic acids lactic ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7009387/>
 80. Evaluation of Peracetic Acid and Lactic Acid on the Bacterial,
https://www.researchgate.net/publication/397413896_Evaluation_of_Peracetic_Acid_and_Lactic_Acid_on_the_Bacterial_Decontamination_of_Chilled_Pork
 81. Reference Document: An/microbial Interventions for Beef,
<https://www.haccpalliance.org/wp-content/uploads/2022/01/Antimicrobial-Interventions-for-Beef.pdf>
 82. The Use of Organic Acids (Lactic and Acetic) as a Microbial ... - MDPI,

- <https://www.mdpi.com/2304-8158/10/10/2293>
83. MICROBIAL CHANGES ON THE SURFACE OF PORK,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20083151868>
 84. EFSA presentation on Peroxyacetic acid | Food Safety,
https://food.ec.europa.eu/system/files/2016-10/biosafety_fh_mc_efsa-presentation-peroxyacetic-acid.pdf
 85. New Technology - C29 Summary,
https://www.fsis.usda.gov/sites/default/files/media_file/2021-08/New_Technology_C29_Summary_FY2003.pdf
 86. Antimicrobial Interventions for Beef - BIFSCo,
<https://www.bifSCO.org/Media/BIFSCO/Docs/antimicrobial-interventions-for-beef.pdf>
 87. (PDF) Evaluation of the safety and efficacy of the organic acids lactic,
https://www.researchgate.net/publication/329604217_Evaluation_of_the_safety_and_efficacy_of_the_organic_acids_lactic_and_acetic_acids_to_reduce_microbiological_surface_contamination_on_pork_carcasses_and_pork_cuts
 88. Effect of Peracetic Acid Solutions and Lactic Acid on Microorganisms,
https://refubium.fu-berlin.de/bitstream/fub188/27993/1/Thomas_Effect_2020.pdf
 89. Spatiotemporal Distribution of Microbiota During Fresh Pork,
<https://www.iastatedigitalpress.com/mmb/article/id/20198/>
 90. Carcass Steam Sanitation in Industrial Slaughterhouses - Jarvis Italia,
<https://jarvisitalia.it/en/steam-sanitation-of-carcasses-reducing-bacterial-load-without-compromise/>
 91. Carcass decontamination methods in slaughterhouses: a review,
<https://scispace.com/pdf/carcass-decontamination-methods-in-slaughterhouses-a-review-556n5hproi.pdf>
 92. Steam-vacuum treatment of pig carcass surfaces - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/41443440/>
 93. AVMA Guidelines for the Humane Slaughter of Animals: 2016 Edition,
<https://www.cnr-bea.fr/wp-content/uploads/2019/10/AVMA-Humane-Slaughter-Guidelines.pdf>
 94. How to butcher a Hog, Step by Step, the Entire Breakdown. - YouTube,
<https://www.youtube.com/watch?v=LSqJ0pr2RUk>
 95. Livestock at Slaughtering Establishments - SCARM Report 79,
<https://www.publishing.csiro.au/ebook/download/pdf/2975>
 96. How to Butcher a Pig | Every Cut Explained Plus Ham and Sausage,
<https://www.youtube.com/watch?v=JkdgdKlpy9I>
 97. Preslaughter and slaughter factors affecting meat quality in lambs,
<https://mro.massey.ac.nz/server/api/core/bitstreams/26052293-ac6e-479e-b30f-b297ddb556f8/content>
 98. Meat Preservation and Processing | Veterian Key,
<https://veteriankey.com/meat-preservation-and-processing/>
 99. The effects of spray and blast-chilling on carcass shrinkage and pork,
<https://pubmed.ncbi.nlm.nih.gov/22060743/>
 100. Chilling pig carcasses: Effects on temperature, weight loss and,

<https://www.semanticscholar.org/paper/Chilling-pig-carcasses%3A-Effects-on-temperature%2C-and-Wal-Engel/88379dc36f9a70f11c603920734b98621f198345>

101. Quality Variation of Pork Bellies by Cutting Manner and Quality Grade,
<https://www.mdpi.com/2304-8158/13/19/3129>
102. Effect of Thickness of Pork Belly on Physicochemical and Sensory,
<https://agris.fao.org/search/en/providers/122646/records/65bbb72540e64bb074ab0aef>