

Educational Manual of Ethnoveterinary Botanicals and Fundamental Livestock Surgical Interventions

The management of a dual-purpose or specialized livestock herd represents a highly sophisticated intersection of veterinary pathology, ethnoveterinary pharmacology, and agronomic science. In contemporary agricultural paradigms—particularly those operating under sustainable and organic legislative frameworks such as the European Union’s Regulation (EU) 2018/848—there is a statutory mandate to prioritize phytotherapeutic and homeopathic products over chemically synthesized allopathic veterinary medicinal products, provided their therapeutic efficacy is guaranteed¹. This regulatory shift is largely driven by the escalating global crisis of antimicrobial resistance (AMR), which requires a systemic reduction in prophylactic and metaphylactic antibiotic usage across all animal husbandry sectors¹. Consequently, modern veterinary practitioners and herd managers must possess a profound understanding of not only traditional surgical interventions and diagnostic modalities but also the complex pharmacognosy of botanical remedies. This manual provides an exhaustive educational overview of diagnostic methodologies, ethnoveterinary botanical applications, and elementary surgical protocols critical to the management of common livestock ailments. It establishes the critical baseline that surgical and botanical interventions cannot succeed in a vacuum; they must be integrated with precise nutritional engineering, epidemiological surveillance, and a nuanced understanding of animal metabolism.

Nutritional and Metabolic Foundations of Disease Diagnostics

Before initiating targeted botanical therapies or surgical interventions, the metabolic baseline of the herd must be exhaustively evaluated. Prominent dual-purpose breeds, such as the Fleckvieh (German Simmental), the French Normande, and the Shorthorn, exhibit unique physiological traits that make them highly adaptable yet nutritionally demanding⁵. These cattle are expected to deliver substantial lactation yields—often ranging from 7,000 to 11,000 kg of milk per lactation—while simultaneously maintaining a robust, muscled frame capable of superior tissue accretion⁵.

Breed	Primary Geographic Adaptation	Expected Milk Yield (kg/lactation)	Carcass Characteristics	Unique Physiological Traits
Fleckvieh (Simmental)	Temperate / Alpine	7,000 - 11,000	Excellent muscling;	Superior body condition

			57-60% dressing	retention; heat tolerant ⁵ .
Normande	Temperate / Highland	6,000 - 8,000	Well-marbled beef	Exceptionally high butterfat and protein; optimized for cheese ⁵ .
Brown Swiss	Temperate / Subtropical	7,500 - 9,500	Moderate to heavy muscling	High component milk; robust longevity; moderate heat tolerance ⁵ .
Dexter	Marginal / Low-Input	2,000 - 3,500	Lean, tender beef; high dressing	Extreme feed efficiency on poor forage; suitable for smallholdings ⁵ .

To sustain this dual functionality, herd management relies on the accurate prediction and optimization of dry matter intake (DMI). The physical capacity of the rumen, combined with the metabolic feedback mechanisms of the cow, tightly regulates daily consumption⁵. A mature dual-purpose cow typically consumes between 3.0% and 3.5% of its body weight in dry matter per day, peaking at up to 4.0% during mid-lactation when energy demands are at their zenith⁵.

Environmental variables severely impact this metric; ambient temperatures exceeding 25°C trigger heat stress adaptations that can suppress DMI by 10% to 35%, simultaneously depressing milk synthesis and exacerbating negative energy states⁵.

Energy Partitioning and Negative Energy Balance (NEB)

Nutritionists currently rely on the 2023 German Society of Nutrition Physiology (GfE) standards, which transitioned from the Net Energy for Lactation (NEL) system to a unified framework based on Metabolizable Energy (ME)⁵. Under this framework, maintenance energy is calculated

relative to metabolic body weight, expressed mathematically as $0.515 \text{ MJ} \times \text{BW}^{0.75}$ ⁵.

For a standard 650 kg cow, this equates to a daily baseline maintenance requirement of approximately 66.5 MJ ME, to which a standard 10 MJ activity allowance is added⁵.

Despite precise rationing, high-producing cows inevitably encounter a severe Negative Energy Balance (NEB) during the transition period (three weeks pre-calving to three weeks post-calving). Because the volumetric capacity of the rumen limits DMI immediately

post-calving, the caloric energy required for peak milk production vastly outstrips dietary energy absorption⁵. During NEB, the cow enters homeorhesis, mobilizing substantial adipose reserves. This lipolysis elevates concentrations of non-esterified fatty acids (NEFAs) and beta-hydroxybutyrate (BHBA), which overwhelm the liver's oxidative capacity, inducing endoplasmic reticulum stress and subclinical ketosis⁵. Comprehensive lipidomic profiling identifies specific polar lipids, such as Phosphatidylethanolamine (PE 33:1) and Lactosylceramide (LacCer 32:1), as highly accurate biomarkers of severe energy deficits⁵. Surgical interventions performed on animals in severe NEB are fraught with complications, as hepatic lipidosis directly suppresses fibroblast proliferation and immune competence during wound healing.

Subacute Ruminal Acidosis (SARA) and Physically Effective Fiber

In the beef-producing cohort of the herd, achieving rapid tissue accretion requires high-energy rations with concentrate-to-forage ratios frequently exceeding 85%⁵. However, aggressive supplementation with rapidly fermentable carbohydrates introduces the severe risk of Subacute Ruminal Acidosis (SARA). SARA is defined as a state wherein the ruminal pH remains depressed below 5.8 for more than 5.24 hours per day⁵.

The acidic environment induces hyperkeratosis and parakeratosis of the ruminal epithelium, shedding mucosal tissues and allowing pathogens to cross into the portal blood circulation, which frequently culminates in liver abscesses and secondary laminitis⁵. A critical diagnostic indicator of SARA is the physical composition of the feces. During SARA episodes, the fractional outflow rate of the rumen increases abnormally. Fecal analysis revealing that more than 4% of the dry matter consists of particles larger than 4 mm is a strong diagnostic indicator of ruminal dysfunction⁵. Rations must be mathematically balanced with physically effective neutral detergent fiber (peNDF), targeting an optimal inclusion rate near 10.4%, to stimulate rumination and the copious production of endogenous, bicarbonate-rich saliva⁵.

Mineral Antagonisms and Agronomic Toxicology

Livestock maintained on pasture-based rotational grazing systems face unique geochemical hazards. The most economically significant antagonism occurs between copper, molybdenum, and sulfur. Within the anaerobic rumen, microbes reduce dietary sulfates into sulfides, which rapidly react with molybdenum to form thiomolybdates⁵. These compounds possess an extraordinary affinity for copper, binding it into insoluble complexes that render dietary copper entirely unabsorbable⁵. This induces a severe secondary copper deficiency characterized by achromotrichia (fading hair coat), spontaneous bone fractures, and profound suppression of neutrophil activity⁵.

Additionally, lush spring swards frequently exhibit elevated potassium and nitrogen levels alongside depressed sodium. Excessive potassium alters the electrochemical gradient across the ruminal wall, severely depolarizing the apical membrane and depressing active magnesium transport, resulting in fatal hypomagnesemia (grass tetany)⁵. The risk is quantitatively modeled

using the tetany ratio; when the index exceeds 2.2 mEq/kg of dry matter, the incidence of clinical hypomagnesemia escalates exponentially⁵.

Furthermore, if late-season grazing utilizes standing brassicas or heavily fertilized autumn swards, herd managers must vigilantly monitor for nitrate toxicity. When plant growth slows due to frost or drought, metabolic reduction of nitrates ceases, causing them to accumulate in the lower stems⁵. Ruminants reduce these nitrates to nitrites, which aggressively oxidize hemoglobin into methemoglobin. Because methemoglobin is incapable of binding oxygen, the animal suffers rapid tissue hypoxia, chocolate-brown blood, tremors, and fatal asphyxiation⁵.

Nitrate-Nitrogen (NO ₃ -N) Concentration	Toxicity Risk Level	Feeding Recommendation
0 - 1,500 ppm	Safe	Feed without restriction ⁵ .
1,500 - 5,000 ppm	Caution	Limit to 50% of DMI; dilute with safe hay ⁵ .
5,000 - 9,000 ppm	Danger	Limit to 33% of total ration; observe carefully ⁵ .
> 9,000 ppm	Lethal	Do not feed; extreme risk of acute mortality ⁵ .

Parasite Epidemiology and Targeted Selective Treatment (TST)

Gastrointestinal nematodes (GINs) continually threaten pastured cattle, with the brown stomach worm, *Ostertagia ostertagi*, representing the most devastating pathogen in temperate climates⁵. The parasite targets the abomasum, destroying the acid-secreting gastric glands. This halts hydrochloric acid production, raises abomasal pH, and prevents the activation of pepsinogen into pepsin, collapsing protein digestion and inducing profuse diarrhea and massive weight loss⁵.

The parasite possesses an evolutionary survival mechanism known as hypobiosis. Larvae ingested during late autumn arrest their development at the L4 stage, remaining dormant within the abomasal glands throughout the winter⁵. Their synchronous mass emergence the following spring causes Type II ostertagiosis, characterized by acute clinical signs and high mortality⁵.

Historically, parasite control relied on blanket, prophylactic administration of broad-spectrum anthelmintics (such as macrocyclic lactones and benzimidazoles), which precipitated a global crisis of anthelmintic resistance⁵. Modern epidemiological management mandates a transition to Targeted Selective Treatment (TST) protocols to preserve "refugia"—a sub-population of susceptible, unexposed parasites maintained on the pasture to dilute the genetic pool of resistant worms⁵. Blanket treatments are abandoned; interventions are dictated by algorithmic

surveillance. For growing cattle, selectively drenching only animals that fall below an average daily weight gain (ADWG) threshold of 683 g/day maintains herd performance while reducing anthelmintic usage by up to 95%⁵. In mature milking herds, bulk tank milk is evaluated via the *Ostertagia ostertagi* antibody ELISA (the MOO test) to quantify the optical density ratio (ODR) of parasite-specific antibodies, allowing targeted interventions only when pathogenic thresholds are breached⁵.

Diagnostic Protocols for Intramammary Infections (Mastitis)

Clinical and subclinical mastitis exact a severe economic toll through depressed milk yield, compromised physical milk quality, and the costs associated with premature culling⁶. The pathology is driven by bacterial colonization of the mammary tissue, prompting a massive influx of leukocytes (white blood cells) to the gland, which is quantified as the somatic cell count (SCC)⁸.

Somatic Cell Counting and the California Mastitis Test (CMT)

While laboratory-derived SCC remains the international diagnostic standard, routine deployment is economically and logistically prohibitive for daily herd monitoring⁹. The California Mastitis Test (CMT) serves as an indispensable, rapid, cow-side diagnostic assay designed to estimate the SCC of individual quarters⁷.

The CMT utilizes a specialized reagent consisting of a 3% sodium lauryl sulfate anionic detergent coupled with bromocresol purple (acting as a pH indicator)⁸. When equal volumes of foremilk and reagent are agitated in a specialized paddle, the detergent lyses the cellular membranes of the somatic cells⁸. The subsequent release and precipitation of nuclear deoxyribonucleic acid (DNA) instantly transforms the mixture into a viscous gel⁷. The visual thickness and cohesiveness of this gel are directly proportional to the leukocyte concentration, providing a reliable estimation of subclinical inflammation⁷. Simultaneously, the bromocresol purple detects the alkaline pH shift that typically accompanies vascular leakage during severe glandular inflammation, turning the mixture a deep purple⁸.

CMT Score	Visible Reaction Characteristics	SCC Estimate (cells/mL)	Diagnostic Interpretation
Negative (0)	Fluid mixture, no precipitation	<	Healthy Quarter ⁸ .
Trace (T)	Slight precipitate, disperses with movement	200, 000 —	Subclinical Mastitis ⁸ .

Weak Positive (1)	Distinct precipitate without heavy gelation	400, 000 —	Subclinical Mastitis ⁸ .
Distinct Positive (2)	Immediate thickening, clear gel formation	1, 500, 000 —	Serious Infection ⁸ .
Strong Positive (3)	Highly cohesive, convex gel mass	>	Severe Infection ⁸ .

Diagnostic validation studies indicate that when establishing a threshold of 310,000 cells/mL to delineate subclinical mastitis, the sensitivity and specificity of field tests approach 92.6% and 91.5%, respectively¹³. However, practitioners must interpret CMT results with clinical nuance. The test frequently yields false-positive reactions during the very early colostral phase and in late lactation due to naturally elevated cellular shedding unassociated with active infection⁷. Furthermore, rheological variations in the CMT gel can occur despite identical SCC values. Advanced diagnostic frameworks evaluate the differential somatic cell count (DSCC), recognizing that structural differences between cell types—specifically the varying proportions of macrophages versus polymorphonuclear neutrophils (PMNs)—heavily influence the precipitation dynamics of the DNA¹². During the healing phase of mastitis, macrophage proportions increase even as total SCC declines, occasionally triggering prolonged positive CMT reactions¹².

Ethnoveterinary Botanicals and Pharmacognosy

In light of the European Green Deal and evolving organic legislation restricting prophylactic allopathic treatments, the integration of ethnoveterinary medicine (EVM) provides critical therapeutic alternatives¹. EVM relies on the extraction and delivery of biologically active secondary plant compounds—including phenolics, quinones, terpenoids, and alkaloids—formulated as decoctions, infusions, tinctures, and poultices¹⁴.

Phytotherapy for Udder Edema and Mastitis

The intramammary application of botanicals is frequently utilized to augment natural immune defenses without generating antibiotic residues in the bulk tank⁴. Lamiaceae species, specifically thyme (*Thymus vulgaris*), oregano (*Origanum vulgare*), and sage (*Salvia officinalis*), are deployed extensively⁶. The efficacy of these plants relies upon highly concentrated volatile essential oils, predominantly thymol, carvacrol, and 1,8-cineole⁶. These compounds exhibit profound lipophilic properties, enabling them to disrupt bacterial cell membranes and inhibit the extracellular biofilm matrices produced by *Staphylococcus aureus*⁶. Traditional protocols formulate these herbs into concentrated infusions utilized as topical udder washes or post-milking teat dips⁶.

Turmeric (*Curcuma longa*) is another foundational anti-inflammatory botanical. In field

applications, a finely ground paste combining turmeric, ginger, and salt is applied directly to indurated quarters twice daily⁶. The bioactive curcumin suppresses local cyclooxygenase pathways, reducing glandular edema and facilitating milk letdown. Clinical evaluations of this specific botanical matrix demonstrate subclinical mastitis cure rates exceeding 87.5% within seven days, confirmed by rapid normalization of milk pH and precipitous drops in SCC⁶. Additionally, homeopathic formulations such as Phytolac are widely administered orally to cattle to combat systemic fever and mammary congestion, aligning perfectly with organic certification requirements prioritizing highly diluted phytotherapeutics over synthetic antibiotics².

Gastrointestinal Disorders and Neonatal Scours

Diarrhea (scours) in neonatal livestock represents a critical medical emergency. The primary etiology frequently involves rotavirus, coronavirus, or *Cryptosporidium parvum*, leading to acute enteritis and villous atrophy¹⁸. The ensuing pathology is defined not by the bacterial or viral agent itself, but by the catastrophic loss of extracellular fluid, resulting in hypovolemic shock and severe metabolic acidosis¹⁹.

Modern allopathic intervention prioritizes aggressive fluid therapy utilizing advanced oral rehydration solutions (ORS). Second-generation ORS formulations contain bicarbonate precursors, such as acetate and propionate, which consume hydrogen ions upon systemic metabolism to safely correct blood pH without alkalinizing the abomasum²⁰. Advanced fourth-generation solutions incorporate glutamine, an amino acid that fuels the regeneration of damaged enterocytes and actively facilitates sodium absorption across the compromised intestinal mucosa²⁰.

Within the ethnoveterinary framework, gastrointestinal inflammation is managed utilizing dual modalities: mucilaginous and botanical astringents. Plants rich in mucilage, such as flaxseed/linseed (*Linum usitatissimum*) and psyllium, contain complex polysaccharides that swell upon hydration²¹. These compounds coat the irritated intestinal epithelium, slow hypermotility, and physically bind bacterial endotoxins²¹. Conversely, astringent therapy relies upon the aggressive application of hydrolyzable and condensed tannins derived from sweet chestnut bark, tormentil root (*Potentilla erecta*), and oak bark (*Quercus* spp.)²². Administered as a concentrated decoction, these high-molecular-weight polyphenols bind tightly to the mucosal proteins in the gut lining. This interaction generates a coagulated, impermeable protective layer across the epithelium that instantly halts the exudation of bodily fluids and denies bacterial pathogens access to the underlying vasculature²².

The Toxicological Duality of Tannins: Oak and Acorn Poisoning

A foundational principle of veterinary pharmacognosy is the recognition of dose-dependent toxicity. While meticulously prepared decoctions of *Quercus* bark provide vital antidiarrheal action, the uncontrolled grazing of oak leaves and fallen acorns triggers severe, often fatal, systemic toxicosis in cattle and sheep²².

The pathogenesis of oak poisoning is driven exclusively by hydrolyzable tannins (HTs). Unlike condensed tannins, which resist cleavage, HTs are rapidly disassembled by ruminal microbes and enzymes into highly toxic, low-molecular-weight metabolites, principally gallic acid, ellagic

acid, and pyrogallol²⁶. Following absorption across the ruminal wall, these metabolites exert targeted destruction upon the renal tubular epithelium and vascular endothelium. This widespread cellular damage impairs protein reabsorption, causing acute cortical tubular necrosis marked by the precipitation of intratubular protein casts²⁶.

The clinical manifestation of natural acorn toxicosis includes profound depression, hemorrhagic gastroenteritis, polyuria transitioning to terminal anuria, and severe perirenal edema and ascites resulting from collapsed oncotic pressure²⁶. Crucially, species susceptibility varies drastically based on salivary biochemistry. Goats, pigs, and wild cervids are remarkably tolerant to oak ingestion because they secrete specialized tannin-binding salivary proteins (TBSPs)²⁶. These TBSPs form stable, insoluble complexes with the tannins in the oral cavity, effectively neutralizing them before ruminal degradation can occur²⁷. Cattle possess poorly developed TBSP pathways, rendering them acutely vulnerable²⁷. Preventative agronomic management involves restricting access to oak stands following severe storms that drop excessive mast, and supplementing exposed cattle with feed containing 10% hydrated lime (calcium hydroxide) or polyethylene-glycol (PEG), which serve as artificial tannin-inactivating agents within the rumen²⁷.

Dermatological Interventions and Hoof Care

For external lacerations, localized abscesses, and infectious pododermatitis (hoof rot), practitioners deploy a highly specific botanical materia medica. A globally recognized vulnerary agent is St. John's wort (*Hypericum perforatum*)³¹. The flowering aerial parts are subjected to prolonged maceration in olive oil, yielding a bioactive lipid matrix rich in hypericin and hyperforin³². This infused oil exerts profound anti-inflammatory, antimicrobial, and localized analgesic properties, proving highly effective when applied topically to deep dermal abrasions and necrotic hoof tissue³⁴.

Similarly, marigold (*Calendula officinalis*) is formulated into salves and poultices; the active flavonoids stimulate robust granulation tissue formation and accelerate re-epithelialization in slow-healing ulcerations³⁵. In traditional Indian ethnoveterinary practice, complex topical matrices are synthesized for severe foot lesions. A standard preparation requires the thorough grinding of aloe vera (*Aloe barbadensis*), garlic, turmeric, and curry leaves, which are subsequently boiled into sesame oil, filtered, and applied directly to the interdigital cleft to eradicate anaerobic bacterial colonization³⁷.

Alkaloid and Terpene Toxicology in Phytotherapy

Ethnoveterinary practitioners must remain vigilant against the inherent toxicological risks associated with specific botanical families, ensuring that remedies are processed and administered correctly.

Comfrey (*Symphytum officinale*): Historically utilized in topical poultices to reduce joint effusion and accelerate bone repair due to its high allantoin content, comfrey represents a severe hazard if administered orally⁶. The plant synthesizes up to 14 distinct pyrrolizidine alkaloids (PAs), including lasiocarpine and lycopsamine³⁸. Upon ingestion and subsequent hepatic metabolism, these PAs are converted into highly reactive pyrrolic esters that form permanent covalent adducts with hepatic DNA and structural proteins³⁸. This genotoxic

cascade irreversibly destroys liver endothelial cells, culminating in hepatic veno-occlusive disease (VOD), hepatocellular adenomas, and lethal organ failure³⁸. Consequently, its use is strictly relegated to topical application on unbroken skin.

Wormwood (*Artemisia absinthium*): Wormwood is extensively cited in traditional texts as a potent anthelmintic decoction for the expulsion of gastrointestinal parasites⁴¹. However, the

essential oil of *A. absinthium* contains varying concentrations of α -thujone and β -thujone, monoterpene ketones possessing intense neurotoxic capabilities⁴³. Thujone operates as a

non-competitive antagonist of the γ -aminobutyric acid type A (GABA-A) receptor⁴³. By blocking this primary inhibitory neurotransmitter pathway, high systemic doses of thujone trigger massive, unregulated central nervous system excitation⁴³. Clinically, this manifests as dose-dependent tonic-clonic seizures, severe tremors, and paralysis⁴³. Formulations must strictly monitor dosage to prevent crossing the neurotoxic threshold in livestock.

Fundamental Surgical Principles and Regional Anesthesia

Surgical intervention in livestock practice demands an uncompromising integration of anatomical knowledge, rigid asepsis, and localized chemical restraint. Due to the massive weight of the subjects and the severe physiological hazards associated with general anesthesia—specifically the risks of ruminal tympany, regurgitation, and aspiration pneumonia—the vast majority of bovine and small ruminant surgeries are executed on standing, sedated animals utilizing targeted regional nerve blocks⁴⁶.

Aseptic Field Preparation

Sterilization of surgical instruments and the operative field is paramount to prevent iatrogenic infections. Instruments must be autoclaved or subjected to high-level chemical disinfection. While ethanol and methanol solutions provide basic microbial reduction, extended exposure to glutaraldehyde (>2%) solutions is mandated for reliable sporicidal activity against resilient pathogens like *Bacillus* and *Clostridium*⁴⁸. The surgical site on the animal must undergo a rigorous, multi-stage aseptic preparation utilizing 2% to 4% chlorhexidine gluconate or povidone-iodine scrub⁵¹. Practitioners must strictly avoid the inappropriate use of povidone-iodine as an instrument soak, as it lacks necessary sporicidal efficacy for inanimate surfaces and frequently induces severe metallic corrosion on surgical steel⁵³.

Pharmacology and Toxicology of Local Anesthetics

Lidocaine hydrochloride (formulated as a 2% solution) remains the universal pharmacological standard for veterinary local anesthesia. The molecule functions by reversibly binding to and inhibiting voltage-gated sodium channels along the neuronal axon, thereby halting the influx of sodium ions and preventing the propagation of nociceptive action potentials toward the central nervous system⁴⁶.

The systemic absorption of lidocaine requires meticulous calculation, as toxic thresholds vary drastically across species. Overdose triggers a cascade of initial central nervous system

stimulation (twitching, nystagmus, convulsions) followed by catastrophic cardiopulmonary depression⁵⁴. In mature cattle, the maximum safe systemic dose is generously established at 8.0 to 10.0 mg/kg of body weight⁵⁵. Conversely, small ruminants (sheep and goats) exhibit an extreme sensitivity to local anesthetics due to rapid systemic absorption from highly vascularized injection sites⁵⁵. The absolute toxic threshold in caprines and ovines is reached at approximately 10 to 12 mg/kg, with profound neurotoxicity presenting rapidly⁵⁷. Therefore, the maximum clinical safe dose for sheep and goats must never exceed 4.0 to 6.0 mg/kg⁵⁵. If large physical volumes of anesthetic are required for extensive regional blocks in small ruminants, practitioners must dilute the standard 2% lidocaine solution 1:1 with sterile normal saline to yield a 1% solution, thereby providing adequate infiltration volume without breaching fatal toxic thresholds⁵⁵.

Regional Anesthesia Techniques

Targeted desensitization is achieved via specific neural blockade techniques, allowing for complex interventions without general anesthesia.

1. The Cornual Nerve Block (Dehorning and Disbudding): The horn corium and its surrounding integument are innervated by the cornual branch of the zygomaticotemporal nerve⁶⁰. In cattle, this nerve courses relatively superficially along the frontal crest. A single subcutaneous injection of 5 to 10 mL of 2% lidocaine deposited midway between the lateral canthus of the eye and the base of the horn, immediately beneath the temporal ridge, yields complete anesthesia within 10 minutes⁵⁵. Small ruminants present a more complex anatomical challenge; their horn base receives collateral sensory innervation from the infratrochlear nerve branching near the medial canthus, requiring a dual-site block or a complete subcutaneous ring block around the horn base to achieve total desensitization⁶⁰.

2. Flank Anesthesia: The Inverted L-Block and Paravertebral Blocks: For standing laparotomies, cesarean sections, and rumenotomies, the paralumbar fossa must be desensitized. The inverted L-block involves the serial injection of local anesthetic in a continuous, overlapping line bordering the dorsal and cranial margins of the proposed incision site⁶¹. While technically simple to execute, it requires a massive volume of lidocaine, risks localized tissue hematomas that delay healing, and frequently fails to provide total muscle relaxation of the deeper peritoneal layers⁴⁶.

Consequently, the thoracolumbar paravertebral nerve block is vastly superior. This technique aims to desensitize the dorsal and ventral branches of the T13, L1, and L2 spinal nerves, providing uniform analgesia and profound muscle relaxation of the entire flank⁴⁶. In the *proximal approach*, the needle is driven vertically at the cranial edge of the transverse processes of the lumbar vertebrae, depositing anesthetic directly at the intervertebral foramina where the nerve roots exit the spinal cord⁴⁶. In the *distal approach*, the nerves are blocked further down their anatomical path by fanning anesthetic dorsally and ventrally at the extreme lateral tips of the transverse processes of L1, L2, and L4⁴⁶.

3. Intravenous Regional Anesthesia (IVRA / Bier's Block): IVRA is the undisputed gold standard for performing digit amputations, complex hoof resections, and treating severe pododermatitis (hoof rot)⁶⁴. The traditional Bier's block involves placing a heavy pneumatic or

elastic tourniquet circumferentially around the mid-metacarpus or metatarsus to completely occlude both arterial inflow and venous return⁶⁴. A superficial vein—typically the dorsal common digital vein or radial vein—is catheterized, and lidocaine is injected retrograde, forcing the anesthetic backward against the venous valves⁶⁷. The drug diffuses rapidly out of the isolated venous bed into the surrounding neural tissue, providing profound analgesia and total motor akinesia of the distal limb within 10 to 15 minutes⁶⁵.

A significant clinical limitation of standard IVRA is the large total volume of lidocaine required, which predisposes the animal to systemic toxicity immediately upon tourniquet release, alongside the complication of severe tourniquet pain⁶⁴. To mitigate this, advanced surgical protocols utilize a modified **Distal IVRA (DIVRA)** technique. Here, the tourniquet is positioned significantly lower, immediately distal to the dewclaws⁶⁴. By utilizing the axial digital vein for infusion, the physical volume of the isolated vascular bed is drastically reduced. This allows surgeons to utilize exactly half the standard dose of lidocaine⁶⁴. To address the lack of postoperative pain control inherent to lidocaine, the anesthetic is frequently admixed with an

α_2 -agonist, such as xylazine, medetomidine, or dexmedetomidine⁶⁸. This combination achieves rapid sensory blockade while providing prolonged, intense analgesia long after the tourniquet is removed⁶⁸. Tourniquets must never be left in place for more than 60 to 90 minutes to prevent ischemic tissue necrosis, and deflation must be conducted gradually to prevent a sudden bolus of anoxic, anesthetic-laden blood from entering systemic circulation and causing cardiac arrhythmias⁶⁷.

Foundational Surgical Interventions

Rumenostomy and Trocarization

Acute ruminal tympany (bloat) is a catastrophic, rapidly progressive emergency. In instances of secondary free-gas bloat (frequently caused by a physical esophageal obstruction or vagal indigestion), the passing of an orogastric tube or the insertion of a sharp, large-bore trocar and cannula directly through the abdominal wall into the left paralumbar fossa can rapidly vent the trapped gas⁷⁰.

However, in primary pasture bloat, the gas is deeply entrapped in a highly stable, viscous foam generated by soluble leaf proteins and saponins⁷⁰. This dense foam will instantly clog a standard trocar⁷⁰. If the oral administration of chemical antifoaming agents (e.g., poloxalene, mineral oils) fails to break the surface tension and respiratory collapse is imminent due to diaphragmatic compression, an emergency rumenotomy or temporary rumenostomy is indicated⁷⁰.

A rumenostomy involves creating a semi-permanent, surgically controlled fistula to the exterior. Following a paravertebral or inverted L-block, a circular incision equal to the diameter of two ribs is made through the skin in the exact center of the left paralumbar fossa⁶¹. The surgeon dissects through the external abdominal oblique muscle, and then utilizes a blunt, muscle-splitting technique (gridding) through the fibers of the internal abdominal oblique and transversus abdominis⁶¹. This gridding aligns with the natural orientation of the muscle fibers, creating an endogenous sphincter or valve mechanism⁶¹. The underlying ruminal wall is grasped

with tissue forceps, exteriorized gently, and meticulously sutured directly to the cut skin edges⁶¹. This critical step prevents the highly contaminated, bacteria-rich ruminal fluid from leaking into the sterile peritoneal cavity, a technical failure that guarantees fatal peritonitis⁶¹. The stoma is allowed to heal by secondary intention, slowly contracting and closing naturally once the tympanic crisis has resolved⁶¹.

Abscess Excision and Drainage

Subcutaneous abscesses in livestock are frequently caused by pyogenic pathogens such as *Trueperella pyogenes*, *Corynebacterium pseudotuberculosis*, or *Staphylococcus aureus* following blunt trauma, vaccination site reactions, or deep puncture wounds⁷².

Surgical drainage should never be attempted until the abscess is fully encapsulated and mature. Maturity is indicated clinically by focal alopecia, localized heat, and fluctuance—a distinct, fluid-filled sensation generated by the degradation of tissue into purulent exudate upon deep palpation⁷². Prematurely lancing a hard, unorganized cellulitis mass is contraindicated, as it drives the pathogenic bacteria deeper into unprotected fascial planes⁷². Following the local infiltration of lidocaine around the periphery, a single vertical stab incision (3 to 5 cm) is made with a surgical scalpel at the absolute lowest (most dependent) point of the mass to ensure continuous, gravity-assisted drainage⁷². The purulent exudate is evacuated, and the cavity is aggressively lavaged⁷². Crucially, hydrogen peroxide must *never* be injected into a closed, deep abscess cavity; its potent oxidative mechanism is highly cytotoxic to newly forming fibroblasts and fragile endothelium, severely delaying healing and risking the creation of deep subcutaneous gas pockets⁷². Lavage is strictly performed using copious volumes of sterile, warm isotonic saline (0.9% NaCl) or extremely diluted antiseptic solutions (e.g., 0.05% chlorhexidine or a 1:10 dilution of povidone-iodine)⁷². The incision is never sutured closed; the cavity must remain open to drain and heal by secondary intention (from the inside out)⁷². Because puncture tracts and necrotic tissue cores present strictly anaerobic microenvironments, the administration of tetanus prophylaxis (toxoid or antitoxin) is a mandatory adjunct to surgical debridement to prevent the lethal proliferation of *Clostridium tetani*⁷⁴.

Ophthalmic Procedures: Subconjunctival Injections

Infectious Bovine Keratoconjunctivitis (IBK), commonly known as pinkeye, is caused by the bacterium *Moraxella bovis* and results in severe corneal ulceration, intense photophobia, and potential globe rupture⁷⁷. When the frequent application of topical ointments is impractical for large, extensively grazed herds, subconjunctival injections provide a highly effective method for the sustained, localized release of antibiotics (e.g., penicillin) and corticosteroids directly to the inflamed ocular tissues⁷⁷.

Following the application of a topical corneal anesthetic and physical restraint, a 25-gauge needle is carefully inserted either directly beneath the thin bulbar conjunctiva covering the sclera (epibulbar) or into the palpebral conjunctiva lining the inner surface of the upper eyelid⁷⁸. Approximately 1 mL of medication is slowly infused, creating a visible subconjunctival bleb⁷⁷. This reservoir of medication slowly diffuses across the corneal surface over several days⁷⁷. In

cases exhibiting severe, deep corneal ulceration threatening imminent perforation, surgical intervention is required. A temporary tarsorrhaphy (suturing the upper and lower eyelids together) or the surgical suturing of the third eyelid over the globe is performed to act as a living biological bandage, eliminating friction from blinking and drastically reducing UV light exposure⁷⁷.

Applied Principles of Surgical Suturing

The selection of the appropriate surgical suture pattern dictates the mechanical integrity of the closure, the distribution of tension, and the ultimate rate of tissue healing⁸³.

1. Inverting Patterns (Cushing and Lembert): When closing incisions in hollow, fluid-filled viscera—such as the rumen wall, abomasum, urinary bladder, or uterus—the tissue edges must be mechanically inverted (turned inward toward the lumen)⁸³. Inverting patterns achieve a flawless watertight seal, effectively burying the mucosal layer and preventing highly contaminated luminal contents from wicking along the suture line into the sterile peritoneal cavity⁸³.

- **The Cushing Pattern:** A continuous, running pattern that penetrates the serosa, muscularis, and submucosa, but deliberately *never* breaches the internal mucosal lumen. The needle is driven parallel to the incision line, crossing back and forth across the defect to roll the tissue inward over itself⁸⁴.
- **The Lembert Pattern:** Functionally similar to the Cushing, but the needle bites are taken perpendicular to the incision line, driving deeper into the tissue to provide superior holding power for organ walls under high tension⁸³. Both patterns minimize the amount of exposed suture material left on the serosal surface, which drastically reduces the foreign-body reaction and the formation of post-operative intra-abdominal adhesions⁸³.

2. Appositional Patterns (Ford Interlocking): For dermatological closures where exact anatomical alignment is required, appositional patterns are utilized⁸⁵. The Ford Interlocking pattern (often referred to as the blanket stitch) is a continuous pattern used extensively in large animal practice, specifically for closing heavy flank laparotomies⁸⁶. Before the surgeon pulls a new tissue bite tight, the needle is passed directly through the loop of the previous throw⁸⁷. This creates a robust locking mechanism running parallel to the entire incision line. This architecture distributes mechanical tension flawlessly across the thick bovine hide and ensures that if a single loop breaks or is rubbed open by the animal, the entire suture line does not rapidly unravel⁸⁵.

Conclusion

The successful execution of integrated livestock health management demands a practitioner capable of operating across multiple scientific disciplines. The foundation of this system rests upon recognizing that physiological stress, specifically nutritional deficits and metabolic imbalances like SARA and NEB, precipitates clinical disease. By deeply understanding the pharmacognosy of ethnoveterinary botanicals—leveraging the antimicrobial properties of the Lamiaceae family, the astringent power of hydrolyzable tannins, and respecting the acute toxicological dangers of plants like *Symphytum officinale* and *Artemisia absinthium*—practitioners can deploy highly effective, residue-free therapeutics that comply

with stringent organic regulations. When conservative phytotherapy and systemic support are insufficient, the precise execution of localized regional anesthesia, strict aseptic technique, and anatomically correct surgical intervention ensures the rapid restoration of animal health, safeguarding both welfare and productivity within modern agrarian systems.

Works cited

1. Thieme E-Journals - Homeopathy / Volltext,
<https://www.thieme-connect.com/products/ejournals/html/10.1055/s-0044-1781448?device=desktop&innerWidth=412&offsetWidth=412&lang=de>
2. EU law states categorically: “Homeopathy is First-Line-Medicine,
<https://edzardernst.com/2024/12/eu-law-states-categorically-homeopathy-is-first-line-medicine-really-no/>
3. Organic livestock farming | Request PDF - ResearchGate,
https://www.researchgate.net/publication/240382658_Organic_livestock_farming
4. Integrative medical approach to the antimicrobial resistance problem,
<https://cam-europe.eu/wp-content/uploads/2019/11/Integrative-medical-approach-to-the-antimicrobial-resistance-problem-Weiermayer-2019.pdf>
5. 43 Dual-Purpose Cattle Nutrition Plan.pdf
6. (PDF) Herbal Remedies for Bovine Mastitis: From Traditional,
https://www.researchgate.net/publication/400859721_Herbal_Remedies_for_Bovine_Mastitis_From_Traditional_Knowledge_To_Biotechnological_Validation
7. California Mastitis Test as a Screening Test - Academia.edu,
https://www.academia.edu/39743586/California_Mastitis_Test_as_a_Screening_Test
8. California Mastitis Test Procedure Guide | PDF - Scribd,
<https://www.scribd.com/document/889378056/California-Mastitis-Test>
9. Use of somatic cell counts and California mastitis test results from,
<https://avmajournals.avma.org/view/journals/javma/224/3/javma.2004.224.419.pdf>
10. Correlation between some direct and indirect tests for screen,
[http://www.ifrj.upm.edu.my/21%20\(03\)%202014/59%20IFRJ%2021%20\(03\)%202014%20Abd%20EI%20648.pdf](http://www.ifrj.upm.edu.my/21%20(03)%202014/59%20IFRJ%2021%20(03)%202014%20Abd%20EI%20648.pdf)
11. comparison of california Mastitis test and somatic cell counts for,
<https://www.jvas.in/article/comparison-of-california-mastitis-test-and-somatic-cell-counts-for-detection-of-subclinical-mastitis-in-crossbred-cattle/pdf>
12. Association between differential somatic cell count and California,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9709608/>
13. Threshold somatic cell count for delineation of subclinical mastitis,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6048081/>
14. Ethnoveterinary Medicine in Animal Disease Management - Vetscraft,
<https://www.vetscraft.com/ethnoveterinary-medicine-in-animal-disease-management/>
15. Review on Ethnoveterinary Medicinal Practice against Poultry,
[https://idosi.org/bjps/8\(2\)19/4.pdf](https://idosi.org/bjps/8(2)19/4.pdf)
16. salvia officinalis salvia: Topics by Science.gov,

- <https://www.science.gov/topicpages/s/salvia+officinalis+salvia>
17. Phytolac – Udder inflammation in cows – 1 L - Produits-Veto.com,
<https://en.produits-veto.com/produit/phytolac-boiron/>
 18. Calf Scour | NADIS,
<https://www.nadis.org.uk/disease-a-z/cattle/calf-management/calf-scour-beef-and-dairy-rearer-units/>
 19. Treatment of the Scouring Calf - Veterinary Partner - VIN,
<https://veterinarypartner.vin.com/default.aspx?pid=19239&catId=253974&id=4952596>
 20. Diarrhoea in calves and rehydration therapies - Vet Times,
<https://www.vettimes.com/news/vets/livestock/diarrhoea-in-calves-and-rehydration-therapies>
 21. Phytotherapy - Vetmeduni,
<https://www.vetmeduni.ac.at/en/donation-communication/phytotherapy>
 22. Traditional ethnoveterinary knowledge of indigestion or diarrhoea,
https://www.researchgate.net/publication/353659701_Traditional_ethnoveterinary_knowledge_of_indigestion_or_diarrhoea_treatments_in_cattle_on_the_Bilogora_hills_in_Croatia
 23. DB ETVET - ethnobotany, <https://www.ethnobotany.nl/db-etvet/>
 24. (PDF) Medicinal plants - Prophylactic and therapeutic options for,
https://www.researchgate.net/publication/303877066_Medicinal_plants_-_Prophylactic_and_therapeutic_options_for_gastrointestinal_and_respiratory_disease_in_Calves_and_piglets_A_systematic_review
 25. Tannin in Ruminant Nutrition: Review - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9738529/>
 26. Histopathologic and ultrastructural findings in oak (Quercus spp),
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11559815/>
 27. A Dry Fall and a Heavy Acorn Crop: Keep an Eye out for Acorn Toxicity,
<https://ucanr.edu/sites/default/files/2021-01/342444.pdf>
 28. Department of Animal Science - Plants Poisonous to Livestock,
<http://www.ansci.cornell.edu/plants/toxicagents/tannin.html>
 29. Acorn Toxicity | Helping farmers in Scotland - Farm Advisory Service,
<https://www.fas.scot/article/acorn-toxicity/>
 30. Acorn Poisoning in Cattle - American Angus Association,
https://www.angus.org/angus-media/angus-beef-bulletin/abb-extra/old-most-accessed/hn_acorn-poisoning-in-cattle
 31. hypericum perforatum treatment: Topics by Science.gov,
<https://www.science.gov/topicpages/h/hypericum+perforatum+treatment.html>
 32. Medicinal and local food plants in the south of Alava (Basque ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4675496/>
 33. An ethnoveterinary study on plants used for in the treatment of,
<https://or.niscpr.res.in/index.php/IJNPR/article/download/4216/1620/14740>
 34. medicinal plant hypericum: Topics by Science.gov,
<https://www.science.gov/topicpages/m/medicinal+plant+hypericum>
 35. Ethnoveterinary medicines used for ruminants in British Columbia,

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC1831764/>
36. Veterinary Herbal Medicine: A Systems-Based Approach - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7151902/>
37. Ethnoveterinary practices in animal care - Agriculture - Vikaspedia,
<https://en.vikaspedia.in/viewcontent/agriculture/best-practices/animal-husbandry/ethnoveterinary-practices-in-animal-care>
38. METABOLISM, GENOTOXICITY, AND CARCINOGENICITY OF,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC5894094/>
39. Hazards Associated with the Use of Herbal and Other Natural Products,
<https://veteriankey.com/hazards-associated-with-the-use-of-herbal-and-other-natural-products/>
40. Comfrey toxicity revisited - Eclectic School of Herbal Medicine,
<https://eclecticschoolofherbalmedicine.com/wp-content/uploads/2025/03/Comfrey-Toxicity-Revisited-Rhode.pdf>
41. MEDICINAL PLANTS USED IN TRADITIONAL VETERINARY,
https://veterinarymedicinejournal.usamv.ro/pdf/2020/issue_2/Art14.pdf
42. Low-Thujone A. absinthium L. (Wormwood) Essential Oils and,
<https://www.mdpi.com/1420-3049/31/10/1551>
43. Wormwood (Artemisia absinthium L.)--a curious plant with ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/20542104/>
44. Thujone: Psychedelic, Potent Cancer Treatment, or Poison? |,
<https://sites.evergreen.edu/plantchemeco/thujone-psychedelic-potent-cancer-treatment-or-poison/>
45. Wormwood toxicity - Botanical-online,
<https://www.botanical-online.com/en/medicinal-plants/wormwood-toxicity>
46. NIGERIAN VETERINARY JOURNAL - CABI Digital Library,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20230447430>
47. Local anesthesia in animals | PPTX - Slideshare,
<https://www.slideshare.net/slideshow/local-anesthesia-in-animals/230726627>
48. Sterilization of Surgical Tools: Removing Bacterial Endospores with,
<https://microbiologyjournal.org/sterilization-of-surgical-tools-removing-bacterial-endospores-with-a-combination-of-povidone-iodine-chlorhexidine-gluconate-ethanol-and-methanol/>
49. Sterilization of Surgical Tools: Removing Bacterial Endospores with,
<https://pdfs.semanticscholar.org/e05e/468bedbb01941089f46da0967dc9b8bd2bb7.pdf>
50. Disinfection of Healthcare Equipment | Infection Control - CDC,
<https://www.cdc.gov/infection-control/hcp/disinfection-sterilization/healthcare-equipment.html>
51. Aseptic technique and minor office surgery | Ento Key,
<https://entokey.com/aseptic-technique-and-minor-office-surgery/>
52. Disinfection and sterilisation procedures - VISION 2020 e-resource,
https://v2020eresource.org/content/files/Disinfection_Sterilisationprocedures.pdf
53. EQUIPMENT STERILIZATION DISINFECTION,
<https://ldh.la.gov/assets/oph/Center-PHCH/Center-CH/infectious-epi/EpiManual/>

[DisinfectionSterilization.pdf](#)

54. LOCAL ANESTHESIA/ANALGESIA,
<https://www.westernu.edu/mediafiles/veterinary/vet-anesthesia-analgesia/local-anesthesia-analgesia.pdf>
55. Local and Regional Anesthesia in Food Animals | Veterian Key,
<https://veteriankey.com/local-and-regional-anesthesia-in-food-animals/>
56. Local anesthesia – Large Animal Surgery – Supplemental Notes,
<https://open.lib.umn.edu/largeanimalsurgery/chapter/local-anesthesia/>
57. Toxicity and Pharmacokinetic Studies of Lidocaine and Its Active,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6115778/>
58. ANESTHESIA AND PAIN MANAGEMENT IN SMALL RUMINANTS,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20163308908>
59. Anaesthesia of sheep, goats, and other herbivores | Veterian Key,
<https://veteriankey.com/anaesthesia-of-sheep-goats-and-other-herbivores/>
60. Nerve blocks and dehorning- By: Dr. Najmu Saaqib Reegoo DVM,
<https://www.slideshare.net/slideshow/nerve-blocks-and-dehorning-by-dr-najmu-saaqib-reegoo-dvm/140154730>
61. How to – Rumenostomy – Large Animal Surgery,
<https://open.lib.umn.edu/largeanimalsurgery/chapter/how-to-rumenostomy/>
62. Veterinary Anesthesia Overview Guide | PDF | Anesthesia | Medical,
<https://www.scribd.com/document/840642380/Anesthesia-DVM>
63. Local and regional anesthesia 2019 5 lecture,
https://www.uomosul.edu.iq/public/files/datafolder_117/oldUo_20190821_033057_28_524.pdf
64. Comparison of lidocaine and its combination with ketamine for distal,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20230488533>
65. (PDF) Evaluation of intravenous regional anaesthesia and four-point,
https://www.researchgate.net/publication/320914755_Evaluation_of_intravenous_regional_anaesthesia_and_four-point_nerve_block_efficacy_in_the_distal_hind_limb_of_dairy_cows
66. Effects of adding atracurium to Lidocaine solution during ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12330004/>
67. Foot and Claw Surgery - IHMC Public Cmaps (3),
<https://cmapspublic3.ihmc.us/rid=1QRBKTP2Q-1D6WT6P-1DB7/Drugs%20and%20Anaesthesia.pdf>
68. Buffalo Bulletin (October-December 2021) Vol.40 No.4 - KUOJS,
<https://kuojs.lib.ku.ac.th/index.php/BufBu/article/view/3260/2233>
69. (PDF) Evaluation of Intravenous Regional Anesthesia of Lidocaine,
https://www.researchgate.net/publication/341412794_Evaluation_of_Intravenous_Regional_Anesthesia_of_Lidocaine_Medetomidine_and_Lidocaine-medetomidine_in_the_Hind_Limb_of_Goats
70. Bloat in Ruminants - Digestive System - Merck Veterinary Manual,
<https://www.merckvetmanual.com/digestive-system/diseases-of-the-ruminant-f-orestomach/bloat-in-ruminants>
71. Surgery of the Forestomach | Request PDF - ResearchGate,

- https://www.researchgate.net/publication/308910189_Surgery_of_the_Forestomach
72. Skin Abscesses in Livestock and Deer: Description and Treatment, https://www.researchgate.net/publication/404080499_Skin_Abscesses_in_Livestock_and_Deer_Description_and_Treatment_WEC483UW552_42026
 73. Skin Abscesses in Livestock and Deer: Description and Treatment, <https://ask.ifas.ufl.edu/publication/UW552>
 74. Wound Debridement and Lavage Techniques | Dr. Zubair Khalid, <https://www.zubairkhalid.com/knowledge/veterinary-medicine/veterinary-surgery/wound-debridement-lavage-techniques>
 75. Tetanus - Anipedia - University of Pretoria, <https://anipedia.up.ac.za/resources/tetanus/1156.html>
 76. Tetanus wound care and prophylaxis - Starship, <https://www.starship.org.nz/guidelines/tetanus-prophylaxis/>
 77. Manage Pinkeye - FlyBoss, <https://flyboss.com.au/manage-pinkeye/>
 78. How to – Subconjunctival injection – Large Animal Surgery, <https://open.lib.umn.edu/largeanimalsurgery/chapter/how-to-subconjunctival-injection/>
 79. Subconjunctival Injection in eye || Pink Eye Treatment in cow, <https://www.youtube.com/watch?v=yiwsHaXeY1U>
 80. Managing and Preventing Pinkeye - Livestock, <https://livestock.extension.wisc.edu/articles/managing-and-preventing-pinkeye/>
 81. Pinkeye in cattle: Part Three - Beef - MSU Extension, https://www.canr.msu.edu/news/pinkeye_in_cattle_part_three
 82. Managing pinkeye in your herd - Greenham, <https://www.greenham.com.au/managing-pink-eye-in-your-herd/>
 83. Inverting suture patterns – Large Animal Surgery, <https://open.lib.umn.edu/largeanimalsurgery/chapter/suturing-inverting-suture-patterns/>
 84. Cushing Pattern - Veterinary Surgery Online, <https://www.vetsurgeryonline.com/cushing-pattern/>
 85. Overview of Surgical Suture Patterns | PDF - Scribd, <https://www.scribd.com/document/708689946/surgical-suture-patterns>
 86. Skin Closure Methods in Small Animal Surgery - Sustainable Vet, <https://www.sustainablevet.org/blog/skin-closure-methods-small-animal-surgery>
 87. Ford Interlocking Pattern - Veterinary Surgery Online, <https://www.vetsurgeryonline.com/ford-interlocking-pattern/>
 88. Sutures | PPTX - Slideshare, <https://pt.slideshare.net/slideshow/sutures-85859065/85859065?nway-refresh=B>