

Thermodynamic, Psychrometric, and Physiological Management of Subterranean Crop Storage in Temperate Climates

The engineering and operation of earth-sheltered storage facilities—commonly referred to as root cellars—demand a rigorous interdisciplinary synthesis of geotechnical engineering, atmospheric thermodynamics, and postharvest plant physiology. The ultimate objective of such a facility is to decouple the internal storage environment from the extreme diurnal and seasonal fluctuations of the surface climate, establishing a strictly regulated microclimate capable of preserving harvested biological tissues in a state of suspended senescence. This objective is complicated by the highly heterogeneous environmental requirements of different agricultural commodities. For instance, subterranean storage must frequently accommodate root crops that require near-saturation humidity (90–95% relative humidity) to prevent cellular desiccation, alongside allium crops that mandate significantly drier conditions (50–70% relative humidity) to arrest aggressive fungal proliferation¹.

Achieving these divergent psychrometric parameters within a single continuous subterranean airspace, particularly in a temperate maritime-transitional climate such as Berlin, Germany, requires a highly nuanced approach. To successfully operate a multi-zone earth-sheltered crop storage facility, designers and agricultural engineers must synthesize architectural zoning, passive aerodynamics, boundary layer manipulation, and matric potential management. This report exhaustively analyzes the physical, thermal, and biological principles required to successfully execute and regulate high-performance subterranean crop storage.

1. Geothermal Heat Transfer and Subterranean Thermodynamics

To maintain a stable internal temperature of $0-4^{\circ}\text{C}$ ($32-40^{\circ}\text{F}$), a subterranean structure must operate as an active thermodynamic system that meticulously balances the geothermal heat flux of the surrounding regolith against the convective cooling driven by atmospheric ventilation³. The foundational principle is the exploitation of the earth's immense thermal mass, which forces high-frequency surface temperature fluctuations to propagate downward as heavily damped, phase-shifted thermal waves.

1.1 The Kasuda Equation and Thermal Wave Attenuation

Heat transfer through a porous particulate medium, such as the sandy loams native to the Berlin-Brandenburg region, is governed by the principles of thermal conduction. The mathematical prediction of soil temperature at any specific depth and time of year relies on the analytical model formulated by Kasuda and Archenbach, derived from the one-dimensional

Fourier heat conduction equation³. The subsurface temperature, $T_{soil}(z, t)$, is mathematically expressed as:

$$T_{soil}(z, t) = T_{mean} - T_{amp} \cdot \exp\left(-z\sqrt{\frac{\pi}{\alpha\tau}}\right) \cdot \cos\left(\frac{2\pi}{\tau}(t - t_{shift})\right)$$

Where T_{mean} is the mean annual surface temperature, T_{amp} is the amplitude of the annual surface temperature wave, z is the depth below the surface in meters, α is the thermal diffusivity of the soil, τ is the period of the temperature cycle (typically 365 days), t is the current day of the year, and t_{shift} is the day of the year corresponding to the minimum surface temperature³.

The exponential term in this equation dictates amplitude attenuation. Due to the high thermal inertia of the soil, short-period diurnal temperature variations are heavily diminished as depth increases, effectively vanishing beyond a depth of 0.5 to 1.0 meters³. The cosine term dictates the temporal phase lag. In a temperate climate like Berlin, the deep-earth steady-state temperature stabilizes near the regional mean annual air temperature (approximately 9–10°C). Because root crops require storage temperatures strictly near 0–4°C to suppress postharvest respiration, the cellar cannot be buried at an infinite depth¹. Instead, it must be positioned strategically within the shallow active zone (typically 2 to 4 meters deep) to capture the extreme phase-shifted cold of the winter exactly as it arrives in mid-summer, while relying on heavy artificial surface insulation to truncate the downward propagation of the summer heat wave³.

1.2 Soil Thermal Conductivity and the Johansen Method

The velocity and penetration depth of the thermal wave are entirely dependent on the thermal diffusivity (α) of the soil matrix, mathematically defined as the ratio of thermal conductivity (λ) to volumetric heat capacity (C_v):

$$\alpha = \frac{\lambda}{C_v} = \frac{\lambda}{\rho_b \cdot c_p}$$

Soil is a multiphase system consisting of solid mineral particles, liquid water, and interstitial air. Because air possesses a thermal conductivity approximately 20 times lower than that of liquid

water, the volumetric water content (θ) drastically alters the thermodynamic properties of the soil matrix³. In the sandy and loamy soils typical of the Berlin area, thermal diffusivity exhibits a highly non-monotonic relationship with soil water content⁶.

To accurately model this behavior, geotechnical engineers utilize the Johansen method, which calculates the thermal conductivity of the soil as a linear interpolation between its completely dry (λ_{dry}) and completely saturated (λ_{sat}) states, weighted by the dimensionless Kersten number (K_e)³:

$$\lambda = \lambda_{dry} + (\lambda_{sat} - \lambda_{dry})K_e$$

As a dry soil begins to wet, conductivity increases much faster than heat capacity, causing diffusivity to rise sharply. This parameter typically reaches its absolute maximum at a moderate volumetric water content (around 18% to 25%, depending on the specific textural classification of the loam or sand)³. Beyond this threshold, as the soil approaches total saturation, the relative increase in heat capacity overtakes the increase in conductivity, causing the thermal diffusivity to slowly decline³. If a root cellar is constructed in a high-diffusivity medium (e.g., highly compacted, saturated sand), it will suffer from rapid thermal bridging to the surface. In such environments, deeper excavation and significantly augmented engineered insulation become mandatory to preserve the target 0–4°C microclimate³.

1.3 Structural Loads, Hydrostatic Uplift, and Hydrological Defenses

The burial of a rigid structure beneath several meters of earth exposes it to continuous, immense gravitational and geotechnical forces. A flat-roofed subterranean structure must support the static vertical dead load of the backfilled soil, which for a typical compacted loam equates to approximately 120 to 135 pounds per cubic foot (18.8 to 21.2 kN/m^3)³. The lateral forces exerted by the soil against the cellar walls are calculated using the At-Rest Earth Pressure coefficient ($K_0 \approx 1 - \sin(\phi')$), which treats the soil mass as an equivalent fluid pressure forming a triangular load distribution³.

Compounding these structural loads is the perpetual threat of capillary moisture and hydrostatic pressure. The preeminent international framework for addressing external hydrological threats is the European DIN 18533 standard: *Waterproofing of Components in Contact with the Ground*³. For a root cellar constructed in well-draining soils with a functional perimeter subsurface trench drain, the vertical walls operate under Class W1-E (soil moisture and non-accumulating seepage water), while the earth-covered roof operates under W3-E (non-pressing water on earth-covered ceilings)³.

Compliance with DIN 18533 for the W3-E roof mandates a two-layer bituminous or single-layer elastomeric membrane³. Materials such as EPDM rubber must be utilized and explicitly certified

as root-resistant in accordance with DIN EN 13948 to prevent biological degradation and root penetration from topsoil flora³. To successfully decouple the cellar from the thermal bridging of wet soil, engineers deploy "umbrella insulation" systems using cellular glass (e.g., FOAMGLAS or GLAPOR). Cellular glass offers extraordinary compressive strengths (ranging from 0.5 to 2.4 MPa), is completely vapor-tight, and remains fundamentally impervious to rodent tunneling and biological decay over the multi-generational lifespan of the structure³. Furthermore, adherence to the BS EN 1997-1 UPL (Uplift) limit state protocol is required to ensure the upward buoyant force of the displaced groundwater does not exceed the factored downward dead weight of the concrete structure, a failure mode that results in catastrophic foundation fracturing³.

2. Postharvest Physiology and Aerobic Respiration

Once isolated from external thermal and hydrological extremes, the internal environment must be strictly regulated to meet the physiological demands of the stored organic tissues. Fresh fruits and vegetables are living organisms that continue to undergo aerobic respiration postharvest. This metabolic process consumes oxygen and energy reserves (primarily sugars

and starches) while releasing metabolic heat, carbon dioxide (CO_2), water vapor, and potent phytohormones such as ethylene¹.

The primary objective of subterranean storage is to reduce the ambient temperature to the lowest safe threshold without inducing chilling injury, thereby suppressing the respiration rate to a baseline "idle"¹. The relationship between temperature and respiration is characterized by

the Q_{10} temperature coefficient, a measure of the rate of reaction change resulting from a 10°C temperature shift¹⁷. Within the standard $5\text{--}25^\circ\text{C}$ range, the Q_{10} value for most root

crops is between 2.0 and 2.5, meaning that for every 10°C decrease in temperature, the respiration rate is halved, significantly extending shelf life and preserving nutritional density¹⁷.

The rate of heat generated by this process is measured as the Heat of Respiration, typically expressed in British thermal units (Btu) per ton per day. To accurately calculate the necessary passive or active cooling loads for a facility, engineers consult reference materials such as the United States Department of Agriculture (USDA) Agriculture Handbook 66: *The Commercial Storage of Fruits, Vegetables, and Florist and Nursery Stocks*⁹. For example, winter squash

stored at 10°C produces an immense 5,000 to 7,500 Btu/ton/day, whereas topped carrots stored at 0°C generate a highly manageable 800 to 1,400 Btu/ton/day¹⁸.

Commodity	Optimal Temperature ($^\circ\text{C}$ / $^\circ\text{F}$)	Relative Humidity (%)	Expected Storage Life	Heat of Respiration (0°C , Btu/ton/day)	Chilling / Freezing Sensitivity Threshold

Carrots (topped)	0 / 32	98–100%	6–8 months	800–1,400	Freezes at -1.2°C (29.8°F) ¹
Beets (topped)	0 / 32	98–100%	4–6 months	1,200–1,900	Freezes at -0.9°C (30.3°F) ¹
Onions (dry)	0 / 32	65–70%	6–9 months	900–2,500	Freezes at -0.8°C (30.6°F) ²
Garlic	0 / 32	65–70%	5–8 months	900–3,100	Freezes at -0.8°C (30.5°F) ²
Cabbage (late)	0 / 32	98–100%	5–6 months	1,000	Freezes at -0.9°C (30.4°F) ¹⁶
Pumpkins / Winter Squash	10–15 / 50–60	50–70%	2–6 months	5,000–7,500	Highly chilling sensitive; degrades rapidly below 10°C [cite: 2, 18, 21]
Potatoes (Late / Processing)	4.5–13 / 40–55	90–95%	5–10 months	800–2,000	Below 4°C undergoes cold-induced sweetening

					(CIS) ²
--	--	--	--	--	--------------------

Table 1: Postharvest physiological parameters, psychrometric requirements, and thermal generation data for common temperate storage crops, synthesized from USDA Agriculture Handbook 66¹.

3. The Humidity Conflict: Psychrometric Requirements of Divergent Crops

While temperature requirements across temperate winter crops are relatively uniform—clustering tightly around the 0°C (32°F) mark to suppress respiration without inducing ice nucleation—their relative humidity (RH) requirements are fundamentally incompatible.

Root vegetables, such as carrots, beets, turnips, and parsnips, possess thin epidermal layers that offer negligible boundary layer resistance to moisture diffusion¹. If the ambient air is not near saturation, a high Vapor Pressure Deficit (VPD) is established between the water-rich internal tissue of the root and the surrounding atmospheric air, driving rapid transpiration²⁵. Wilting symptoms in carrots become evident with as little as a 3% total weight loss²⁸. To prevent this tissue desiccation, the USDA Handbook 66 strictly mandates storage conditions of 98–100% relative humidity for topped carrots and beets, ensuring tissue turgor and crispness are maintained over multi-month storage periods¹.

Conversely, alliums such as onions and garlic require an entirely different psychrometric environment. While they share the 0°C temperature optimum to arrest sprouting, they demand a much drier atmosphere, strictly maintained between 65% and 70% RH². Postharvest, onions are purposefully cured at elevated temperatures (24–27°C) and moderate humidity (70–80% RH) for approximately two weeks prior to cold storage³⁰. This curing process accelerates the deposition of suberin—a complex, waxy, highly hydrophobic polymer—and lignin within the cell walls of the damaged neck tissue, forming a hardened wound periderm³². This suberized tunic seals the bulb against excessive moisture loss and serves as a physical barrier against necrotrophic pathogens³⁵. Prolonged exposure to the 90–100% RH environment required by root crops destroys this dry tunic, re-hydrating the outer layers and providing the necessary moisture for the germination of fungal spores, inevitably inducing catastrophic decay³⁰.

4. Psychrometric Dynamics of Winter Air and Passive Ventilation

A critical requirement of subterranean storage is the continuous exhaustion of metabolic gases—namely CO₂ and ethylene—and the cyclic replenishment of oxygen. While active commercial agricultural facilities employ mechanical fans to achieve 30 to 60 Air Changes per Hour (ACH), passive root cellars rely entirely on aerodynamic phenomena: primarily

buoyancy-driven stack effect ventilation, supplemented by wind-driven cross ventilation³.

4.1 Buoyancy-Driven Stack Effect Ventilation

The stack effect is driven by the density differential between the colder, denser outdoor ambient air and the relatively warmer, humid air inside the cellar. A low intake pipe introduces dense air near the floor, displacing the lighter, stale air, which escapes through a high exhaust pipe located at the apex of the vaulted ceiling³. The Neutral Pressure Level (NPL) establishes the specific elevation where indoor and outdoor absolute pressures are equal; below this point the enclosure is negatively pressurized (drawing air in), while above it, it is positively pressurized (forcing air out)³.

The volumetric airflow rate (Q) driven by thermal buoyancy is calculated using the standard ASHRAE thermal force equation, integrating Bernoulli's principle for incompressible flow:

$$Q = C_d \cdot A \cdot \sqrt{2g \cdot \Delta H \cdot \frac{T_i - T_o}{T_i}}$$

Where C_d is the discharge coefficient of the duct opening (typically 0.60 to 0.65), A is the cross-sectional free area of the ventilation pipe, g is gravitational acceleration (9.81 m/s^2), ΔH is the vertical stack height (the distance between the intake and exhaust midpoints), and T_i and T_o are the absolute indoor and outdoor temperatures in Kelvin, respectively³. By artificially raising the exterior exhaust stack well above ground level, the height differential (ΔH) is maximized. Because airflow is proportional to the square root of the height differential ($\sqrt{\Delta H}$), this exponentially increases the draft, allowing for the necessary 2 to 4 ACH required to safely flush ethylene gas without mechanical intervention³.

4.2 The Winter Psychrometric Paradox and the Mollier Diagram

The management of ventilation in a temperate climate like Berlin introduces a severe, counter-intuitive psychrometric challenge during the winter months. To mathematically understand this phenomenon, engineers consult the Mollier $h - x$ diagram, the European standard psychrometric chart plotting enthalpy against specific humidity⁴⁰. Relative humidity (RH) is inversely proportional to temperature; warm air is capable of holding exponentially more water vapor than cold air⁴³. During a typical Berlin winter, the outdoor air may register at 0°C to 5°C with a very high relative humidity (e.g., 85% RH)⁴⁵. However, the absolute moisture content (specific humidity or humidity ratio, expressed in grams of water per kilogram of dry air) of this frigid air is extremely low⁴⁵.

When this cold outdoor air is drawn into the root cellar via the stack effect, it is warmed by the massive geothermal inertia of the earth-sheltered walls to the target 4°C (or higher, depending on localized heat of respiration). As the air warms, its capacity to hold moisture expands rapidly, but the absolute mass of water vapor remains entirely constant. Consequently, the relative humidity of the air drops precipitously⁴³. If unchecked, winter ventilation acts as a powerful passive dehumidifier, stripping moisture from the internal environment and aggressively driving the Vapor Pressure Deficit higher. The latent heat of evaporation required to vaporize water from the stored crops further cools the room, but the primary threat is the desiccation of the produce⁴⁷.

This physical reality dictates the architectural necessity of isolated microclimates. Because the macro-environment of the cellar inevitably dries out during winter ventilation—often naturally settling around the 65–70% RH ideal for alliums—the high-humidity root crops must be artificially shielded from the ambient air²³.

5. Microclimate Zoning via Damp Sand Matrices

To reconcile the paradox of requiring a 65% RH atmosphere for onions and a 98% RH atmosphere for carrots within the same physical space, the macro-environment of the cellar must be operated at the lower humidity threshold (65–70% RH). The high-humidity crops are then isolated within localized microclimates by packing them in a damp particulate matrix—specifically, washed quartz or river sand⁵⁰.

5.1 The Physics of Soil Water Retention and Matric Potential

The efficacy of damp sand relies on the principles of soil water potential (Ψ_t), which dictates the thermodynamic energy required for a plant tissue to extract or lose water to its surrounding environment. The total water potential is the sum of its constituent physical forces:

$$\Psi_t = \Psi_g + \Psi_m + \Psi_o + \Psi_h$$

Where Ψ_g is gravitational potential, Ψ_m is matric potential, Ψ_o is osmotic potential, and Ψ_h is hydrostatic pressure⁵³. Within the confines of a storage crate, the matric potential (Ψ_m)—driven by capillary action and adsorptive forces between the water molecules and the solid sand particles—dominates the fluid interaction⁵³.

By burying carrots entirely in damp sand, the roots are shielded from the atmospheric Vapor Pressure Deficit (VPD) of the macroscopic cellar space. The damp sand acts as a sacrificial moisture buffer; as the dry 65% RH air of the cellar passes over the storage crates, water evaporates from the outer layers of the sand matrix rather than from the delicate cellular tissues of the carrots²³.

The behavior of water within this matrix is governed by the Soil Water Characteristic Curve (SWCC), which illustrates the non-linear relationship between matric suction and volumetric

water content⁵⁷. Quartz sand is characterized by large macropores, meaning it possesses a very low air-entry value (bubbling pressure) and releases water at relatively low matric suctions compared to clays or silts⁶⁰. This low air-entry value allows the roots to easily absorb trace amounts of moisture from the sand to maintain turgidity, preventing the rubbery, shriveled texture that inevitably occurs when roots are exposed to dry ambient air⁵². Furthermore, hysteresis in the SWCC—caused by the "ink-bottle" effect and contact angle variations during the draining and re-wetting of the pores—ensures that the sand retains varying degrees of moisture depending on its wetting history, stabilizing the microclimate against rapid fluctuations⁶³.

5.2 Oxygen Diffusion Rates (ODR) and Aerobic Constraints

While the sand matrix must be damp, it must absolutely not reach saturation. Stored root vegetables are living tissues that continue to respire, taking in ambient oxygen and expelling carbon dioxide. If the volumetric water content of the sand is too high, water displaces the air within the interstitial pore spaces. Because oxygen diffuses approximately 10,000 times slower through liquid water than through air, over-wetting the sand causes a catastrophic drop in the Oxygen Diffusion Rate (ODR) surrounding the root epidermis⁶⁶.

When the roots are deprived of oxygen (tissue anoxia), they undergo a rapid metabolic shift from aerobic oxidative phosphorylation to anaerobic fermentation to survive⁶⁶. This emergency pathway converts pyruvate into acetaldehyde and ultimately ethanol via the enzyme alcohol dehydrogenase (ADH), yielding minimal ATP, generating toxic byproducts, and rapidly leading to cellular death and tissue maceration⁶⁶.

Furthermore, waterlogged, anaerobic conditions in the sand matrix provide the ideal proliferation environment for *Pectobacterium carotovorum* (formerly *Erwinia carotovora*), the primary causal agent of bacterial soft rot⁷⁰. *P. carotovorum* relies on excessive moisture and low oxygen thresholds to multiply, secreting powerful pectinolytic cell wall-degrading enzymes that rapidly liquefy the carrot tissue into a foul-smelling slime⁷⁰. Therefore, the packing matrix must be a highly porous, well-washed quartz sand (e.g., 0.4–2.0 mm grain size complying with EN 12904 standards) that drains freely, maintaining a high air-filled porosity while preserving only a thin film of capillary moisture around the individual particles⁷⁴. Organic matrices like peat moss or sawdust, while occasionally referenced in lay literature, are highly susceptible to compacting, waterlogging, and imparting severe off-flavors to the stored produce²³.

6. Pathological and Physiological Failure Modes

Failure to rigorously manage the thermodynamic, psychrometric, and spatial parameters of the cellar inevitably results in catastrophic loss via physiological disorders or aggressive phytopathological infections. The biological interactions between different species of stored produce dictate strict atmospheric segregation protocols.

6.1 Ethylene Cross-Contamination and Isocoumarin Synthesis

Ethylene (C_2H_4) is a highly potent gaseous phytohormone naturally produced by ripening climacteric fruits (e.g., apples, pears, tomatoes) and by rotting organic matter³. While it accelerates senescence and sprouting in potatoes, its physiological effect on carrots is

particularly devastating and irreversible.

When carrots are exposed to ethylene concentrations as low as 0.1 to 0.5 parts per million (ppm) at low temperatures, it triggers a severe biochemical defense response⁷⁸. The presence of ethylene induces the condensation of acetyl-CoA and malonyl-CoA molecules, leading to the rapid synthesis and accumulation of 8-hydroxy-3-methyl-6-methoxy-3,4-dihydro-isocoumarin (commonly referred to as isocoumarin or 6-methoxymellein) in the peel and phloem tissue of the root⁷⁹. Isocoumarin is a phytoalexin that imparts an intensely bitter, turpentine-like off-flavor, rendering the carrots entirely unpalatable and resulting in total commercial or nutritional loss⁷⁸. Therefore, high-ethylene producers like apples must either be rigidly excluded from the cellar entirely or isolated in hyper-ventilated, vapor-tight sub-compartments, relying heavily on the continuous passive stack effect to scavenge and exhaust ambient ethylene³.

In apples, the accumulation of ethylene and the subsequent oxidation of α -farnesene into conjugated trienols (CTs) within the peel is the primary mechanism behind superficial scald, a browning disorder that ruins the fruit's marketability⁸³.

6.2 *Botrytis aclada* and the Necessity of *Allium* Suberization

While onions require the low RH (65–70%) macro-environment created by winter ventilation, they remain highly susceptible to *Botrytis aclada* (syn. *B. allii*), the causal agent of onion neck rot³⁰. *Botrytis aclada* is an opportunistic necrotrophic fungus that often remains entirely asymptomatic in the field but manifests aggressively during storage if the relative humidity exceeds the 70–75% threshold, or if free condensation forms on the bulb surface⁸⁷.

The primary defense against *Botrytis* penetration is not chemical, but physical. Postharvest,

onions must be subjected to a rigorous curing period (typically 2 weeks at $24\text{--}27^{\circ}\text{C}$ / $75\text{--}80^{\circ}\text{F}$ and 70–80% RH) before being introduced into the cold 0°C cellar environment³⁰. As previously established, this curing process accelerates the deposition of a suberin and lignin wound periderm³². If uncured onions are stored, or if the macro-humidity of the cellar rises above 75%, *B. aclada* easily colonizes the succulent neck tissue, softening the scales and producing a dense grey conidial mass, eventually resulting in the complete liquefaction of the bulb from the neck downward³⁹.

6.3 *Sclerotinia sclerotiorum* and Condensation Management

Within the high-humidity microclimate of the damp sand, root crops face the constant threat of *Sclerotinia sclerotiorum*, the fungal pathogen responsible for white mold (sclerotiniosis)⁹⁰. *S. sclerotiorum* thrives in the 95–100% RH zones, producing a cottony white mycelial mat embedded with hard black sclerotia. This fungus rapidly spreads laterally through the sand matrix from infected to healthy roots, causing a watery soft rot that can decimate a densely packed crate in weeks⁹⁰.

The risk of fungal and bacterial proliferation throughout the cellar is exponentially magnified by the presence of free liquid water. The thermodynamic relationship governing condensation is precisely modeled by the Magnus-Tetens equation, which calculates the dew point

temperature (T_d) based on ambient dry-bulb temperature (T) in Celsius and relative humidity (RH)³:

$$\gamma = \frac{17.625 \cdot T}{243.04 + T} + \ln \left(\frac{RH}{100} \right)$$

$$T_d = \frac{243.04 \cdot \gamma}{17.625 - \gamma}$$

For a cellar successfully maintained at a dry-bulb temperature of 4°C with a relative humidity of 90%, the dew point is approximately 2.5°C ³. If any interior surface—such as an uninsulated flat concrete ceiling exposed to surface frost conduction—drops to this threshold temperature, catastrophic condensation will spontaneously form³. Condensation dripping vertically onto stored crops provides the exact film of free water required for the germination of fungal spores. In apples, this moisture facilitates *Penicillium expansum* infections, which lead to blue mold decay and the subsequent accumulation of patulin, a highly toxic and tightly regulated mycotoxin⁹⁷.

To geometrically mitigate this severe psychrometric risk, subterranean architecture must avoid flat roof spans. Integrating a vaulted or arched ceiling ensures that any inevitable condensation adheres to the masonry via capillary action and surface tension. The moisture safely sheds down the curvature of the arch to the vertical walls, bypassing the biological cargo entirely and draining harmlessly into the gravel sub-floor³.

6.4 Sprout Suppression in Potatoes

While potatoes share the $4.5\text{--}13^{\circ}\text{C}$ storage requirements of other root crops, storing them at the lower end of this spectrum (e.g., $2\text{--}4^{\circ}\text{C}$) to indefinitely delay sprouting induces a detrimental physiological response known as Cold-Induced Sweetening (CIS)². During CIS, the starch within the tuber is rapidly degraded into reducing sugars (glucose and fructose), rendering the potato unpalatable and causing severe browning (Maillard reaction) if processed or fried²².

Therefore, potatoes destined for culinary consumption must be stored at warmer temperatures ($8\text{--}12^{\circ}\text{C}$), which inevitably breaks natural dormancy and encourages sprouting²². To combat this, facility managers must employ sprout suppressants. While the synthetic herbicide chlorpropham (CIPC) has been the historical standard, regulatory bans in

regions like the European Union due to its toxic degradation products (e.g., 3-chloroaniline) have necessitated the transition to safer alternatives¹⁰¹. Modern, compliant sprout suppression relies on the application of 1,4-dimethylnaphthalene (1,4-DMN), a naturally occurring potato hormone analog, or the use of essential oils such as carvone (derived from caraway or mint), which inhibit sprout elongation by downregulating specific hormonal pathways without compromising tuber viability or human health¹⁰¹.

7. Conclusion

The long-term preservation of agricultural harvests within an earth-sheltered subterranean structure requires an exact synthesis of geotechnical engineering, fluid dynamics, and postharvest physiology. In temperate maritime-transitional climates such as Berlin, the strategic utilization of the Kasuda equation allows engineers to embed the structure at a depth that decouples the facility from surface temperature extremes. By capturing a persistent

0–4°C geothermal baseline and defending it with high-compressive cellular glass insulation, postharvest aerobic respiration is effectively suppressed.

However, the biological reality of mixed-crop storage demands conflicting humidity parameters. Root crops require near-saturation environments (98% RH) to stave off extreme vapor pressure deficits and lethal wilting, while alliums and fruits demand drier atmospheres (65% RH) to prevent catastrophic fungal colonization by *Botrytis aclada* and *Penicillium expansum*. The introduction of passive, buoyancy-driven winter ventilation solves the accumulation of toxic metabolic gases like ethylene—which otherwise induces bitter isocoumarin synthesis in carrots—but inherently dehumidifies the macro-environment of the cellar as cold outdoor air warms and expands.

The elegant resolution to this thermodynamic conflict is the implementation of decentralized microclimate zoning using damp quartz sand matrices. By leveraging the matric potential, water retention hysteresis, and high oxygen diffusion rates of a coarse sand medium, root crops are sequestered in an optimal, highly localized 98% RH environment. The sand sacrifices its moisture to the atmospheric vapor pressure deficit, protecting the root tissue from desiccation while allowing the macro-space of the cellar to remain at the 65% RH necessary for the preservation of suberized alliums. When combined with arched ceiling geometries to manipulate dew point condensation and compliant organic sprout suppression regimes, these interdisciplinary methodologies ensure the total postharvest stability and safety of the food supply.

Works cited

1. Postharvest Respiration in Produce | PDF | Vegetables | Berry - Scribd, <https://www.scribd.com/document/48950079/Chilling-and-Respiration-of-fruits-and-vegetables>
2. Storage - Manual for the preparation and sale of fruits and vegetables, <https://www.fao.org/4/y4893e/y4893e06.htm>
3. 64 Root Cellar Construction Principles.pdf
4. R PLEA2017 Proceedings Volume III PDF | Building Insulation - Scribd,

- <https://www.scribd.com/document/465978946/R-PLA2017-proceedings-volume-III-pdf>
5. Development of a Simplified Regression Equation for Predicting, <https://pdfs.semanticscholar.org/3866/9daf91c0c7480a1944a5e833b359e4ba2e9e.pdf>
 6. Estimation of soil moisture and its effect on soil thermal, <https://www.ias.ac.in/public/Volumes/jess/123/08/1793-1807.pdf>
 7. Characteristics and Prediction of the Thermal Diffusivity of Sandy Soil, <https://www.mdpi.com/1996-1073/15/4/1524>
 8. The Commercial Storage of Fruits, Vegetables, and - Bio-Nica.info, <http://www.bio-nica.info/biblioteca/LurieTemperature%20Preconditioning.pdf>
 9. Fresh-cut Fruits. In: The Commercial Storage of Fruits, Vegetables, https://www.academia.edu/32669425/Fresh_cut_Fruits_In_The_Commercial_Storage_of_Fruits_Vegetables_and_Florist_and_Nursery_Stocks_USDA_ARS_Agriculture_Handbook_66
 10. The Commercial Storage of Fruits, Vegetables, and Florist and, <https://www.ars.usda.gov/is/np/CommercialStorage/CommercialStorage.pdf>
 11. Range of single layer waterproofing membranes for roofs, façades, https://www.roofgiant.com/resources/files/rt-brochure_resitrix-range.pdf
 12. environmental product declaration - KÖSTER, https://roofing.koester.eu/files/de_en/K%C3%96STER%20TPO%20Pro_en.pdf
 13. RESITRIX® CL W - CCM Europe, <https://www.ccm-europe.com/gb/epdm/resitrix/waterproofing-membranes/resitrix-cl-w>
 14. Usda Handbook 661954 | PDF - Scribd, <https://www.scribd.com/document/780289964/Usda-Handbook-661954>
 15. Opportunities for Prediction Models to Reduce Food Loss and Waste, <https://www.mdpi.com/2071-1050/16/17/7803>
 16. Post-Harvest Handling, <https://nsnewfarmer.ca/wp-content/uploads/sites/5/2018/03/Post-Harvest-Handling-food-safety.pdf>
 17. Postharvest Storage, Packaging and Handling of Specialty Crops, https://www.researchgate.net/publication/293937411_Postharvest_Storage_Packaging_and_Handling_of_Specialty_Crops_A_Guide_for_Florida_Small_Farm_Producers
 18. Engineering Winter Storage Facilities for Vegetable Crops, https://www.umass.edu/agriculture-food-environment/sites/ag.umass.edu/files/pdf-doc-ppt/engineeringwinterstorage_s_belyea_nevfc_2011.pdf
 19. Postharvest Handling and Storage, <https://nevegetable.org/cultural-practices/postharvest-handling-and-storage>
 20. Keeping Crops Fresh for Market - University of Wisconsin Extension, <https://fyi.extension.wisc.edu/cropstorage/files/2016/01/Keeping-Crops-Fresh-for-Market-2015-handout.pdf>
 21. Bulletin #4135, Storage Conditions: Fruits and Vegetables, <https://extension.umaine.edu/publications/4135e/>

22. Sprout suppression on potato: need to look beyond CIPC for ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4711416/>
23. How to Store Beets Long-Term (With or Without a Root Cellar),
<https://thegroundedhomestead.com/post/preserve-beets-after-harvest>
24. A simple method for determining boundary layer resistance in leaf,
https://www.researchgate.net/publication/229673716_A_simple_method_for_determining_boundary_layer_resistance_in_leaf_cuvettes
25. The Effect of Vapor Pressure Deficit Regulation on the Growth of,
<https://www.mdpi.com/2076-3417/12/7/3667>
26. Transpiration response to soil drying versus increasing vapor,
<https://academic.oup.com/jxb/article/74/16/4789/7206755>
27. Transpiration response to vapour pressure deficit ... - ConnectSci,
<https://connectsci.au/fp/article/53/5/FP26008/272339/Transpiration-response-to-vapour-pressure-deficit>
28. Commercial Production and Management of Carrots,
<https://fieldreport.caes.uga.edu/publications/B1175/commercial-production-and-management-of-carrots/>
29. Should Vegetables Be Stored in High or Low Humidity? - Smart Fog,
<https://www.smartfog.com/insights/should-vegetables-be-stored-in-high-or-low-humidity-produce-storage-guide/>
30. Onion, Scallion, and Shallot, <https://nevegetable.org/book/export/html/265>
31. Fast curing: A method to improve postharvest quality of onions in hot,
https://www.researchgate.net/publication/259135590_Fast_curing_A_method_to_improve_postharvest_quality_of_onions_in_hot_climate_harvest
32. The Organic Seed Grower - Amazon S3,
<https://s3.amazonaws.com/ogwlresources/Files/Organic%20Seed%20Grower,%20The%20-%20John%20Navazio.pdf>
33. Vegetable Diseases PDF | PDF | Edible Plants | Crops - Scribd,
<https://www.scribd.com/document/458834401/Vegetable-Diseases-pdf>
34. Full text of "Onions and other vegetable alliums [electronic resource]",
https://archive.org/stream/Onions_and_Other_Vegetable_Alliums_Crop_Production_Science_in_Horticulture_by_Ja/Onions_and_Other_Vegetable_Alliums_Crop_Production_Science_in_Horticulture_by_James_L._Brewster_djvu.txt
35. Post-Harvest Technology in Horticulture | PDF | Vegetables - Scribd,
<https://www.scribd.com/document/612321922/Post-Harvest>
36. When to Harvest Onions: The Tops-Down Signal, Curing, and,
<https://www.bloomingexpert.com/tips/vegetables/when-to-harvest-onions/>
37. Production Practices and Quality Assessment of Food Crops,
https://archive.org/download/springer_10.1007-1-4020-2535-1/10.1007-1-4020-2535-1.pdf
38. The Plant Pathology Journal,
<https://www.ppjonline.org/upload/pdf/PPJ017-03-04.pdf>
39. Botrytis Neck and Bulb Rot / Onion and Garlic / Agriculture - UC IPM,
<https://ipm.ucanr.edu/agriculture/onion-and-garlic/botrytis-neck-and-bulb-rot/>
40. Mollier diagram | www.swegon.com,

<https://www.swegon.com/references-and-insights/technical-guides/mollier-diagram/>

41. Mollier Chart Air Calculator - AFIM Dehumidifying,
<https://afim-dehumidifier.com/en/mollier-chart-air-calculator/>
42. The Mollier diagram and the Psychrometric Chart,
<http://www.conservaionphysics.org/mollier/mol1.php>
43. Best way to raise humidity in house during cold months? - Reddit,
https://www.reddit.com/r/homeowners/comments/2kc5gv/best_way_to_raise_humidity_in_house_during_cold/
44. FOR MOBILE DEHUMIDIFIERS - HWS-Trade,
<https://www.hws-trade.de/wp-content/uploads/2020/04/SelectionGuide-Dantherm-CDT-Bautrockner-ENG.pdf>
45. Humidity considerations when increasing ventilation - Condair UK,
<https://www.condair.co.uk/en/support-and-resources/knowledge-hub/articles/consider-your-humidity-when-increasing-your-ventilation>
46. Engineering Blog — CalcEngineer, <https://calcengineer.com/blog/>
47. Module for Certification of Energy Manager,
<https://objectstorage.ap-dcc-gazipur-1.oraclecloud15.com/n/axvjbngprylg/b/V2Ministry/o/office-sreda/2024/12/8abc1d252d8146ccbae198f57a9465f9.pdf>
48. Download Publication - California Water Efficiency Partnership,
https://calwep.org/wp-content/uploads/2024/01/CII_Volume-II_July_2014.pdf
49. Root Cellars: A Traditional Holistic Homesteading Practice: Module 3,
<https://gracefulhomesteading.com/pages/holistic-homesteading-module-3-lesson-5-root-cellars>
50. Root Cellar: Definition and Growing Context | Garden ROI,
<https://gardenroi.com/glossary/root-cellar/>
51. How to Store Root Vegetables | She Grows Veg,
<https://shegrowsveg.com/hub/how-to-store-root-vegetables/>
52. Storing Root Vegetables in Sand - GrowVeg.com,
<https://www.growveg.com/guides/storing-root-vegetables-in-sand/>
53. Measuring Soil Water Potential for Water Management in Agriculture,
<https://www.mdpi.com/2071-1050/2/5/1226>
54. Measuring Soil Water Potential for Water Management in Agriculture,
https://www.researchgate.net/publication/45267869_Measuring_Soil_Water_Potential_for_Water_Management_in_Agriculture_A_Review
55. A Guide to Soil Moisture - ConnectedCrops,
<https://connectedcrops.ca/the-ultimate-guide-to-soil-moisture/>
56. Delineation of Soil Moisture Potentials and Moisture Balance,
<https://www.intechopen.com/chapters/72283>
57. Improving water retention capacity of an aeolian sandy soil ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6788994/>
58. Soil–Water Retention Curves Derived as a Function of Soil Dry Density,
<https://www.mdpi.com/2624-795X/1/1/2>
59. Water retention curve - Wikipedia,
https://en.wikipedia.org/wiki/Water_retention_curve

60. Size of Sand Grains Controls Pore Structure and Water Dynamics,
<https://www.mdpi.com/2073-445X/15/5/864>
61. (PDF) Soil Water Characteristic Curves (SWCCs) of Mining Sand,
https://www.researchgate.net/publication/310235912_Soil_Water_Characteristic_Curves_SWCCs_of_Mining_Sand
62. Choice of Pedotransfer Functions Matters when Simulating Soil,
https://e-docs.geo-leo.de/bitstream/11858/8651/1/JAME_JAME21325.pdf
63. Main Drying and Wetting Curves of Soils - Biblioteka Nauki,
<https://bibliotekanauki.pl/articles/64139981.pdf>
64. Hysteresis in Soil Moisture Curves | PDF - Scribd,
<https://www.scribd.com/document/453772783/Water-Retention-Curve-pdf>
65. 3.3 Soil Water Retention – Rain or Shine - OPEN OKSTATE,
<https://open.library.okstate.edu/rainorshine/chapter/3-3-soil-water-retention/>
66. Try or Die: Dynamics of Plant Respiration and How to Survive Low,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8780655/>
67. 10. Measurement of Oxygen Diffusion Rate - ResearchGate,
https://www.researchgate.net/publication/302444715_10_Measurement_of_Oxygen_Diffusion_Rate
68. R2101 – Aerobic Respiration and Crop Storage,
<https://www.edinburghgardenschool.com/2017/05/r2101-aerobic-respiration-and-crop-storage/>
69. Ethylene and 1-MCP regulate major volatile biosynthetic pathways,
https://www.researchgate.net/publication/282487091_Ethylene_and_1-MCP_regulate_major_volatile_biosynthetic_pathways_in_apple_fruit
70. Pectobacterium carotovorum (bacterial soft rot) | CABI Compendium,
<https://www.cabidigitallibrary.org/doi/full/10.1079/cabicompendium.21913>
71. HDC Project Self Assessment And Report Form,
https://projectbluearchive.blob.core.windows.net/media/Default/Research%20Papers/Horticulture/PC%20291_Report_Final%202011.pdf
72. Extracellular Enzymes and Metabolites Involved in Pathogenesis of,
https://www.researchgate.net/publication/226892209_Extracellular_Enzymes_and_Metabolites_Involved_in_Pathogenesis_of_Botrytis
73. Development and Evaluation of a Smart Soil Moisture-Based ... - MDPI,
<https://www.mdpi.com/2624-7402/8/5/193>
74. Quality of Root Vegetables during Prolonged Storage,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20173059932>
75. Quartz Sand and Quartz Gravel | EVERS GmbH & Co. KG,
<https://www.evers.de/en/products/everzitr-filter-materials/filtration/quartz-sand-and-quartz-gravel>
76. a comparison of oxygen diffusion rates in sand moistened,
<https://www.international-agrophysics.org/pdf-107056-37865?filename=A-comparison-of-oxygen-di.pdf>
77. March 2025 - Sustainable Market Farming,
<https://www.sustainablemarketfarming.com/2025/03/>
78. Ethylene Management in Postharvest Handling: A Practical Guide,

- <https://www.biopac.com.au/ethylene-management-postharvest/>
79. Respiration rates (A and B) of intact carrots and isocoumarin levels,
https://www.researchgate.net/figure/Respiration-rates-A-and-B-of-intact-carrots-and-isocoumarin-levels-C-and-D-in-the_fig1_279893817
 80. Factors Influencing Ethylene-induced Isocoumarin Formation and,
<https://journals.ashs.org/downloadpdf/view/journals/jashs/121/3/article-p537.pdf>
 81. Isocoumarin content of carrots as influenced by ethylene,
https://www.researchgate.net/publication/284315882_ISOCOUMARIN_CONTENT_OF_CARROTS_AS_INFLUENCED_BY_ETHYLENE_CONCENTRATION_STORAGE_TEMPERATURE_AND_STRESS_CONDITIONS
 82. Effect of 1-methylcyclopropene, potassium permanganate and silica,
https://horticultureresearch.net/jah/Effect%20of%201-methylcyclopropene,%20potassium%20permanganate_1.pdf
 83. Pre- and Postharvest Factors Control the Disease Incidence of,
<https://www.mdpi.com/2073-4395/10/4/464>
 84. Strategies to Preserve Postharvest Quality of Horticultural Crops and,
<https://www.mdpi.com/2076-3921/9/4/356>
 85. Metabolomic Change Precedes Apple Superficial Scald Symptoms,
<https://pubs.acs.org/doi/10.1021/jf901571g>
 86. EPPO Standards, <https://gd.eppo.int/download/standard/610/pp2-004-2-en.pdf>
 87. Peeling the Onion: Towards a Better Understanding of Botrytis,
<https://apsjournals.apsnet.org/doi/10.1094/PHYTO-06-20-0258-IA>
 88. Botrytis aclada (grey mould of onion) | CABI Compendium,
<https://www.cabidigitallibrary.org/doi/full/10.1079/cabicompendium.9608>
 89. Neck rot (Botrytis allii) of bulb onions - ResearchGate,
https://www.researchgate.net/publication/230056298_Neck_rot_Botrytis_allii_of_bulb_onions
 90. Secondary spread of Sclerotinia rot among carrot roots in storage.,
https://www.researchgate.net/figure/Secondary-spread-of-Sclerotinia-rot-among-carrot-roots-in-storage_fig7_249304208
 91. effects-of-fungal-biodeterioration-on-food-values-qualities-of-carrot,
<https://www.scivisionpub.com/pdfs/effects-of-fungal-biodeterioration-on-food-values-qualities-of-carrot-and-fungicidal-effects-of-plant-oils-of-cymbopogon-citratus--4045.pdf>
 92. Challenges and Opportunities in the Sustainable Improvement of,
<https://www.mdpi.com/2223-7747/13/15/2092>
 93. Storing potatoes and root vegetables: cold, dark, kept apart,
<https://gartenkern.de/en/blog/kartoffeln-wurzelgemuese-einlagern>
 94. Sclerotinia Rot of Carrot - APS Journals,
<https://apsjournals.apsnet.org/doi/pdf/10.1094/PDIS.2003.87.5.456>
 95. Final Report for ONE13-176 - SARE Grant Management System,
<https://projects.sare.org/project-reports/one13-176/>
 96. (PDF) Salinity and nitrogen source affect productivity and nutritional,
https://www.researchgate.net/publication/373140262_Salinity_and_nitrogen_source_affect_productivity_and_nutritional_value_of_edible_halophytes

97. Ecophysiology of *Penicillium expansum* and patulin production in,
<https://bibliotecadigital.ipb.pt/bitstreams/a74705ba-de95-45e1-8009-ef1fde7a36d2/download>
98. Mycotoxins in Food: Cancer Risks and Strategies for Control - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11545588/>
99. Patulin accumulation in apples during postharvest - ResearchGate,
https://www.researchgate.net/publication/248511756_Patulin_accumulation_in_apples_during_postharvest_Effect_of_controlled_atmosphere_storage_and_fungicide_treatments
100. Damage–Safety Trade-Offs in the Transition of Apple Harvesting,
<https://www.mdpi.com/2304-8158/15/16/2787>
101. Sprout Suppressants in Potato Storage: Conventional Options and,
<https://www.mdpi.com/2071-1050/14/11/6382>
102. effect of storage temperature and postharvest tuber treatment with,
<https://bibliotekanauki.pl/articles/55789028.pdf>
103. Achieving sprout suppression in stored potatoes,
<https://ir.library.oregonstate.edu/downloads/08612w507?locale=es>
104. Sprout Inhibition in Potatoes - PotatoPro,
<https://www.potatopro.com/about/sprout-inhibitors>