

Biological Mechanisms of Postharvest Ethylene Production and Structural Mitigation Strategies in Subterranean Root Cellars

The co-storage of postharvest agricultural commodities presents a highly complex physiological and thermodynamic challenge. Climacteric fruits, such as apples and tomatoes,

synthesize and emit significant volumes of ethylene gas (C_2H_4) as a natural byproduct of their ripening and senescence phases¹. When enclosed in unventilated or poorly managed storage environments, this volatile phytohormone accumulates to phytotoxic levels, subsequently acting upon non-climacteric stored commodities such as root vegetables. In root crops like carrots and potatoes, exogenous ethylene triggers profound metabolic shifts, inducing bitter isocoumarin accumulation, breaking or altering dormancy, and accelerating cold-induced sweetening⁴.

To engineer an effective subterranean root cellar—particularly in the specific geotechnical and climatic context of Berlin, Germany—it is necessary to synthesize an understanding of plant biochemistry with advanced structural and thermodynamic engineering. The facility must

maintain a precise microclimate of $0 - 4^{\circ}C$ with $80 - 95\%$ relative humidity, while simultaneously purging metabolic gases⁷. This exhaustive analysis details the molecular biology of ethylene synthesis and signal transduction, the biochemical spoilage mechanisms induced in root vegetables, and the requisite structural, aerodynamic, and geotechnical countermeasures dictated by European standards (such as DIN 18533 and DIN EN 1997-1) to ensure the long-term preservation of mixed agricultural harvests.

The Molecular Biology of Ethylene Biosynthesis in Climacteric Fruits

Ethylene is a simple gaseous unsaturated hydrocarbon that serves as an essential phytohormone, regulating a myriad of plant developmental processes, from seed germination and leaf abscission to the response to abiotic stress⁹. In climacteric fruits, the biological generation of ethylene is characterized by two distinct regulatory systems¹. System 1 represents the basal, autoinhibitory state present in preclimacteric fruit and vegetative tissues, wherein exogenous ethylene actively suppresses the transcription of endogenous biosynthetic enzymes¹.

As the fruit reaches physiological maturity, a developmental transition shifts the tissue into System 2, initiating an autocatalytic surge in ethylene production¹. During this climacteric phase, transcription factors such as RIN-MADS (a MADS-box transcription factor that targets promoter regions) and HB-1 interact directly with the promoters of biosynthetic genes,

triggering an explosion in enzymatic synthesis¹³. Once System 2 is engaged, any localized accumulation of ethylene within a confined storage environment acts as a positive feedback loop, radically accelerating the ripening and senescence of the surrounding biological matter¹.

The Methionine Pathway and the Yang Salvage Cycle

The biochemical synthesis of ethylene is a highly regulated, ATP-dependent pathway that initiates with the sulfur-containing amino acid methionine¹⁰. Methionine is first converted into S-adenosyl-L-methionine (SAM) via the enzyme SAM synthetase, a reaction that consumes one molecule of ATP¹⁰. Because the intracellular pool of free methionine is highly limited and the rate of ethylene emission during the climacteric peak is immense, the plant cannot rely solely on the de novo synthesis of this amino acid.

To sustain autocatalytic ethylene production, climacteric fruits utilize a vital recycling mechanism known as the Methionine Salvage Pathway, or the Yang Cycle¹⁴. During the subsequent enzymatic conversion of SAM, a byproduct known as 5-methylthioadenosine (MTA) is generated⁹. Through a series of phloem-specific enzymatic reactions—often involving 5-methylthioribose kinase (MTK1)—MTA is efficiently recycled back into methionine¹⁶. This continuous recycling ensures that the essential sulfur group is conserved, allowing the fruit tissue to maintain extreme rates of ethylene production without exhausting its primary amino acid reserves¹⁶.

ACC Synthase (ACS): The Rate-Limiting Step

The first committed and historically considered rate-limiting step in ethylene biosynthesis is the conversion of SAM to 1-aminocyclopropane-1-carboxylic acid (ACC), catalyzed by the enzyme 1-aminocyclopropane-1-carboxylate synthase (ACS; EC 4.4.1.14)⁹. ACS is a cytosolic enzyme

belonging to the α, γ -elimination class of pyridoxal-5'-phosphate (PLP)-dependent lyases¹⁷. PLP, the active form of Vitamin B6, acts as a critical electron sink during the reaction, stabilizing the conjugate base and countering the excess electron density generated during deprotonation¹⁷.

The catalytic mechanism of ACS is initiated when the aldehyde group of the PLP coenzyme

forms a Schiff base (an internal aldimine linkage) with the ϵ -amino group of a highly conserved lysine residue, specifically Lys278, within the enzyme's active site¹⁷. Upon the entry of the substrate SAM, an imine exchange occurs, transferring the Schiff base from Lys278 to the amino group of SAM, forming an external aldimine²⁴. A subsequent deprotonation generates a quinonoid intermediate¹⁷.

Crucial to this cyclization reaction is a highly conserved tyrosine residue (Tyr152). Molecular models, crystallography, and site-directed mutagenesis studies on tomato ACS (LeACS2) demonstrate that the hydroxyl group of Tyr152 plays an essential role in stabilizing the ring formation¹⁷. Tyrosine facilitates the nucleophilic attack that cleaves the sulfur-bound carbon, releasing MTA and closing the three-membered cyclopropane ring of ACC¹⁷.

Post-translational regulation of ACS is highly sophisticated, determining the lifespan and activity of the enzyme. ACS enzymes are categorized into three distinct types based on their

C-terminal phosphorylation sites. Type 1 ACS contains both Calcium-Dependent Protein Kinase (CDPK) and Mitogen-Activated Protein Kinase (MAPK) target sites; phosphorylation by these kinases significantly boosts enzyme stability and ethylene production¹⁰. Type 2 ACS contains only CDPK sites and is heavily regulated by ETO1-containing E3 ubiquitin ligases, which target the enzyme for rapid proteasomal degradation via the TOE domain unless phosphorylation prevents ubiquitination¹⁰. Type 3 ACS lacks these C-terminal regulatory sites entirely¹⁰.

Furthermore, ACS exhibits dual enzymatic behavior, possessing $C_\beta - S$ lyase activity in addition to its primary function, allowing it to cleave cystine into thiocysteine, ammonia, and pyruvate²⁷.

ACC Oxidase (ACO): The Final Oxidation

The terminal step of ethylene biosynthesis is catalyzed by 1-aminocyclopropane-1-carboxylate oxidase (ACO; EC 1.4.3.3), which converts ACC into ethylene, carbon dioxide (CO_2), and

hydrogen cyanide (HCN)⁹. The toxic HCN byproduct is rapidly detoxified by β -cyanoalanine synthase to prevent cellular damage, but it also acts to continuously activate the ACO enzyme during reaction turnover²⁸. Accumulating evidence indicates that under specific physiological conditions, ACO, rather than ACS, becomes the rate-limiting factor in ethylene production⁹.

Unlike ACS, ACO is a non-heme iron(II)-dependent oxygenase²⁹. The enzyme requires absolute structural and chemical prerequisites to function: an active site containing a

"2-His-1-carboxylate facial triad" that binds a single ferrous ion (Fe^{2+}), molecular oxygen (O_2) as an electron acceptor, and ascorbate acting as a two-electron donating co-substrate²⁸.

Furthermore, ACO requires bicarbonate (HCO_3^-) or carbon dioxide as a critical activator; empirical evidence points to specific basic residues, such as Arg175, acting as the necessary binding site for this bicarbonate activation²⁸.

The Michaelis-Menten kinetic parameters of ACO suggest that the enzyme exhibits a relatively

low affinity for ACC. In vitro studies on plant ACO reveal an apparent K_m of approximately $156 \pm 8.3 \mu M$ for ACC, with a V_{max} of $5.4 \pm 0.08 \text{ mmol g}^{-1} \text{ h}^{-1}$ on a fresh matter basis³³. The reaction pathway initiates when ACC and ascorbate bind to the active site,

causing the iron center to become five-coordinate²⁹. This allows O_2 to bind and activate, forming a high-valent iron(IV)-oxo species that abstracts an electron from the amine of the ACC substrate, generating a radical cation²⁹. A structural rearrangement ensues, leading to the fragmentation of the cyclopropane ring into ethylene and cyanoformic acid, the latter

spontaneously degrading into CO_2 and cyanide²⁹. Because ACO fundamentally relies on ascorbate and molecular oxygen, placing climacteric fruit in ultra-low oxygen (ULO) environments directly deprives ACO of its requisite electron acceptor, effectively halting the

final biochemical step of ethylene synthesis³⁶.

Biosynthetic Enzyme	Cellular Location	Cofactors / Co-substrates Required	Primary Reaction Mechanism	Kinetic / Regulatory Characteristics
ACC Synthase (ACS)	Cytosol	Pyridoxal-5'-phosphate (PLP), ATP (for SAM)	α, γ -elimination forming a cyclopropane ring	Rate-limiting; regulated by MAPK/CDPK phosphorylation and ETO1 E3-ligase degradation.
ACC Oxidase (ACO)	Cytosol / Apoplast	Fe^{2+} , Ascorbate, O_2 , Bicarbonate (HCO_3^-)	Two-electron oxidation generating a radical cation	Apparent $K_m \approx 156 \mu M$ for ACC. Requires molecular oxygen to complete structural rearrangement.

Ethylene Perception and the Signal Transduction Cascade

The mechanism by which plant tissues perceive and respond to ethylene is a linear, de-repressive signal transduction cascade¹¹. Ethylene perception occurs at the endoplasmic reticulum (ER) membrane via a family of highly specific receptor proteins, the most extensively characterized being ETR1 (Ethylene Response 1), alongside ERS1, ETR2, ERS2, and EIN4¹¹.

The ETR1 Receptor and Copper Coordination

The ETR1 receptor functions as a homodimer and contains a highly conserved ethylene-binding domain (EBD) situated within its three N-terminal transmembrane helices³⁹. The functional

capacity to bind ethylene is entirely dependent on a single copper (Cu^+) cofactor located within the hydrophobic pocket of the receptor dimer⁴⁰. Molecular modeling and site-directed mutagenesis demonstrate that the copper ion is coordinated by a cysteine residue (Cys65) and

a histidine residue (His69) on transmembrane helix II, alongside a critical aspartate residue (Asp25) on helix I⁴⁰. The *etr1-1* mutation, a well-documented genetic variation where Cys65 is substituted with tyrosine, completely abolishes the receptor's ability to chelate copper, rendering the plant dominantly insensitive to ethylene⁴⁰.

The De-Repression Mechanism

In the absence of ethylene, the ETR1 receptor is in an active "on" state³⁷. In this conformation, the receptor actively recruits and stimulates a Raf-like serine/threonine protein kinase called Constitutive Triple Response 1 (CTR1)¹¹. Active CTR1 continually phosphorylates the ER-localized transmembrane protein Ethylene Insensitive 2 (EIN2) at specific residues (Ser645 and Ser924)⁴⁴. This phosphorylation serves as an inhibitory mark, targeting EIN2 for rapid degradation via the 26S proteasome by the F-box proteins ETP1 and ETP2, thereby completely silencing the ethylene response⁴⁴.

When ethylene gas permeates the cellular space and binds to the copper cofactor in the ETR1 receptor, it induces a conformational change that inactivates the receptor³⁷. Consequently, CTR1 is deactivated, and EIN2 is no longer phosphorylated⁴⁴. The stabilization of EIN2 allows an unknown cytoplasmic protease to cleave its cytosolic C-terminal end domain (CEND)⁴⁴. The liberated EIN2-CEND translocates into both the nucleus and specific cytoplasmic foci known as P-bodies⁴⁴. In the P-bodies, EIN2-CEND binds to the 3' untranslated regions (3'UTRs) of the mRNAs encoding the F-box proteins EBF1 and EBF2, effectively repressing their translation⁴⁴. Concurrently, in the nucleus, the absence of EBF1 and EBF2 prevents the continuous proteasomal degradation of the master transcription factors EIN3 and EIL1⁴³. The rapid accumulation and stabilization of EIN3/EIL1 directly activates the transcription of hundreds of downstream ethylene-responsive genes, unleashing the vast array of biochemical changes associated with ripening, senescence, and stress responses⁴⁴.

Ethylene-Induced Phytotoxicity in Stored Root Vegetables

If metabolic gases generated by climacteric fruits are not purged from a mixed agricultural root cellar, exogenous ethylene will readily diffuse into the surrounding non-climacteric root vegetables. Because root vegetables are metabolically active tissues, the perception of ethylene via their own ETR1-homologous receptors initiates a cascade of severe biochemical stress responses that rapidly degrade their organoleptic and commercial qualities⁴.

Isocoumarin Synthesis and Bitterness in Carrots

Carrots (*Daucus carota* L.) are exceptionally susceptible to exogenous ethylene exposure, which triggers the massive accumulation of bitter, phenolic secondary metabolites⁴. The primary compound responsible for this bitter off-flavor is a phytoalexin with a dihydro-isocoumarin skeleton known as 6-methoxymellein (6-MM), structurally defined as

8-hydroxy-6-methoxy-3-methyl-3,4-dihydroisocoumarin ($C_{11}H_{12}O_4$)⁴⁹.

Under standard subterranean storage conditions, 6-MM levels in healthy carrots remain

negligible⁴⁹. However, exogenous ethylene acts as a potent biological elicitor, tricking the carrot tissue into initiating a comprehensive pathogen-defense response⁴. The activated ethylene signal transduction pathway upregulates the phenylpropanoid and polyketide biosynthetic pathways. Specifically, ethylene exposure significantly increases the enzymatic activity of phenylalanine ammonia-lyase (PAL), the gateway enzyme responsible for the deamination of phenylalanine to cinnamic acid, driving phenolic synthesis⁵³.

Downstream in the pathway, ethylene induces the expression of 6-hydroxymellein synthase, a multifunctional polyketide synthase that constructs the foundational isocoumarin ring structure⁵⁴. Subsequently, the enzyme 6-hydroxymellein-O-methyltransferase catalyzes the critical methylation step that converts 6-hydroxymellein to the final 6-MM product⁵⁵. Empirical

studies show that continuous exposure to an ethylene concentration as low as $1 \mu\text{l/L}$ (1 ppm) can reliably induce the formation of 6-MM⁵⁷. The human palate detects a distinct, highly unpleasant bitterness when the concentration of 6-MM exceeds a sensory threshold of

approximately 100 mg/kg of fresh weight⁵⁸. Furthermore, ethylene concurrently stimulates the Embden-Meyerhof-Parnas (EMP) glycolytic pathway in carrots, vastly increasing oxygen uptake and driving the de novo synthesis of other stress-related phenols like isochlorogenic acid and eugenin⁴.

Dormancy Alteration and Cold-Induced Sweetening in Potatoes

Potatoes (*Solanum tuberosum* L.) stored in a multi-use root cellar present a complex, highly concentration- and time-dependent physiological response to ethylene⁵. Short, transient exposures to ethylene rapidly break endodormancy, triggering immediate respiratory spikes and promoting premature sprouting⁵. Conversely, long-term or continuous exposure to ethylene severely inhibits the elongation of those sprouts by suppressing the internal production of auxin and inhibiting auxin signal transduction pathways⁵⁹.

While continuous ethylene might superficially seem advantageous for sprout suppression, its presence during standard cold storage conditions ($0 - 4^{\circ}\text{C}$) exacerbates a catastrophic metabolic disorder known as Cold-Induced Sweetening (CIS)⁶. During cold storage, potatoes naturally undergo starch hydrolysis as a response to thermal stress, accumulating sucrose. Ethylene acts synergistically with cold stress to heavily upregulate the activity of acid invertase⁶³. Invertase rapidly cleaves the non-reducing sucrose into the reducing sugars glucose and fructose⁶³.

When potatoes heavily loaded with glucose and fructose are eventually subjected to high-temperature culinary processing (e.g., frying), these reducing sugars react aggressively with free amino acids, particularly asparagine, via the non-enzymatic Maillard reaction⁶. This yields two highly undesirable outcomes: the development of dark, blackish-brown melanoidin pigments that ruin the visual appearance of the potato, and the synthesis of acrylamide, a potent neurotoxin and probable human carcinogen⁶. To mitigate this, processors are often forced to engage in temperature reconditioning, holding the tubers at elevated temperatures (

14 – 16°C) for weeks to force the reconversion of reducing sugars back into starch, a costly and time-consuming energetic burden⁶.

Root Commodity	Primary Metabolic Disorder	Key Enzymatic / Molecular Drivers	Organoleptic Consequence
Carrot (<i>Daucus carota</i> L.)	Isocoumarin Phytoalexin Synthesis	Phenylalanine ammonia-lyase (PAL), 6-hydroxymellein synthase, O-methyltransferase	Accumulation of 6-methoxymellein (6-MM) causing severe bitterness above 100 mg/kg .
Potato (<i>Solanum tuberosum</i> L.)	Cold-Induced Sweetening (CIS)	Upregulation of Acid Invertase	Cleavage of sucrose into glucose/fructose, leading to Maillard browning and acrylamide formation upon heating.

Thermodynamics and Geotechnical Engineering of Subterranean Storage in Berlin

To defend stored produce from ambient weather extremes while avoiding the financial and energetic burdens of active refrigeration, a root cellar must strictly manipulate geothermal physics. Building a high-performance root cellar in the specific geological context of Berlin presents unique challenges. Berlin is situated within the Warsaw-Berlin *Urstromtal* (a massive glacial meltwater valley), characterized by thick deposits of highly permeable quaternary sands, gravels, and localized layers of *Geschiebemergel* (glacial till/marl)⁶⁷. The engineering must mathematically synthesize soil thermodynamics, extreme dead loads, and rigorous hydrological defenses.

Soil Physics and the Kasuda Equation

The foundational principle of an earth-sheltered cellar is the decoupling of the internal space from high-frequency, high-amplitude diurnal surface temperature swings. Instead, surface temperature fluctuations are forced through the surrounding soil matrix, propagating downward as phase-shifted, highly damped thermal waves⁷.

To precisely calculate the required excavation depth to maintain $32 - 40^{\circ}F$ ($0 - 4^{\circ}C$) through the winter and summer, geotechnical engineers utilize the Kasuda Equation⁷. Derived from the one-dimensional Fourier heat conduction equation, the subsurface temperature

$T_{soil}(z, t)$ is expressed as:

$$T_{soil}(z, t) = T_{mean} - T_{amp} \cdot \exp\left(-z\sqrt{\frac{\pi}{\alpha\tau}}\right) \cdot \cos\left(\frac{2\pi}{\tau}(t - t_{shift}) - z\sqrt{\frac{\pi}{\alpha\tau}}\right)$$

Where T_{mean} is the mean annual surface temperature (approximately $9.5^{\circ}C$ in Berlin),

T_{amp} is the annual surface temperature amplitude, z is the depth in meters, α is the thermal diffusivity of the soil, τ is the period of the temperature cycle (365 days), t is the current day of the year, and t_{shift} is the phase offset corresponding to the minimum surface temperature⁷.

The exponential term in the equation dictates amplitude attenuation, while the cosine term dictates the temporal phase lag. To ensure the extreme cold of mid-winter reaches the cellar

exactly during the peak heat of mid-summer, the thermal diffusivity (α) of Berlin's specific soil profile must be quantified. Thermal diffusivity is defined as the ratio of thermal conductivity (λ) to volumetric heat capacity (C_v) ($\alpha = \frac{\lambda}{\rho_b \cdot c_p}$)⁷.

In Berlin's sandy soils, moisture heavily impacts α . The Johansen method utilizes the dimensionless Kersten number (K_e) to interpolate thermal conductivity between dry and fully saturated states⁷. As the sandy soil wets from rainfall, its thermal conductivity increases exponentially faster than its heat capacity⁷. Consequently, damp sand exhibits maximum thermal diffusivity (often rising from 0.20 to $0.70 \times 10^{-6} m^2/s$), leading to rapid thermal bridging that defeats the phase-lag effect entirely⁷. To mitigate this in a Berlin *Urstromtal* environment, heavy "umbrella insulation" must be deployed. Specifically, cellular glass (e.g., FOAMGLAS or GLAPOR systems)—manufactured by heating crushed recycled glass with carbon to create hermetically sealed, vapor-tight cells—must be extended horizontally below the topsoil to artificially lengthen the thermal path, forcing the heat wave to travel laterally rather than directly down into the roof slab⁷.

Soil Classification	Moisture Condition	Thermal Conductivity λ (W/m·K)	Thermal Diffusivity α ($10^{-6}m^2/s$)	Volumetric Heat Capacity C_v ($106J/m^3\cdot K$)

Clean Sand (Berlin Profile)	Dry	0.30 - 0.58	0.20 - 0.25	1.2 - 1.6
Clean Sand	Saturated	1.70 - 2.40	0.50 - 0.70	3.1 - 3.4
Sandy Loam	Wet (~21%)	0.70 - 1.12	0.35 - 0.45	2.1 - 2.4

Structural Load and Hydrostatic Defenses

The subterranean burial of a concrete structure generates immense, unyielding static forces. A flat cellar roof buried beneath just 0.91 meters (3 feet) of soil must support a continuous vertical dead load exceeding 17.2 kPa (360 psf)⁷. Simultaneously, the vertical walls face massive lateral earth pressures. Because the walls are rigidly propped by the floor slab and roof diaphragm, they cannot yield or deflect to mobilize active pressure, meaning engineers must calculate lateral thrust based on the At-Rest Earth Pressure coefficient (K_0)⁷. Using Jaky's empirical relation ($K_0 \approx 1 - \sin(\phi')$), Berlin's silty sands (with an internal friction angle $\phi' \approx 30^\circ$) yield a K_0 of approximately 0.50⁷. This equates to an Equivalent Fluid Pressure (EFP) of roughly 60 psf/ft , necessitating heavily reinforced concrete walls (typically 200 mm to 300 mm thick depending on depth) to prevent shear failure and maintain restrictive crack widths below 0.2 mm⁷.

However, the most catastrophic failure mode in subterranean architecture involves groundwater. Under the Eurocode 7 (DIN EN 1997-1) standard, structural designs must be strictly verified against the UPL (Uplift) limit state⁷¹. If the local water table in Berlin's low-lying areas rises, the effective unit weight of the soil shifts to its submerged weight, and full hydrostatic pressure acts vertically upward against the floor slab⁷. The upward buoyant force

(calculated as internal basement area $\times$ head of water $\times 9.81 \text{ kN/m}^3$) must be counteracted by the favored dead weight of the concrete structure, to which a strict safety factor of 0.9 must be applied⁷. If buoyancy exceeds this factored gravity, the entire cellar will float, heaving violently out of the ground and fracturing the foundations.

To prevent this, the surrounding sub-grade must include a rigorous perimeter drainage system utilizing perforated pipes wrapped in geotextile filter fabric, adhering strictly to the DIN 4095 standard (*Planning, design and installation of drainage systems protecting structures against water in the ground*)⁷⁴. This drainage intercepts subsurface seepage and routes it to daylight down-slope, ensuring hydrostatic head pressure never accumulates against the walls.

Hydrological Defenses and DIN 18533 Waterproofing

Standards

Even in the absence of bulk hydrostatic pressure, capillary moisture must be absolutely excluded. The exterior envelope of the cellar must be sealed according to the rigorous German standard DIN 18533 (*Waterproofing of components in contact with the ground*)⁵². DIN 18533 categorizes external hydrological threats into distinct Water Exposure Classes (W-E) that dictate precise material specifications⁵².

For a cellar in well-drained Berlin sand equipped with functional DIN 4095 drainage, the vertical exterior walls operate under Class W1.1-E (soil moisture and non-accumulating seepage water in highly permeable soils) or W1.2-E (less permeable soils with drainage)². Under these classes, the concrete must be coated with a highly elastic Polymer-Modified Bitumen Coating (PMBC)⁵². According to DIN 18533-3, the waterproofing applicator must verify that a minimum

dry film thickness (d_{min}) of 3.0 mm is achieved across the walls to bridge structural

micro-cracks (Crack Class R1-E, $\leq 0.2\text{ mm}$ width changes)⁵².

Because PMBC is an emulsion containing water that evaporates during curing, an allowance must be made for shrinkage and substrate unevenness; thus, an application of approximately

4.0 mm to 4.3 mm wet film thickness is strictly required to satisfy the standard

(calculated as $d_z = d_u + d_v$, where d_u accounts for substrate fluctuations and d_v for application variables)⁸¹.

For the earth-covered roof slab (Class W3-E: non-pressing water on earth-covered ceilings), precipitation percolates through the topsoil and rests directly on the structure. Therefore, the PMBC must be reinforced with an embedded glass mesh fabric inlay, requiring a minimum dry

film thickness of 4.0 mm (approximately 5.5 mm wet film thickness)⁵². To protect the bitumen from mechanical damage and biological degradation, it must be overlaid with a rigid dimpled drainage board and a dedicated root-resistant elastomeric membrane (e.g., EPDM rubber) prior to backfilling, guaranteeing that the highly aggressive root systems of surface flora cannot breach the waterproofing envelope⁷.

DIN 18533 Exposure Class	Hydrological Condition	Required Waterproofing Strategy & PMBC Specification
W1.1-E / W1.2-E	Soil moisture and non-accumulating seepage water in permeable soils (or with drainage).	Two-layer PMBC. Minimum Dry Film: 3.0 mm . Minimum Wet Film: $\approx 4.0 - 4.3\text{ mm}$.

W2.1-E	Moderate pressing water (water column $< 3m$).	Two-layer PMBC with glass mesh reinforcement. Minimum Dry Film: 4.0 mm . Wet: $\approx 5.5\text{ mm}$.
W3-E	Non-pressing water on earth-covered ceilings/roof slabs.	Two-layer PMBC with glass mesh reinforcement. Minimum Dry Film: 4.0 mm . Wet: $\approx 5.5\text{ mm}$.
W4-E	Splash water at the wall base and vertical capillary rise.	PMBC bridging the plaster transition. Minimum Dry Film: 3.0 mm . Wet: $\approx 4.0\text{ mm}$.

Psychrometrics, Aerodynamics, and Ethylene Purging Systems

Even with superior thermal and hydrological envelopes, internal environmental parameters must be actively managed. Root vegetables demand a relative humidity (RH) between 80% and 95% to prevent the desiccation, wilting, and shriveling of stored tissues⁷. Operating at this extreme boundary places the atmospheric condition perilously close to the dew point.

Condensation Control and the Magnus-Tetens Equation

The psychrometric relationship between ambient dry-bulb temperature, relative humidity, and the point of condensation is determined by the Magnus-Tetens formulation⁷. To calculate the

dew point (T_d):

$$\gamma = \frac{17.625 \cdot T}{243.04 + T} + \ln\left(\frac{RH}{100}\right)$$

$$T_d = \frac{243.04 \cdot \gamma}{17.625 - \gamma}$$

If the cellar is successfully maintained at 4°C and 90% RH, the Magnus-Tetens equations

yield a dew point of approximately 2.5°C ⁷. If localized thermal bridging through the concrete drops any interior surface temperature to 2.5°C or below—a highly likely scenario during deep winter—atmospheric water vapor will spontaneously condense⁷. In psychrometric enthalpy calculations, the total heat removed from the air incorporates both the sensible heat of the dry air and the latent heat of vaporization of the moisture, making condensation a highly energetic physical process⁸⁵.

Condensation dripping vertically onto root crops promotes devastating bacterial soft rot and rapid fungal proliferation⁷. Therefore, flat ceilings are structurally and psychrometrically unviable. Subterranean architecture universally dictates the use of vaulted or arched ceilings. The steep pitch ensures that condensed droplets do not exceed their surface tension threshold to drip vertically; instead, they adhere to the concrete via capillary action and safely shed laterally down the arch profile, descending the vertical walls and draining harmlessly into a gravel floor⁷.

Aerodynamics: Buoyancy-Driven Stack Ventilation

To safely flush the continuous emission of metabolic ethylene and CO_2 from stored apples and tomatoes, the cellar must rely on passive buoyancy-driven stack effect ventilation. While mechanical ventilation achieves 30-60 Air Changes per Hour (ACH), a passive root cellar must guarantee a minimum of 2 ACH to safely flush ethylene⁷.

The stack effect operates based on the density differential between the internal conditioned air and the external ambient air⁷. The Neutral Pressure Level (NPL) forms a specific horizontal plane within the cellar where internal and external absolute pressures equalize⁷. Below the NPL, negative pressure draws cold, dense ambient air through a lower intake pipe. Above the NPL, positive pressure forces the warmer, lighter internal air containing accumulated ethylene and CO_2 out through an elevated exhaust pipe positioned at the apex of the ceiling vault⁷.

The volumetric airflow rate (Q , in m^3/s) driven by thermal buoyancy is modeled by the standard ASHRAE thermal force equation incorporating Bernoulli's principle for incompressible flow:

$$Q = C_d \cdot A \cdot \sqrt{2g \cdot \Delta H \cdot \frac{T_i - T_o}{T_i}}$$

Where C_d is the discharge coefficient, which drops from approximately 0.60 for unobstructed sharp-edged PVC pipes down to 0.35 if heavy louvered vents are utilized to deter pests⁷. A is the cross-sectional free area of the ventilation pipe, g is the acceleration due to gravity, ΔH is the vertical stack height differential between intake and exhaust, and T_i and T_o are the absolute indoor and outdoor temperatures in Kelvin⁷.

To increase the volumetric removal of ethylene without cutting larger diameter penetrations through the DIN 18533 waterproofing membrane, engineers artificially increase ΔH by extending the exterior exhaust stack higher above the surface grade⁷. Because airflow is directly proportional to the square root of the height differential ($\sqrt{\Delta H}$), this exponentially increases the draft force, ensuring that ethylene concentrations remain below the $1 \mu\text{l/L}$ threshold that triggers 6-methoxymellein synthesis in carrots⁷.

Active Ethylene Mitigation Strategies

In mixed storage scenarios where passive stack ventilation is insufficient to drop the ethylene concentration during the System 2 climacteric peak of large apple harvests, active chemical and biological interventions are mandatory to preserve root crops.

Chemical Scrubbing: Potassium Permanganate (KMnO_4)

One primary postharvest method involves the deployment of potassium permanganate (KMnO_4) impregnated onto highly porous activated alumina beads⁹³. Potassium permanganate is a powerful oxidizing agent that chemically degrades ethylene into carbon dioxide and water⁹⁵. The beads undergo a visible colorimetric change from purple to brown as the compound is exhausted, allowing for visual monitoring⁹⁴.

Trials demonstrate that alumina beads impregnated with $4 \text{ g}/100 \text{ g}$ of KMnO_4 can remove 90% of atmospheric ethylene within 3 hours in a closed system at 20°C ⁹³. However, there is a fundamental thermodynamic limitation when utilizing this technology in a root cellar. Because root cellars must operate at $80 - 95\%$ RH, competitive adsorption occurs⁹³. The highly concentrated water vapor dominates the specific surface area of the alumina carrier, physically blocking ethylene molecules from accessing the KMnO_4 active sites, which significantly diminishes the scrubber's reactivity (efficiency drops to approximately 15% of its dry-air capacity)⁹³.

Receptor Inhibition: 1-Methylcyclopropene (1-MCP)

Consequently, a more direct molecular intervention at the receptor level is widely utilized prior to storage: the application of 1-methylcyclopropene (1-MCP)³. Commercially recognized as a revolutionary advancement in postharvest technology and approved for use within the European Union under Directive 91/414/EEC, 1-MCP is a synthetic cyclic alkene that acts as a potent, irreversible competitive inhibitor of ethylene perception³.

1-MCP possesses a vastly higher affinity for the plant's ETR1 receptors than ethylene itself³. By permeating the tissue as a gas, 1-MCP binds directly to the copper cofactor (Cys65, His69, Asp25) inside the transmembrane EBD of the ETR1 receptor³⁷. This binding locks the receptor in its "on" state³⁷. As established previously, this ensures that the CTR1 kinase remains

continuously active, EIN2 remains phosphorylated and targeted for proteasomal destruction, and the downstream EIN3/EIL1 transcription factors remain completely suppressed³⁷. Because 1-MCP binds irreversibly, the plant tissue remains insensitive to ethylene until the cells transcribe and translate entirely new receptor proteins³⁹. By exposing both the climacteric apples (to halt autocatalytic System 2 production) and the non-climacteric root vegetables (to prevent the perception of any ambient ethylene) to 1-MCP prior to cellar placement, the risk of ethylene-induced phytotoxicity is effectively neutralized at the molecular level, outperforming even advanced Dynamic Controlled Atmosphere (DCA) storage techniques in suppressing aroma volatiles and respiration⁹⁸.

Conclusion

The successful co-storage of climacteric fruits and non-climacteric root vegetables demands a rigorous synthesis of plant molecular biology and subterranean structural engineering. A failure to purge ethylene—synthesized via the methionine cycle and the specific enzymatic actions of PLP-dependent ACS and iron-dependent ACO—inevitably leads to receptor-mediated phytotoxicity in root crops. This manifests biochemically as bitter 6-methoxymellein accumulation in carrots and cold-induced sweetening in potatoes, the latter leading to catastrophic Maillard browning and acrylamide formation upon processing.

To counteract these biological realities, the construction of an earth-sheltered root cellar in the highly permeable sandy soils of Berlin requires exact calculation of soil thermal diffusivity via

the Kasuda equation to guarantee an unbroken $0 - 4^{\circ}C$ thermal baseline, aided by pressure-resistant cellular glass insulation. To protect this delicate microclimate, the structure must be engineered to withstand immense At-Rest Earth Pressures and hydrostatic uplift forces in accordance with Eurocode 7 (DIN EN 1997-1). Furthermore, the implementation of DIN 18533-compliant PMBC waterproofing, combined with a precise aerodynamic stack-ventilation design utilizing vaulted ceilings and calculated discharge coefficients, ensures that humidity remains optimally high without inducing destructive condensation. By coupling these advanced geotechnical defenses with physiological inhibitors like 1-MCP or $KMnO_4$ scrubbers, the long-term, high-quality preservation of mixed agricultural harvests can be mathematically, structurally, and biologically assured.

Works cited

1. Structural representation of the ethylene biosynthesis and the Met or, https://www.researchgate.net/figure/Structural-representation-of-the-ethylene-biosynthesis-and-the-Met-or-Yang-cycle-with_fig1_230850626
2. Ethylene Control of Fruit Ripening: Revisiting the Complex Network, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4677914/>
3. Mechanism unravelling the effect of 1-Methylcyclopropene (1-MCP, https://www.researchgate.net/publication/374142711_Mechanism_unravelling_the_effect_of_1-Methylcyclopropene_1-MCP_through_molecular_docking_approach_and_postharvest_analysis_in_papaya

4. Naturally-Occurring and Ethylene-Induced Phenolic Compounds in, <https://pubmed.ncbi.nlm.nih.gov/30812265/>
5. (PDF) Dual Effects of Ethylene on Potato Dormancy and Sprout Growth, https://www.researchgate.net/publication/7121849_Dual_Effects_of_Ethylene_on_Potato_Dormancy_and_Sprout_Growth
6. Impact of the Temperature Reconditioning of Cold-Stored Potatoes, <https://www.mdpi.com/2304-8158/13/5/652>
7. 64 Root Cellar Construction Principles.pdf
8. Abiotic Stress Hormesis in Carrots: Disease Resistance ... - Corpus UL, <https://corpus.ulaval.ca/bitstreams/78f3fbd7-7699-4e52-9ed3-7c7a1656cb32/download>
9. The Ethylene Biosynthetic Enzymes, 1-Aminocyclopropane-1 ... - MDPI, <https://www.mdpi.com/2218-273X/14/1/90>
10. The Fascinating Role of Ethylene in the Control of Plant, <https://www.preprints.org/manuscript/202309.1166>
11. (PDF) Ethylene Signaling in Regulating Plant Growth, Development, https://www.researchgate.net/publication/388256472_Ethylene_Signaling_in_Regulating_Plant_Growth_Development_and_Stress_Responses
12. Ethylene and Fruit Ripening | Request PDF - ResearchGate, https://www.researchgate.net/publication/229718538_Ethylene_and_Fruit_Ripening
13. Targeted Systems Biology Profiling of Tomato Fruit Reveals ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3490579/>
14. Mutations in Tomato ACC Synthase2 Uncover Its Role in ... - bioRxiv, <https://www.biorxiv.org/content/10.1101/2020.05.12.090431v2.full-text>
15. Preharvest application of aminoethoxyvinylglycine (AVG) modifies, https://www.researchgate.net/publication/221956340_Preharvest_application_of_aminoethoxyvinylglycine_AVG_modifies_harvest_maturity_and_cool_storage_life_of_'Arctic_Snow'_nectarines
16. Yang Cycle Reactions in Bacteria, Yeast, and Plants. - ResearchGate, https://www.researchgate.net/figure/Yang-Cycle-Reactions-in-Bacteria-Yeast-and-Plants_fig1_51096291
17. 1-Aminocyclopropane-1-carboxylate synthase - Wikipedia, https://en.wikipedia.org/wiki/1-Aminocyclopropane-1-carboxylate_synthase
18. Tomato ACS4 is necessary for timely start of and progression, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2014.00466/full>
19. Accumulation and Transport of 1-Aminocyclopropane-1-Carboxylic, <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2017.00038/full>
20. Phloem-Specific Expression of Yang Cycle Genes and Identification, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3123959/>
21. Random mutagenesis of 1-aminocyclopropane-1-carboxylate, <https://www.pnas.org/doi/10.1073/pnas.95.17.9796>
22. Characterization and sequencing of the active site of 1 ... - PubMed,

- <https://pubmed.ncbi.nlm.nih.gov/2122449/>
23. ACC Synthase - Biology,
<https://biology.kenyon.edu/BMB/jmol2014/1IAX/indexACCSynthase.html>
 24. Mechanistic Studies of 1-Aminocyclopropane-1-carboxylate,
<https://pubs.acs.org/bichaw/article/50/11/1950/772325/Mechanistic-Studies-of-1-Aminocyclopropane-1>
 25. Review - Frontiers Publishing Partnerships,
https://www.frontierspartnerships.org/journals/acta-biochimica-polonica/articles/10.18388/abp.2002_3784/pdf
 26. Tyr152 plays a central role in the catalysis of 1-aminocyclopropane,
<https://academic.oup.com/jxb/article/56/418/2203/471052>
 27. Dual activities of ACC synthase: Novel clues regarding the ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8580319/>
 28. (PDF) 1-Aminocyclopropane-1-carboxylic acid oxidase reaction,
https://www.researchgate.net/publication/258639842_1-Aminocyclopropane-1-carboxylic_acid_oxidase_reaction_mechanism_and_putative_post-translational_activities_of_the_ACCO_protein
 29. The nature of O₂ activation by the ethylene-forming enzyme ... - PNAS,
<https://www.pnas.org/doi/pdf/10.1073/pnas.0711626105>
 30. 1-Aminocyclopropane-1-carboxylic acid oxidase determines the fate,
<https://www.biorxiv.org/content/10.1101/2024.02.01.578397v1.full-text>
 31. How Do Electrostatic Perturbations of the Protein Affect the,
<https://pubs.acs.org/doi/10.1021/acs.jpca.1c00141>
 32. ETHYLENE BIOSYNTHESIS - Annual Reviews,
<https://www.annualreviews.org/doi/pdf/10.1146/annurev.pp.44.060193.001435>
 33. Detection of 1-aminocyclopropane-1-carboxylate oxidase activity in,
<https://www.scielo.br/j/txpp/a/9dpvQhGx6Mc5GvcNQfsxGSH/?format=pdf&lang=en>
 34. The role of 1-aminocyclopropane-1-carboxylate (ACC) oxidase, and,
https://www.researchgate.net/publication/229775290_Ethylene_biosynthesis_The_role_of_1-aminocyclopropane-1-carboxylate_ACC_oxidase_and_its_possible_evolutionary_origin
 35. US8247352B2 - Method of inhibiting ethylene production in plants,
<https://patents.google.com/patent/US8247352B2/en>
 36. Regulation by Carbon Dioxide of Wound-induced Ethylene,
https://www.researchgate.net/publication/234022545_Regulation_by_Carbon_Dioxide_of_Wound-induced_Ethylene_Biosynthesis_in_the_Peel_of_Citrus_Fruit
 37. Novel Protein-Protein Inhibitor Based Approach to Control Plant,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2017.01528/full>
 38. Recent advances in ethylene research - Semantic Scholar,
<https://pdfs.semanticscholar.org/c645/b049cd17c05ad32f950e5108e62029810924.pdf>
 39. Ligand-Induced Alterations in the Phosphorylation State of Ethylene,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3440222/>

40. Basis for high-affinity ethylene binding by the ethylene receptor,
<https://www.pnas.org/doi/10.1073/pnas.2215195120>
41. Model of ethylene binding site. Models are shown for the membrane,
https://www.researchgate.net/figure/Model-of-ethylene-binding-site-Models-are-shown-for-the-membrane-topology-of-the-ETR1_fig2_225429002
42. Western-blot analysis indicates mutant ETR1 proteins form,
https://www.researchgate.net/figure/Western-blot-analysis-indicates-mutant-ETR1-proteins-form-disulfide-linked-dimers_fig3_12816835
43. The RING E3 ligase SDR1 destabilizes EBF1/EBF2 and ... - PNAS,
<https://www.pnas.org/doi/10.1073/pnas.2024592118>
44. Endoribonuclease DNE1 Promotes Ethylene Response by ... - MDPI,
<https://www.mdpi.com/1422-0067/25/4/2138>
45. Mechanistic Insights in Ethylene Perception and Signal Transduction,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4577421/>
46. Ethylene-triggered subcellular trafficking of CTR1 enhances the,
<https://www.biorxiv.org/content/10.1101/2020.06.12.148858.full>
47. mRNA decapping proteins regulate EIN2-dependent ethylene,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2026.1897707/full>
48. PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2929093/>
49. Occurrence of 6-Methoxymellein in Fresh and Processed Carrots,
https://www.researchgate.net/publication/8237190_Occurrence_of_6-Methoxymellein_in_Fresh_and_Processed_Carrots_and_Relevant_Effect_of_Storage_and_Processing
50. Structural and Sensory Characterization of Compounds Contributing,
https://www.researchgate.net/publication/10715606_Structural_and_Sensory_Characterization_of_Compounds_Contributing_to_the_Bitter_Off-Taste_of_Carrots_Daucus_carota_L_and_Carrot_Puree
51. 6-Methoxymellein - Wikipedia, <https://en.wikipedia.org/wiki/6-Methoxymellein>
52. Processing Effects on Carrot Phytonutrients - ASHS Journals,
<https://journals.ashs.org/downloadpdf/view/journals/hortsci/41/1/article-p74.pdf>
53. UVA, UVB and UVC Light Enhances the Biosynthesis of Phenolic,
<https://www.mdpi.com/1420-3049/22/4/668>
54. Regulation of biosynthesis of carrot phytoalexin 6-methoxymellein,
<https://agris.fao.org/search/en/providers/122535/records/65dfdf0663b8185d9caf9468>
55. Storage of shredded carrots induces accumulation of chlorogenic,
https://www.researchgate.net/publication/394271769_Storage_of_shredded_carrots_induces_accumulation_of_chlorogenic_acid_and_other_phenolics_without_generating_toxic_or_anti-nutritional_metabolites_A_metabolomics_study
56. PHYTOALEXINS: What Have We Learned After 60 Years?,
https://www.researchgate.net/publication/11652503_PHYTOALEXINS_What_Have_We_Learned_After_60_Years
57. (PDF) Rapid analysis of 6-methoxymellein in carrots by boiling water,
https://www.researchgate.net/publication/222126423_Rapid_analysis_of_6-metho

[xymellein_in_carrots_by_boiling_water_extraction_solid_phase_extraction_and_H PLC](#)

58. sweet and bitter taste in organic carrot - SLU,
<https://pub.epsilon.slu.se/3083/1/IntroLarsKj2007pdf.pdf>
59. Dual effects of ethylene on potato dormancy and sprout growth.,
<https://www.semanticscholar.org/paper/Dual-effects-of-ethylene-on-potato-dormancy-and-Rylski-Rappaport/1f11241539afd1119c4494878faa76e9988bb80a>
60. Involvement of Ethylene in Potato Microtuber Dormancy - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC34794/>
61. Integrative Phytohormone and Transcriptome Analyses Reveal the,
<https://www.mdpi.com/2311-7524/10/3/286>
62. Impact of Agronomic and Storage Practices on Acrylamide in,
https://www.researchgate.net/publication/324712146_Impact_of_Agronomic_and_Storage_Practices_on_Acrylamide_in_Processed_Potatoes
63. Screening potato cultivars for low sugar accumulation during,
https://www.researchgate.net/publication/318232749_Screening_potato_cultivars_for_low_sugar_accumulation_during_storage_at_various_storage_temperatures
64. Heat stress affects carbohydrate metabolism during cold-induced,
https://www.researchgate.net/publication/311246419_Heat_stress_affects_carbohydrate_metabolism_during_cold-induced_sweetening_of_potato_Solanum_tuberosum_L
65. Extensive Variation in Fried Chip Color and Tuber Composition in,
https://www.researchgate.net/publication/41138200_Extensive_Variation_in_Fried_Chip_Color_and_Tuber_Composition_in_Cold-Stored_Tubers_of_Wild_Potato_Solanum_Germplasm
66. An overview of the factors affecting sugar content of potatoes,
https://www.researchgate.net/publication/227769746_An_overview_of_the_factors_affecting_sugar_content_of_potatoes
67. On the age of the young morainic morphology in the area ascribed,
https://www.researchgate.net/publication/248478873_On_the_age_of_the_young_morainic_morphology_in_the_area_ascribed_to_the_maximum_extent_of_the_Weichselian_glaciation_in_north-eastern_Germany
68. R PLEA2017 Proceedings Volume III PDF | Building Insulation - Scribd,
<https://www.scribd.com/document/465978946/R-PLA2017-proceedings-volume-III-pdf>
69. 1 TITLE: On farm and fresh produce management AUTHORS - T-Stor,
<https://t-stor.teagasc.ie/bitstreams/f95bdeac-972e-41f4-bd68-4f99039a5fc5/download>
70. Technical data sheet COMBIDIC -1K - Schomburg,
https://www.schomburg.com/de/en/files/technical-data-sheet_combidic-1k-088f555c-fd7e-4e14-9f8f-9490d936a4c6
71. Modern Geotechnical Design Codes of Practice Implementation,
<https://www.scribd.com/document/724209138/Modern-Geotechnical-Design-Codes-of-Practice-Implementation-Application>
72. Eurocode 7: Pile Design Practices in Europe | PDF | Deep Foundation,

- <https://www.scribd.com/document/508476435/2016-Proceedings-ETC-Volume-II>
73. (PDF) Foundation Systems for High-Rise Structures - Academia.edu,
https://www.academia.edu/95677873/Foundation_Systems_for_High_Rise_Structures
 74. Delta-Technical-Planning-Brochure.pdf,
<https://www.deltamembranes.com/app/uploads/2018/10/Delta-Technical-Planning-Brochure.pdf>
 75. KALZIP® SYSTEMS Products and applications,
<https://www.kalzip.com/wp-content/uploads/2026/02/Kalzip-Products-and-Application-January-2025-USA.pdf>
 76. Practical Sealing System Manual,
https://www.servicesealingsolutions.co.uk/wp-content/uploads/2019/04/Practical_Sealing_System_Manual_EN.pdf
 77. Reduction of Ethylene-induced Physiological Disorders of Carrots,
<https://journals.ashs.org/downloadpdf/view/journals/hortsci/35/7/article-p1312.pdf>
 78. Planning Guide Building Entries - FHRK,
https://fhrk.de/wp-content/plugins/cortexmagick-wp/images/cache/2023/03/Planungshilfe_Uebersetzung_engl_260303.pdf
 79. DIN 18533 - BITUTEC Private Label GmbH 13.4.14,
<https://www.bitutec.de/en/areas-of-application/translate-to-english-din-18533>
 80. Materials, selection, execution – foundation waterproofing (part 1),
<https://nexler.com/en/knowledge-base/materials-selection-execution-foundation-waterproofing-part-i>
 81. DOBAU®-2KS | BIGUMA Baltic,
<https://biguma.lt/en/katalogas/statiniu-apsauga/dobau-2ks>
 82. DOBAU®-2KS 15 L | 2K PMBC Bitumen Thick Coating | DIN 18533,
<https://daemmstoffhandel.de/en/dobau-2ks-2k-pmbc-bitumen-thick-coating-din-18533-15l.html>
 83. Technical Data Sheet W 251 KÖSTER Bikuthan 1C,
https://www.koester.eu/files/tds/files/de_en/TechDataSheet_W%20251_K%C3%96STER%20Bikuthan%201C_de_en_2019-07-16_143132.pdf
 84. dobau®- 2ks,
https://dga.de/_Downloads/technical_data_sheet/en/dobau-2ks_en_tm_dga.pdf
 85. Enthalpy Calculator — kJ/kg, BTU/lb,
<https://calcengineer.com/hvac/enthalpy-calculator/>
 86. Psychrometrics - Energy-Models.com,
<https://energy-models.com/psychrometrics>
 87. Psychrometric Calculations in HVAC Design (With Real Project,
<https://www.nexoradesign.net/post/psychrometric-calculations-in-hvac-design>
 88. Challenges and Opportunities in the Sustainable Improvement of,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11314595/>
 89. Understanding the fundamental role of ethylene in potato storage,
<https://potatoes.ahdb.org.uk/understanding-the-fundamental-role-of-ethylene-in-potato-storage>
 90. Dual Effects of Ethylene on Potato Dormancy and Sprout Growth,

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC541416/>
91. Residential passive ventilation systems : Evaluation and design,
https://www.aivc.org/sites/default/files/members_area/medias/pdf/Technotes/TN54%20RESIDENTIAL%20PASSIVE%20VENTILATION.pdf
 92. WIND-DRIVEN NATURAL VENTILATION IN COURTYARD AND,
<https://etheses.whiterose.ac.uk/id/eprint/3000/1/DX197102.pdf>
 93. Efficacy of Potassium Permanganate Impregnated into Alumina,
https://www.researchgate.net/publication/392352859_Efficacy_of_Potassium_Permanganate_Impregnated_into_Alumina_Beads_to_Reduce_Atmospheric_Ethylene
 94. Activated Alumina PermOx - Sorb-Tech,
<https://sorb-tech.com/activated-alumina-permox>
 95. Controlling Ethylene Levels to Extend Shelf Life of Produce,
<https://agriculture.institute/food-chemistry-and-physiology/controlling-ethylene-levels-extend-shelf-life/>
 96. Postharvest handling of ethylene with oxidative and absorptive means,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10933227/>
 97. The Effects of 1-Methylcyclopropene Treatment on the Fruit Quality,
<https://www.mdpi.com/2077-0472/10/11/490>
 98. DNA Methylation Is Crucial for 1-Methylcyclopropene Delaying,
<https://www.mdpi.com/1422-0067/26/1/168>
 99. Inhibition of Ethylene Binding and Biosynthesis Maintains Fruit,
<https://www.hst-j.org/articles/article/kqZn/>
 100. The Application of 1-MCP in Combination with GABA Reduces,
<https://www.mdpi.com/2077-0472/14/11/2040>
 101. Preharvest Application of 1-Methylcyclopropene Influences Fruit,
<https://journals.ashs.org/view/journals/hortsci/45/4/article-p610.xml>
 102. Impact of Innovative Controlled Atmosphere Storage Technologies,
<https://pubs.acs.org/doi/abs/10.1021/jf802054y>