

Advanced Protocols in Teleost Processing: Mass Transfer Kinetics, Thermodynamic Smoking Modalities, and Defect Mitigation

The global commercial exploitation of teleost fish represents a cornerstone of the modern food industry, demanding rigorous post-harvest interventions to arrest rapid biochemical spoilage while maximizing organoleptic quality and economic yield¹. Fish muscle is inherently highly perishable, characterized by a rapid post-mortem deterioration cascade driven by autolytic enzymatic activity, microbial proliferation, and the chemical oxidation of lipids¹. To mitigate these degradation pathways, the application of sodium chloride (NaCl) combined with thermal dehydration and wood smoke application—collectively known as curing and smoking—has evolved from a rudimentary preservation technique into a highly controlled, multiphase biochemical engineering process³.

The successful execution of hot and cold smoking modalities relies on an intricate equilibrium between osmotic mass transfer, protein solubilization, and thermodynamic control. Central to this equilibrium is the brining protocol. The exact concentration of the saline solution dictates the kinetics of salt diffusion into the sarcoplasm, the solubilization of structural myofibrillar proteins, and the ultimate biosecurity of the final product⁴. However, improper calibration of the brine concentration—specifically exceeding critical thresholds—initiates catastrophic physical defects, most notably the efflorescence and crystallization of salt on the epidermal layer during the drying phase³.

This exhaustive analysis delineates the physical chemistry, thermodynamics, and biosecurity parameters governing the brining and smoking of teleost fish, with a specific focus on the utilization of 70 to 80-degree Salometer brines to achieve uniform flesh penetration without inducing superficial dermal crystallization. Furthermore, it addresses the precise temporal protocols, alternative salting methodologies, and the indispensable formulation of Water Phase Salt (WPS) to neutralize the threat of *Clostridium botulinum*.

Morphological Architecture and Lipid Distribution Dynamics

Before examining the physicochemical mechanisms of brining, it is imperative to establish the structural and compositional parameters of the teleost muscle matrix, as these directly dictate the kinetics of mass transfer. The primary skeletal framework of a round fish consists of a central spinal column from which ribs radiate ventrally to protect the visceral cavity, while neural spines project dorsally¹. The muscle tissue is organized into lateral blocks known as myomeres, separated by collagenous connective tissue called myosepta¹.

The diffusion of aqueous sodium chloride through this highly structured biological matrix is

heavily impeded by the spatial distribution of lipids. Commercially vital species such as Atlantic salmon (*Salmo salar*) and Atlantic mackerel (*Scomber scombrus*) are classified as pelagic fatty fish, possessing lipid contents that can exceed 20% to 30% during peak seasonal feeding migrations². The fat content in fish muscle is highly variable, ranging from mere tenths of a gram in lean species to upwards of 25 grams per 100 grams of product in highly active pelagic species⁸. In salmonids, these lipid deposits are highly concentrated in the ventral belly flap and the dorsal loin regions⁹.

Because lipids are inherently hydrophobic, they act as physical barriers that obstruct the diffusion pathways of aqueous salt solutions². Consequently, highly localized variations in salt concentration arise within a single fillet; lean sections absorb salt rapidly, while fatty adipose tissues demonstrate significantly retarded uptake rates². Thinner pieces and leaner sections reach higher equilibrium salt levels significantly faster than thicker, lipid-dense cuts, which suffer from diffusion limitations¹⁰. This compositional heterogeneity demands a brining protocol capable of overcoming the hydrophobic resistance of lipid-dense regions without simultaneously oversaturating the leaner epaxial musculature.

Pre-Processing Biosecurity and Parasite Inactivation

Prior to the commencement of brining, the raw teleost material must be subjected to rigorous pre-processing controls to ensure baseline biosecurity. The internal muscular tissue of a freshly caught, healthy teleost is generally sterile; the natural microbial load is concentrated in the mucosal slime layer of the epidermis, the respiratory surfaces of the gills, and the dense microbiome of the gastrointestinal tract¹. Processing interventions, including scaling and evisceration, must be executed meticulously to prevent the translocation of these surface and visceral pathogens directly into the sterile muscle tissue¹. Evisceration is legally required for all fish intended for smoking that exceed five inches in total length, ensuring the removal of the viscera and gills, which naturally harbor *Clostridium botulinum* spores¹¹.

For cold-smoked products, which do not undergo a thermal lethality step, the inactivation of zoonotic parasites is an absolute necessity. Teleosts frequently host infective third-stage (L3) larvae of the nematode *Anisakis simplex*¹. To eliminate viable Anisakid larvae and prevent human anisakiasis, stringent freezing protocols have been established by regulatory bodies¹. The core temperature of the fish must be rapidly lowered and maintained at -20°C for no less than 24 hours, or at -35°C for at least 15 hours¹. Laboratory freezing kinetics demonstrate that rapid freezing minimizes the time required for the biological core to reach the critical thermal minimum, thereby preventing the formation of macro-ice crystals that would otherwise rupture the delicate myofibrillar sarcolemma and compromise the structural integrity of the fillet prior to brining¹.

Osmotic Mass Transfer and Fickian Diffusion Kinetics

The incorporation of salt into the fish muscle during wet brining is fundamentally a process of osmotic dehydration and solute impregnation. When a fish fillet is submerged in a hypertonic brine solution, two simultaneous, counter-current fluxes occur: the migration of sodium and chloride ions from the brine into the tissue, and the osmotic exudation of intracellular water

from the tissue into the brine⁵. This mass transfer phenomenon is mathematically modeled utilizing Fick's Second Law of Diffusion, which describes the non-steady-state diffusion of a solute through a permeable medium. For a one-dimensional mass transfer into a slab geometry (approximating a fish fillet), the partial differential equation is expressed as:

$$\frac{\partial C}{\partial t} = D_{eff} \frac{\partial^2 C}{\partial x^2}$$

where C represents the concentration of NaCl in the tissue, t is the brining time, x is the spatial distance from the fillet surface, and D_{eff} is the effective diffusion coefficient¹⁵. The effective diffusion coefficient is an aggregate metric that accounts for the inherent complexities of the biological matrix, including cellular tortuosity, fiber orientation, shrinkage, pH, and localized lipid barriers¹⁶.

Empirical investigations utilizing photon correlation spectroscopy and moisture analysis demonstrate that the D_{eff} for NaCl in teleost muscle typically ranges from 0.47×10^{-10} m²/s to 1.33×10^{-9} m²/s, heavily dependent on the ambient brining temperature and the initial brine concentration¹⁵. Modern processing advancements have explored kinetic enhancers to manipulate this coefficient. For instance, the application of ultrasound during the brining process has been shown to enhance the salt effective diffusion coefficient by 402.8% to 653.21%, creating microscopic cavitation bubbles that accelerate mass transfer through the muscle fibers¹⁹. Conversely, the addition of molecular mediators such as xylitol has been

observed to predictably reduce the D_{eff} of NaCl, allowing for more controlled, slower curing profiles¹⁸.

An increase in the NaCl concentration of the surrounding brine directly steepens the concentration gradient, thereby amplifying the thermodynamic driving force and accelerating the diffusion rate⁵. However, the relationship between brine concentration and mass transfer is non-linear due to the specific biochemical response of the muscle proteins to varying ionic strengths. At lower salt concentrations (approximately 5% to 6% NaCl by weight, equivalent to a physiological ionic strength of 1.0 M), the electrostatic repulsion between protein filaments increases, causing the myofibrillar lattice to swell and uptake water, thereby maximizing the water holding capacity (WHC) of the muscle⁵.

Conversely, when the ambient brine concentration exceeds 9% to 10% NaCl, the high ionic strength induces severe protein dehydration⁵. The proteins form stronger protein-protein bonds, causing the muscle fibers to violently shrink and expel moisture⁵. Therefore, utilizing excessively concentrated brines artificially inhibits deep diffusion, as the superficial muscle fibers rapidly dehydrate, shrink, and form a densified physical barrier—a phenomenon commonly referred to in the industry as case hardening.

Divergent Salting Methodologies: Dry, Wet, and Injection Modalities

The specific methodology utilized to introduce sodium chloride into the teleost matrix profoundly influences the final yield, texture, and degree of fillet gaping. Commercial processors primarily select between dry salting, brine immersion (wet salting), and injection salting, or a highly calibrated mixture of these techniques²⁰.

Dry salting involves the direct application of crystalline NaCl to the surface of the fillet. This establishes a highly aggressive osmotic gradient, rapidly extracting intracellular water to the surface to dissolve the salt and form a spontaneous brine layer¹⁴. Because salt penetration relies on a counter-current flow against the exiting intramuscular water, diffusion into the tissue is markedly slow²⁰. Furthermore, dry salting causes severe myofibrillar fiber shrinkage and extensive water loss, which predictably results in a much lower final product yield¹⁴. However, dry salting generally produces a firmer texture and minimizes the incidence of fillet gaping²⁰. Brine immersion, or wet salting, submerges the fillet entirely in a prepared hypertonic saline solution. In this environment, the muscle cells accumulate the brine and swell²⁰. The chloride ions intercalate between the myofilaments and myosin molecules, freeing up spatial volume that is subsequently occupied by water molecules²⁰. This swelling phenomenon directly explains the superior weight yield achieved by brine-salted fillets compared to dry-salted fillets¹⁴. Nevertheless, the increased hydration and protein solubilization associated with brine salting can lead to a softer texture, paler flesh coloration, and a significantly higher incidence of gaping (the physical separation of muscle flakes along the myosepta)²⁰.

To optimize both throughput speed and yield, industrial processors frequently employ automated brine injection systems. These systems utilize arrays of hypodermic needles to directly deposit precise volumes of hypertonic brine deep into the epaxial and hypaxial musculature, physically bypassing the epidermal diffusion barrier²⁰. Injection salting massively accelerates the curing timeline and delivers the highest overall product yields, sometimes increasing retained weight by up to 5.3% depending on injection pressure and needle speed²¹. However, the mechanical trauma inflicted by the needles, combined with internal hydrostatic pressure, severely exacerbates fillet gaping and structural degradation²¹.

To capture the benefits of multiple techniques, advanced processors utilize mixed salting methods, such as pre-brining followed by dry salting². This hybrid approach involves a preliminary injection or short immersion in saturated brine, immediately followed by the surface application of dry salt². Mixed salting rapidly speeds up the overall permeation of the muscle matrix without inducing the extreme yield losses of pure dry salting or the catastrophic gaping associated with pure injection, maintaining relatively uniform pH, color, and texture profiles across the fillet².

The Salometer Paradigm and Brine Formulation

To exert absolute control over the mass transfer kinetics and prevent the detrimental effects of superficial tissue dehydration during wet brining, the seafood processing industry relies on

precise hydrometric measurements to formulate brines. The specific gravity of the saline solution is quantified using a Salometer (or salinometer), a specialized hydrometer calibrated specifically for sodium chloride solutions at a standard temperature of 60°F (15.5°C)²⁵. The Salometer scale ranges from 0 degrees (representing pure, distilled water) to 100 degrees, which represents a fully saturated brine solution²⁵. At 60°F, a 100-degree Salometer brine contains exactly 26.395% sodium chloride by weight, representing the absolute maximum solubility limit of the salt; beyond this threshold, any additional salt will precipitate out of solution²⁵. Alternative density metrics, such as the Baumé scale, are also utilized in chemical formulations, where the specific gravity of liquids heavier than water is determined by the equation: $\text{Specific Gravity} = 145 / (145 - ^\circ \text{Bé})$ ²⁸. However, the Salometer remains the paramount instrument for fishery applications due to its direct, linear correlation with NaCl saturation percentages.

Salometer Degrees (°SAL)	Baumé Degrees (°Bé)	% NaCl by Weight	Specific Gravity	Lbs of Salt per US Gallon of Water
10	3.7	2.64%	1.018	0.22
30	11.0	7.91%	1.058	0.71
50	18.0	13.19%	1.096	1.26
70	25.0	18.47%	1.139	1.88
80	28.5	21.11%	1.159	2.23
90	32.0	23.75%	1.182	2.59
100	35.0	26.39%	1.204	2.98

For the commercial production of smoked fish, global regulatory and industry standards—including the rigorous methodologies developed by the Torry Research Station—mandate the utilization of brines calibrated strictly between 70 and 80 degrees Salometer (approximately 18.5% to 21.1% NaCl by weight)³. An 80-degree brine represents the optimal mathematical compromise between processing speed and product integrity³. At this concentration, the osmotic gradient is sufficiently steep to drive salt rapidly into the deepest geometric center of the loin within a commercially viable timeframe³. Simultaneously, the 80-degree threshold is purposefully calculated to remain below the catastrophic saturation

limits that cause irreversible superficial dehydration and subsequent aesthetic defects. If processors attempt to accelerate throughput by utilizing ultra-heavy brines exceeding 90 degrees Salometer (approaching the 26.4% absolute saturation limit), the hypertonic shock instantly dehydrates the outermost epaxial myomeres³. While the overall brining time might be marginally reduced, this aggressive strategy guarantees a severe, uneven spatial distribution of salt, leaving the core under-cured while the epidermis becomes highly supersaturated with NaCl³. This supersaturation is the direct, proximate cause of devastating aesthetic defects during the subsequent atmospheric drying stages.

Precise Brining Protocols and Residence Times

The execution of a successful brining operation demands strict temporal management, precisely calibrated against the geometric thickness, lipid content, and ultimate smoking modality of the target species. The residence time in an 80-degree Salometer brine must be perfectly timed; premature extraction results in dangerous under-curing, while excessive immersion leads to organoleptic rejection due to hyper-salinity³.

The protocols differ significantly depending on whether the product is intended for hot or cold smoking. Cold-smoked products generally require longer brine residence times to achieve the deeper salt penetration necessary for preservation, as they will not benefit from a thermal lethality step³. Hot-smoked products utilize the brine primarily for flavor enhancement and surface firming, relying on the subsequent heat of the kiln for microbial destruction, thus allowing for shorter immersion durations³.

Fish Cut / Thickness	Cold Smoking (80° Brine)	Hot Smoking (80° Brine)
Thin Fillets (e.g., Perch, Trout)	5 - 15 minutes	5 - 10 minutes
1/2-inch Fillets	30 minutes	15 minutes
1-inch Fillets	1 hour	30 minutes
1.5-inch Fillets	2 hours	1 hour
Small Whole Fish	2 - 3 hours	1 - 3 hours
Large Whole Fatty Fish (e.g., Herring)	4 - 5 hours	4 hours

During the entire duration of the brining protocol, the thermodynamic state of the solution must be strictly controlled. The ambient temperature of the strong brine should never exceed

60°F (15.5°C)³. Elevated brine temperatures exponentially accelerate the enzymatic degradation of the raw flesh and provide a permissive environment for the rapid logarithmic proliferation of contaminating pathogens³. If the calculated brining residence time exceeds 4 hours, the entire solution must be physically relocated into a refrigerated environment (38°F / 3.3°C), or re-usable ice packs must be introduced to stabilize the temperature³. The direct addition of raw ice to the brine is generally discouraged, as the phase change and melting will dilute the solution, drastically lowering the Salometer reading and altering the calculated mass transfer kinetics³.

Furthermore, when brining fatty pelagic species such as salmon, mackerel, or herring, the fish must remain completely submerged beneath the surface of the brine, often requiring the application of physical weight plates³. If the lipid-dense tissues are permitted to float and breach the surface, exposing the flesh to ambient oxygen, the exposed regions rapidly develop brownish, necrotic-looking lesions—a highly undesirable oxidative condition known colloquially in the industry as "rusting," which severely compromises the aesthetic and chemical stability of the final product³.

Thermodynamics of Deliquescence, Efflorescence, and Epidermal Crystallization

One of the most critical quality control failures in smoked fish production is the manifestation of white, chalky salt deposits across the epidermal layer of the finished product. This phenomenon is not merely an aesthetic blemish; it indicates a fundamental failure in thermodynamic and moisture control. The emergence of these crystals is governed by the atmospheric phase transitions of sodium chloride—specifically, the Deliquescence Relative Humidity (DRH) and the Efflorescence Relative Humidity (ERH)⁶.

Deliquescence is a first-order phase transition wherein a solid crystalline particle spontaneously absorbs atmospheric water vapor to form an aqueous solution droplet³¹. For pure macroscopic sodium chloride, the DRH is an intrinsic thermodynamic property established at exactly 75.3% relative humidity (RH) at ambient temperatures⁶. If the ambient humidity exceeds 75.3%, solid NaCl will liquefy into an aqueous phase. Conversely, efflorescence represents the reverse kinetic pathway. When an aqueous solution of NaCl is subjected to an environment where the relative humidity drops precipitously, water continuously evaporates from the droplet³¹.

However, the droplet does not immediately crystallize upon dropping below the 75.3% DRH threshold. Instead, the solution enters a metastable, highly supersaturated state³².

Crystallization eventually occurs via homogeneous or heterogeneous nucleation only when the ambient humidity falls to the Efflorescence Relative Humidity (ERH), which for bulk NaCl exists dynamically between 43% and 47% RH⁶.

The physics of these phase transitions become vastly more complex when examining the micro-topography of fish skin. Nanoporous biological structures profoundly alter the thermodynamic behavior of salt solutions. Systematic measurements indicate that the DRH of sodium chloride crystals confined within nanopores decreases significantly compared to bulk

crystals³³. For example, the DRH drops from 75.3% down to 58% as the effective pore diameter decreases to 8 nm, a shift primarily driven by the concave curvature of the liquid-vapor interface and the Kelvin effect³². Consequently, salt solutions trapped within the microscopic dermal ridges and scale pockets of a fish exhibit highly unpredictable efflorescence behaviors when subjected to evaporative stress.

When a teleost fillet is immersed in a brine exceeding 90 degrees Salometer, the micro-topography of the epidermal mucosal layer traps a microscopic boundary layer of highly concentrated, near-saturated brine³. As the fish is transferred from the brining vat to the mechanical smoking kiln, it is subjected to the preliminary drying phase. Kiln drying relies on high-velocity airflows and elevated temperatures to rapidly evaporate surface moisture³. As this evaporation occurs, the relative humidity in the immediate microenvironment of the fish skin plummets.

If the initial boundary layer of brine was excessively concentrated (>90° Salometer), the rapid evaporation of the remaining solvent water forcibly drives the solution past its saturation point and deep into the metastable supersaturated zone³. As the kiln air continues to strip away moisture, the localized relative humidity at the skin boundary inevitably drops below the 43% ERH threshold⁶. Upon breaching this thermodynamic limit, the dissolved sodium chloride undergoes spontaneous, rapid nucleation³¹. The salt violently transitions from its aqueous state back into a solid crystalline lattice, precipitating directly onto the pigmented surface of the skin as a highly visible, unattractive white efflorescence (creeping salt structures)³.

To entirely neutralize the threat of efflorescence, two strict procedural parameters must be enforced. First, the primary brine strength must be absolutely restricted to the 70 to 80-degree Salometer window³. This specific concentration ensures that the residual boundary layer liquid contains sufficient solvent water to resist immediate supersaturation during the initial drying phase. Second, following extraction from the 80-degree brine, the fillets must undergo a mandatory, brief post-brining hydrostatic rinse in cold, potable water³. This crucial mechanical intervention flushes away the stagnant, high-molarity boundary layer resting on the skin, replacing it with a highly dilute fluid film. Consequently, when the fish is exposed to the evaporative stresses of the kiln, the surface liquid reaches complete evaporation long before the NaCl concentration can achieve the critical mass required for heterogeneous nucleation, ensuring the final smoked skin remains perfectly glossy and free of solid efflorescence³.

Myofibrillar Solubilization and Pellicle Genesis

While preventing efflorescence on the epidermal side is critical for aesthetic presentation, the primary objective of the preliminary drying phase is the intentional structural modification of the exposed flesh. Exposing the raw, wet musculature directly to hot wood smoke invites the deposition of bitter, uncombusted creosote tars and acidic soot particles, which aggressively degrade the flavor profile³. To facilitate the elegant adsorption of desirable volatile phenols (such as guaiacol and syringol) while protecting the structural integrity of the meat, the processor must engineer the formation of a pellicle³⁰.

The pellicle is a robust, glossy, semi-permeable biopolymer membrane that forms across the cut surface of the fillet³⁰. Its genesis is entirely dependent on the precise biochemical

interaction between the 70 to 80-degree Salometer brine and the underlying endogenous muscle proteins⁴. Teleost muscle is predominantly composed of myofibrillar proteins (MPs)—specifically myosin, actin, actomyosin, paramyosin, and troponin—which collectively account for approximately 50% of the total protein matrix⁴. In their native physiological state, MPs are functionally insoluble in water or low-ionic-strength media, requiring aggressive osmotic intervention for extraction⁴.

When the 80-degree brine diffuses into the epaxial musculature, the high localized ionic strength (significantly exceeding 0.3 M NaCl) acts as a powerful chaotropic agent⁴. The chloride ions aggressively intercalate between the myosin heavy chains, disrupting the electrostatic linkages and hydrogen bonds holding the thick and thin filaments together⁴. This severe electrostatic repulsion causes the myofibrillar lattice to completely depolymerize and solubilize into a viscous, sticky proteinaceous exudate²⁰. This solubilized matrix of extracted myosin and actin is drawn to the external surface of the fillet during the osmotic exchange of the brining process.

Upon entering the preliminary drying phase of the kiln—typically conducted under high-velocity airflow at temperatures between 30°C and 38°C—the rapid evaporation of surface water initiates the complex cross-linking and polymerization of this extracted exudate³. The sticky surface desiccates and solidifies into the highly resilient, tacky biopolymer membrane known as the pellicle³⁰.

This membrane serves two profound thermodynamic and functional purposes. First, it acts as a molecular sponge; the tacky polymerized proteins possess a high affinity for capturing and binding the vapor-phase aromatic compounds generated during wood pyrolysis, ensuring deep flavor infusion³⁰. Second, during the extreme thermal stress of hot smoking (where internal temperatures approach 80°C), the pellicle acts as a rigid, impermeable physical barrier that strictly prevents the catastrophic exudation and loss of valuable intramuscular lipids and moisture³⁸. If the preliminary solubilization provided by the 80-degree brine is skipped, or if the drying phase is rushed (failing to allow the necessary 1 to 12 hours for proper pellicle maturation), the membrane fails to form, resulting in a dry, friable, and severely texturally degraded final product³.

Execution of Smoking Modalities: Hot and Cold Protocols

Following the successful establishment of the pellicle, the teleost must be subjected to the highly controlled thermal and atmospheric conditions of the smoking chamber. Commercial processing is starkly divided into two distinct processing modalities—cold smoking and hot smoking—each defined by rigid thermodynamic boundaries and entirely different biosecurity philosophies¹¹.

Cold Smoking Kinetics

Cold smoking is explicitly engineered to impart deep aromatic complexity and partial dehydration without initiating any thermal coagulation or denaturation of the myofibrillar proteins⁴³. The objective is to preserve the native, translucent, sashimi-like texture of the raw

flesh. To achieve this, the ambient temperature within the smokehouse must be stringently restricted. According to international processing standards and the Torry Advisory protocols, the internal temperature of a cold-smoking kiln must never exceed the critical threshold of 32°C to 33°C (approximately 90°F) at any point during the cycle⁴⁵.

Because cold smoking fundamentally operates in a mesophilic temperature zone that is highly permissive to the rapid proliferation of pathogenic bacteria, the process itself offers absolutely zero thermal lethality¹¹. A standard cold smoking cycle does not achieve a 6D (99.9999%) logarithmic reduction of target organisms¹¹. Consequently, the absolute safety of a cold-smoked product rests entirely on the synergistic hurdle effect established during the pre-processing phases. The processor must rely completely on the aggressive lowering of

water activity (a_w) achieved through osmotic dehydration, the bacteriostatic toxicity of the deposited wood smoke phenols, and most critically, the precise chemical concentration of salt within the aqueous phase of the tissue⁴⁵.

Hot Smoking Execution: The Torry Kiln Standard

Conversely, hot smoking is a destructive thermal process explicitly designed to forcefully denature the proteins, fully cook the flesh, and deliver significant microbial lethality²⁹. The execution of a commercial hot smoke requires highly precise, sequential temperature ramping to ensure the core of the fish achieves pasteurization temperatures without violently rupturing the exterior musculature. The modern standard utilizes mechanical forced-convection systems, such as the Torry Kiln, progressing through three distinct thermodynamic stages²⁹.

1. **Stage 1: Preliminary Desiccation (30°C).** The kiln is initially set to 30°C with maximal smoke output and high volumetric airflow (air inlet half to three-quarters open)²⁹. Sustained for 30 to 60 minutes, this low-temperature phase is strictly dedicated to evaporating superficial moisture and completing the polymerization of the pellicle. Attempting to apply high heat immediately to wet flesh results in the violent boiling of surface water, which irreversibly shreds the delicate myomeres and ruins the physical appearance³.
2. **Stage 2: Coagulation and Aromatic Deposition (50°C).** The thermostat is elevated to 50°C, and the fresh air intakes are heavily restricted (quarter open) to dramatically elevate the relative humidity within the chamber²⁹. Elevated humidity is crucial at this stage to prevent excessive total weight loss (yield reduction)²⁹. Held for 30 to 45 minutes, this phase initiates the primary heat-induced denaturation of the myosin heavy chains, causing the flesh to firm significantly while maximizing the deep tissue absorption of phenolic smoke compounds²⁹.
3. **Stage 3: Thermal Lethality (80°C).** The final stage demands an aggressive temperature spike to 80°C²⁹. At this temperature, the remaining actin and sarcoplasmic proteins fully coagulate, transforming the tissue from a translucent gel into an opaque, rigid matrix²⁹. The ambient 80°C environment must be maintained until the deepest geometric core of the thickest dorsal loin registers a temperature sufficient to guarantee the destruction of vegetative pathogens²⁹.

For highly specialized products, processors frequently deviate from the standard three-stage ramp to achieve unique culinary outcomes. For example, in the production of dogfish flaps (Schillerlocken), the kiln temperature is set directly to a harsh 90°C from the moment the fish enters the chamber²⁹. Operating with the air inlet minimally open to retain extreme heat and maximum smoke density, the flaps are subjected to an intense 60-minute thermal shock, yielding a dark, heavily caramelized, and rapidly cooked exterior²⁹. Similarly, buckling (hot-smoked whole herring) undergoes the three-stage ramp but requires careful intra-kiln rotation and partial cooling within the chamber using diverted ambient air to set the structure before the fish is moved to a chillroom²⁹.

Biosecurity, Water Phase Salt (WPS), and *Clostridium botulinum*

While hot smoking delivers a thermal kill step for vegetative bacteria (such as *Listeria monocytogenes* or *Salmonella* spp.), the absolute paramount toxicological threat in the commercial seafood industry is the germination and subsequent neurotoxin production by the spore-forming obligate anaerobe *Clostridium botulinum*⁷. The marine environment is heavily contaminated with *C. botulinum* Type E, as well as non-proteolytic Types B and F⁵². Unlike their terrestrial proteolytic counterparts, these specific aquatic strains are psychrotrophic; they possess the terrifying biological capability to grow, metabolize, and synthesize lethal botulinum neurotoxin at standard commercial refrigeration temperatures as low as 38°F (3.3°C)⁵².

The threat of botulism is profoundly magnified by modern packaging technologies. To inhibit the growth of aerobic spoilage organisms (such as *Pseudomonas*) and extend the physical shelf life of the smoked product, manufacturers predominantly utilize Reduced Oxygen Packaging (ROP) or Modified Atmosphere Packaging (MAP)¹. By vacuum-sealing the smoked fish and eradicating ambient oxygen, the processor inadvertently creates the perfect, optimal anaerobic microenvironment required for *C. botulinum* spores to germinate²⁹. Because cold smoking does not destroy the highly heat-resistant spores, and even hot smoking parameters often fail to achieve absolute sterilization, the industry must rely on strict biochemical formulation to ensure safety¹¹.

The primary metric enforced by global food safety authorities—specifically the United States Food and Drug Administration (FDA)—to neutralize *C. botulinum* is the Water Phase Salt (WPS) concentration⁵¹. WPS is a mathematically derived parameter that quantifies the actual concentration of sodium chloride residing strictly within the biologically available aqueous portion of the tissue, completely disregarding the inert lipid and solid protein fractions⁵⁴. The mathematical formula is defined as:

$$\text{WPS}(\%) = \left(\frac{\% \text{NaCl}}{\% \text{NaCl} + \% \text{Moisture}} \right) \times 100$$

This distinction is biochemically vital. Microorganisms do not inhabit lipids; they exist exclusively

within the aqueous sarcoplasm⁵⁶. In highly fatty teleosts like salmon or mackerel, the massive volume of intramuscular lipid physically displaces water⁸. Consequently, two fish fillets (one lean cod and one fatty salmon) could both register a total overall salt content of 2.0% by total weight. However, because the salmon possesses significantly less water due to its high fat content, the salt dissolved within that limited remaining water is forced into a much higher concentration. Thus, the salmon will register a significantly higher, more lethal WPS percentage than the cod, exposing contaminating pathogens to much higher osmotic stress⁵⁴.

To legally and safely package smoked fish in anaerobic or reduced-oxygen environments, the FDA and corresponding international health agencies mandate that the brining protocol must achieve a minimum WPS of 3.5% in the deepest, thickest portion of the loin muscle⁵¹. If the formulation achieves this absolute 3.5% threshold, the extreme osmotic pressure permanently arrests the germination of *C. botulinum* Type E spores, even if the product is subjected to moderate temperature abuse during the logistics chain⁵². Failure to hit this 3.5% WPS mark requires the implementation of alternative, highly restrictive hurdles, such as freezing the product permanently or utilizing highly regulated chemical preservatives like sodium nitrite⁵⁷. This 3.5% WPS threshold is also highly effective against other devastating pathogens. Empirical challenge studies demonstrate that a WPS ranging between 2.39% and 4.36% successfully arrests the proliferation of *Listeria monocytogenes* in cold-cured salmon products stored at 3°C for up to 30 days⁵⁸. Therefore, the 70 to 80-degree Salometer brining step is not merely a culinary enhancement; it is a legally mandated, life-saving toxicological intervention.

Biochemical Degradation, Lipid Peroxidation, and Spoilage Pathways

While the attainment of a 3.5% WPS is non-negotiable for biosecurity against anaerobic pathogens, processors must recognize that the application of sodium chloride introduces a severe biochemical paradox. The muscle tissue of teleost fish is exceptionally rich in highly unsaturated, long-chain omega-3 polyunsaturated fatty acids (PUFAs), most notably eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)⁸. These fragile lipid molecules, while vital for human nutrition, are extraordinarily susceptible to oxidative degradation during processing².

When NaCl is infused into the muscle matrix during brining, it acts as a highly aggressive pro-oxidant⁶¹. The presence of sodium chloride accelerates the catalytic oxidation of lipids, generating free radicals that rapidly cleave the delicate double bonds of the PUFAs⁷. This peroxidation cascade yields secondary oxidative byproducts, primarily malondialdehyde (MDA) and volatile aliphatic aldehydes, which manifest organoleptically as severe rancidity and putrid, cardboard-like off-flavors³⁹. The progression of this lipid oxidation is objectively quantified using the Thiobarbituric Acid Reactive Substances (TBARS) index⁴⁸. Fortunately, the deposition of phenolic compounds derived from the pyrolysis of hardwood chips during the smoking phase acts as a powerful counterbalance, providing significant antioxidant effects that partially neutralize the pro-oxidant activity of the salt, thereby stabilizing the lipid fraction for extended chill storage⁵⁰.

Furthermore, the brining and smoking processes are engaged in a perpetual race against the organism's own endogenous autolytic programming¹. Upon harvest and the cessation of aerobic respiration, the rapid post-mortem accumulation of lactic acid drives the intracellular pH downward¹. This acidification ruptures cellular lysosomes, unleashing aggressive proteolytic enzymes known as cathepsins (specifically Cathepsins D and L) alongside calcium-activated calpains directly into the sarcoplasm¹. These enzymes systematically dismantle the structural actin and myosin networks, leading to extreme tissue softening, gaping, and the total collapse of the fillet's structural integrity¹.

Simultaneously, surviving psychrotrophic bacteria (such as *Shewanella* and *Pseudomonas* species) exploit the free amino acids liberated by this proteolysis. These spoilage organisms rapidly reduce Trimethylamine Oxide (TMAO)—a natural marine osmolyte—into Trimethylamine (TMA), the highly volatile compound universally responsible for the pungent, rotting odor of stale seafood¹. The accumulation of TMA, alongside ambient ammonia, is measured collectively as Total Volatile Basic Nitrogen (TVB-N)¹. TVB-N serves as the primary threshold indicator utilized by global food safety authorities to quantify late-stage bacterial spoilage, with smoked fish generally deemed unfit for human consumption when TVB-N values exceed 35 mg N/100 g⁴⁸.

More dangerously, specific enteric bacteria possessing histidine decarboxylase enzymes will attack free histidine, converting it directly into histamine¹. The consumption of histamine-laden fish induces scombroid toxicosis, triggering severe, potentially life-threatening anaphylactic-like cardiovascular events¹. Because histamine synthesis is entirely enzyme-driven and accelerates exponentially when ambient temperatures breach the 4.4°C threshold, strict thermodynamic management must be maintained continually from the harvest vessel, through the brining vats (which must never exceed 15.5°C), until the fish reaches the lethal temperatures of the hot-smoking kiln or the cryogenic temperatures of the freezing hold¹.

Conclusion

The commercial execution of hot and cold smoking for teleost fish demands an absolute, uncompromising mastery of thermodynamic principles, mass transfer kinetics, and biochemical engineering. The processing facility must navigate a precarious biological tightrope. The primary mass transfer mechanism—wet brining—must be executed using solutions calibrated strictly between 70 and 80 degrees Salometer. This precise specific gravity optimizes the driving force of Fickian diffusion, ensuring that the deepest epaxial musculature achieves the legally mandated 3.5% Water Phase Salt required to neutralize the lethal threat of *Clostridium botulinum* Type E.

Crucially, restricting the brine concentration to a maximum of 80 degrees Salometer, combined with a mandatory post-brine hydrostatic rinse, physically prevents the superficial boundary layer from breaching the Efflorescence Relative Humidity (ERH) threshold during kiln dehydration. This effectively neutralizes the threat of heterogeneous nucleation and the disastrous formation of macroscopic salt crystals upon the epidermal tissue. Simultaneously, this optimal ionic strength selectively solubilizes the necessary myofibrillar proteins, extracting myosin and actin to the surface to polymerize into a robust, protective pellicle. By harmonizing

these mass transfer kinetics with the rigid, three-stage thermodynamic controls of the modern smoking kiln, the processing industry can successfully arrest the autolytic and microbial spoilage cascades, delivering a teleost product of unparalleled biosecurity, structural integrity, and organoleptic excellence.

Works cited

1. 60 Fish Preparation and Safety Guide.pdf
2. Effect of dry salt versus brine injection plus dry salt on the ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9637810/>
3. Curing Fish - Meats and Sausages, <https://www.meatsandsausages.com/fish/processing/curing>
4. Solubilization of Myofibrillar Proteins in Water or Low Ionic Strength, https://www.researchgate.net/publication/286375991_Solubilization_of_Myofibrillar_Proteins_in_Water_or_Low_Ionic_Strength_Media_Classical_Techniques_Basic_Principles_and_Novel_Functionalities
5. Effects of brining process on nutrient composition of fish species, <https://scispace.com/pdf/effects-of-brining-process-on-nutrient-composition-of-fish-4pyuwiywu9.pdf>
6. Size Matters in the Water Uptake and Hygroscopic Growth of, <https://pubs.acs.org/jpcafh/article/119/19/4489/1397904/Size-Matters-in-the-Water-Uptake-and-Hygroscopic>
7. Influence of hot-smoking on the stability of fresh and frozen-thawed, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11266935/>
8. Canned Fish in Brine—Variability in Macronutrient and Fatty Acid, <https://www.mdpi.com/2079-7737/15/5/381>
9. Spatial fat distribution images for whole salmon fillets - ResearchGate, https://www.researchgate.net/figure/Spatial-fat-distribution-images-for-whole-salmon-fillets_fig2_283266196
10. Influence of brine concentration of Atlantic Salmon fillet salting, https://www.researchgate.net/publication/223472450_Influence_of_brine_concentration_of_Atlantic_Salmon_fillet_salting
11. Understanding Hot and Cold Smoked Fish Processing and Safety, <https://www.food-safety.com/articles/10016-understanding-hot-and-cold-smoked-fish-processing-and-safety>
12. The Impact of Frozen Storage of Raw Material Prior to Processing on, <https://www.preprints.org/manuscript/202506.2197>
13. Solubilization of fish myofibrillar proteins in NaCl and KCl solutions, https://www.researchgate.net/publication/377994379_Solubilization_of_fish_myofibrillar_proteins_in_NaCl_and_KCl_solutions_A_DIA-based_proteomics_analysis
14. and brine-salted salmon later treated with liquid smoke flavouring, <https://journal.fi/afs/article/download/6020/5217/>
15. Influence of the process parameters on osmotic dehydration ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4711434/>
16. Mathematical model and dependence of fish salting process,

- <https://www.scielo.br/j/bjft/a/9zq69zSyxQZPMWxj4bsTHLt/?format=pdf&lang=en>
17. Salting kinetics and salt diffusivities in farmed Pantanal caiman,
<https://repositorio.unesp.br/server/api/core/bitstreams/8229ffb3-c38c-4298-9c7b-2541ffcb1bab/content>
 18. Simulation Study of Xylitol-Mediated Effect on NaCl Diffusion ... - MDPI,
<https://www.mdpi.com/2304-8158/12/7/1451>
 19. Mass transfer enhancement of tuna brining with different NaCl,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8971329/>
 20. (PDF) Effect of dry salt versus brine injection plus ... - ResearchGate,
https://www.researchgate.net/publication/364724650_Effect_of_dry_salt_versus_brine_injection_plus_dry_salt_on_the_physicochemical_characteristics_of_smoked_salmon_after_filleting
 21. Product Yield and Gaping in Cold-smoked Atlantic Salmon (Salmo,
https://www.researchgate.net/publication/229812635_Product_Yield_and_Gaping_in_Cold-smoked_Atlantic_Salmon_Salmo_salar_Fillets_as_Influenced_by_Different_Injection-salting_Techniques
 22. The quality of cold smoked salmon Influence of raw material and,
https://orbit.dtu.dk/files/5585167/PhD-afhandling_220507.pdf
 23. Effect of salting and cold-smoking procedures on Atlantic salmon,
<https://archimer.ifremer.fr/doc/00423/53475/54382.pdf>
 24. unuftp.is Final Project 2017 The effects of different salting,
https://www.grocentre.is/static/gro/publication/495/document/Moe%20Phyu%20OO_final%20project_2018.pdf
 25. Sodium Chloride Brine Density Tables | PDF - Scribd,
<https://www.scribd.com/document/383468680/Brine-Calculator>
 26. Making brine - Meats and Sausages,
<https://www.meatsandsausages.com/sausage-making/curing/making-brine>
 27. Salometer Guide for Brine Measurement | PDF | Weight - Scribd,
<https://www.scribd.com/document/73567264/Salometer-Usage>
 28. Tech Info Baume Scale - ProMinent Fluid Controls,
<http://prominent.us/webroot/promx/pdf/tech%20info%20baume%20scale.pdf>
 29. TORRY ADVISORY NOTE No. 82 (Revised) Hot smoking of fish,
<https://www.fao.org/4/x5953e/x5953e00.htm>
 30. Little Chief Smoked Salmon Deluxe Recipe - Recipeland,
<https://recipeland.com/recipe/v/little-chief-smoked-salmon-deluxe-39924>
 31. Nucleation Kinetics in Mixed NaNO₃/Glycerol Droplets Investigated,
<https://pubs.acs.org/doi/10.1021/acs.jpcb.5b12442>
 32. Efflorescence relative humidity of airborne sodium chloride particles,
https://www.researchgate.net/publication/222995747_Efflorescence_relative_humidity_of_airborne_sodium_chloride_particles_A_theoretical_investigation
 33. Deliquescence of NaCl Confined in Nanoporous Silica - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/36037488/>
 34. Hygroscopic behavior of NaCl–MgCl₂ mixture particles as nascent,
<https://acp.copernicus.org/articles/15/11273/2015/acp-15-11273-2015.pdf>
 35. (PDF) Measurement of relative humidity dependent light scattering,

- https://www.researchgate.net/publication/40892792_Measurement_of_relative_humidity_dependent_light_scattering_of_aerosols
36. Investigation of NaCl deliquescence in porous substrates using RH, https://www.researchgate.net/publication/225702498_Investigation_of_NaCl_deliquescence_in_porous_substrates_using_RH-XRD
 37. Kinetics of Salt Creeping on a Free Surface: From Nucleation ... - arXiv, <https://arxiv.org/html/2604.05789v2>
 38. Skinned fish soaking in 13% (w/v) brine - ResearchGate, https://www.researchgate.net/figure/Skinned-fish-soaking-in-13-w-v-brine_fig3_313221194
 39. Research Progress on the Mechanism of the Impact of Myofibrillar, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11506927/>
 40. Earth and Space Science - Nova Scotia Curriculum, <https://curriculum.novascotia.ca/sites/default/files/documents/resource-files/science%20%20TRfiles.pdf>
 41. Impact of Extraction Method on the Detection of Quality Biomarkers, <https://www.mdpi.com/2304-8158/10/5/1097>
 42. Protein Extraction From Chicken Myofibrils Irrigated with Various, <http://lib3.dss.go.th/fulltext/Journal/Journal%20of%20food%20science/2000%20v.65/no.1/jfsv65n1p0096-0100ms19990513%5B1%5D.pdf>
 43. TECHNOLOGY OF MEAT, FISH AND POULTRY PRODUCTS | DOCX, <https://www.slideshare.net/slideshow/technology-of-meat-fish-and-poultry-products-149114245/149114245>
 44. Smoked Fish Processing Standards | PDF - Scribd, <https://www.scribd.com/document/707543240/PNS-FDA-27-2010-Recommended-Code-of-Practice-for-the-Processing-and-Handling-of-Smoked-Fish>
 45. (PDF) CHANGES IN MICROBIOLOGICAL QUALITY AND SAFETY, https://www.researchgate.net/publication/351167813_CHANGES_IN_MICROBIOLOGICAL_QUALITY_AND_SAFETY_OF_VACUUM-PACKED_COLD-SMOKED_SALMON_Salmo_salar_DURING_STORAGE
 46. Recommendations for the Preparation of Smoked Salmon, <https://www.fao.org/4/x5886e/x5886e00.htm>
 47. 1 Heat Process Validation Though consumers may not know ... - FDA, <https://www.fda.gov/media/100346/download>
 48. Microbial and Physico-Chemical Characterization of Cold Smoked, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10378578/>
 49. Fish Smoking Procedures Guide | PDF - Scribd, <https://www.scribd.com/document/59316777/Fish-Smoking-Procedures>
 50. The Effect of Storage Temperature on the Chemical and Sensorial, <https://www.trjfas.org/pdf.php?id=408>
 51. United States v. Nova Scotia Food Products Corp., 417 F. Supp., <https://law.justia.com/cases/federal/district-courts/FSupp/417/1364/1455247/>
 52. Fish and Fishery Products Hazards and Controls Guidance - FDA, <https://www.fda.gov/files/food/published/Fish-and-Fishery-Products-Hazards-and-Controls-Guidance-Chapter-13-Download.pdf>

53. Potential Hazards in Cold-Smoked Fish: Clostridium botulinum type E,
https://www.researchgate.net/publication/229827859_Potential_Hazards_in_Cold-Smoked_Fish_Clostridium_botulinum_type_E
54. F.B.14.14 Salting | Fish plant guidelines | ontario.ca,
<https://www.ontario.ca/document/fish-plant-guidelines/fb1414-salting>
55. Cured salted and smoked fish establishments good manufacturing,
https://www.afdo.org/wp-content/uploads/2020/11/Cured_salted_and_smoked_fish_establishments_good_manufacturing_practices_acc_updated_2019.pdf
56. Water-Phase Salt: What It Means in Meat Curing - The Curesmith,
<https://www.thecuresmith.com/codex/water-phase-salt/>
57. Labeling Requirements for Alaska Seafood Processors,
https://alaskaseagrant.org/?sdm_process_download=1&download_id=9828
58. Effect of Water Phase Salt Content and Storage Temperature on,
https://www.researchgate.net/publication/5850632_Effect_of_Water_Phase_Salt_Content_and_Storage_Temperature_on_Listeria_monocytogenes_Survival_in_Chum_Salmon_Oncorhynchus_keta_Roe_and_Caviar_Ikura
59. Effect of brine concentration on the quality of salted large yellow,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11422311/>
60. Effect of Different Cooking Methods on Lipid Content and Fatty Acid,
<https://journal.pan.olsztyn.pl/pdf-159651-87710?filename=87710.pdf>
61. Influence of Storage Time and Method of Smoking on the Content of,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8890880/>
62. EFFECTS OF SMOKING WITH DIFFERENT WOOD CHIPS AND,
<https://dergipark.org.tr/en/pub/gida/article/626414>
63. Solubilization of fish myofibrillar proteins in NaCl and KCl solutions,
<https://pubmed.ncbi.nlm.nih.gov/38354641/>
64. Effect of brine time on quality of wet-salted fish - ResearchGate,
https://www.researchgate.net/publication/343305109_Effect_of_brine_time_on_quality_of_wet-salted_fish
65. The effect of storage temperature on the chemical and sensorial,
https://www.researchgate.net/publication/287549800_The_effect_of_storage_temperature_on_the_chemical_and_sensorial_quality_of_hot_smoked_Atlantic_bonito_Sarda_sarda_Bloch_1838_packed_in_aluminium_foil