

Comprehensive Analysis of Artisanal Dairy Systems: From Reproductive Physiology to Thermal Kinetics and Material Safety

The successful execution of artisanal dairy processing—particularly systems relying on low-technology interventions, hand-milking, and batch thermal pasteurization in wood-fired copper kettles—demands an uncompromising synthesis of veterinary obstetrics, microbiological hygiene, fluid thermodynamics, and material science. While industrial, high-throughput dairy manufacturing relies on automated, closed-loop systems constructed of passivated stainless steel, artisanal producers frequently operate at the intersection of historical methodology and modern food safety legislation. In such environments, the microbiological quality of the final product is inextricably linked to the holistic management of the bovine herd. This continuum begins long before lactation, deeply rooted in the reproductive endocrinology, transition nutrition, and obstetrical management of the dairy cow.

Following successful parturition, ensuring public health and product integrity requires the meticulous execution of validated pre-harvest sanitary protocols, a rigorous understanding of pathogenic thermal inactivation kinetics, and the precise management of food-contact materials to prevent toxicological contamination and lipid oxidation. This report provides an exhaustive, multi-disciplinary analysis of the artisanal milk processing continuum, from the foundational physiology of herd replenishment to the fluorimetric verification of thermal pasteurization.

Foundations of Lactation: Reproductive Physiology and Herd Management

The baseline somatic and microbiological quality of raw milk is profoundly influenced by the systemic health and reproductive efficiency of the herd. Achieving a steady supply of milk alongside consistent herd replenishment requires a comprehensive strategy that balances competing physiological demands, particularly in dual-purpose cattle such as the Fleckvieh, Normande, Shorthorn, and Brown Swiss, which are expected to deliver substantial lactation yields (ranging from 7,000 to 11,000 kg per lactation) while maintaining robust muscular frames¹.

Ovarian Dynamics and Estrous Cyclicity

The bovine female is a non-seasonal polyestrous animal, maintaining continuous reproductive cyclicity interrupted only by gestation or severe metabolic deficits¹. The estrous cycle spans an average of 21 days, regulated by a complex hypothalamic-pituitary-ovarian axis characterized by interdependent follicular waves and the cyclical synthesis and regression of the corpus luteum¹. Estrus (Day 0) is initiated by peak concentrations of estradiol secreted by the dominant preovulatory follicle, leading to a massive preovulatory surge of luteinizing hormone (LH) from

the anterior pituitary¹. Bovine follicular growth manifests in distinct biological waves, driven by transient elevations in follicle-stimulating hormone (FSH)¹. Observational tracking via transrectal ultrasonography reveals that the vast majority of bovine estrous cycles are composed of either two or three distinct follicular waves.

Parameter	Two-Wave Cycle	Three-Wave Cycle
Average Cycle Length	19.4 to 20.7 days	21.5 to 23.0 days
Wave 1 Emergence	Day 0 to 2	Day 0 to 2
Wave 2 Emergence	Day 9 to 11	Day 7 to 9
Wave 3 Emergence	N/A	Day 15 to 16
Ovulatory Follicle Origin	Wave 2	Wave 3
Preovulatory Growth Phase	9.8 days	5.0 days

Table 1: Bovine ovarian follicular wave dynamics and temporal phase lengths.

[cite: 1]

Understanding these waves is critical for the precise timing of artificial insemination (AI), which represents a profound genetic optimization and a vital biosecurity measure over natural service¹.

Pathogen Eradication via Low-Technology Artificial Insemination

Historically, reproduction relied on community-owned herd sires (the *Gemeindebulle*), which inherently posed severe biosecurity risks regarding venereal pathogens¹. The flagellated protozoan *Tritrichomonas foetus* and the gram-negative bacterium *Campylobacter fetus subsp. venerealis* asymptomatically colonize the epithelial crypts of the mature bull's prepuce, making the male a silent reservoir¹. During natural coitus, these pathogens ascend the female reproductive tract, colonizing the endometrium, inducing purulent pyometra, mid-term abortions, and causing pregnancy rates to plummet to 20%¹.

To circumvent this, low-technology artificial insemination utilizing liquid semen preservation is employed. When cryogenic liquid nitrogen is unavailable, semen is diluted in complex biochemical extenders and maintained at ambient or refrigerated temperatures¹. Extenders such as the Illini Variable Temperature (IVT) extender achieve this by bubbling carbon dioxide gas through a buffered solution to drop the pH to 6.3, exploiting the glyoxylate mechanism to

reversibly arrest sperm respiration at room temperature¹. Caprogen utilizes nitrogen gassing, caproic acid, and catalase to actively neutralize toxic hydrogen peroxide generated by lipid peroxidation¹. Utilizing the recto-vaginal method with sterilized reusable glass pipettes allows technicians to deposit this stabilized semen precisely into the uterine body, rapidly amortizing the operational costs of AI through superior genetic milk-yield dividends¹.

Nutritional Biochemistry and Postpartum Metabolic Health

Ensuring persistent high-volume lactation from dual-purpose breeds requires exacting nutritional precision. Nutritionists utilize dynamic frameworks, such as Metabolizable Energy (ME)

systems, calculating baseline maintenance mathematically as $0.515 \text{ MJ} \times BW^{0.75}$ ¹.

Transition Metabolism and Subclinical Pathologies

During early lactation, cows invariably enter a state of severe Negative Energy Balance (NEB) as volumetric rumen capacity lags behind the explosive caloric demands of lactogenesis¹. The cow mobilizes substantial adipose reserves, elevating circulating non-esterified fatty acids (NEFAs). Lipidomic profiling reveals that specific polar lipids, such as Phosphatidylethanolamine (PE 33:1), act as accurate biomarkers of this energetic deficit¹. Mismanagement of this transition leads to Subacute Ruminal Acidosis (SARA)—characterized by ruminal pH dropping below 5.8—and trace mineral antagonisms. For instance, high potassium from lush grasses depolarizes the ruminal epithelium, paralyzing active magnesium transport when the tetany ratio of $K^+ / (Ca^{2+} + Mg^{2+})$ exceeds 2.2 mEq/kg of dry matter¹.

Furthermore, the onset of lactation places an instantaneous drain on calcium reserves, requiring up to 23 grams of calcium to synthesize initial colostrum¹. When serum calcium concentrations fall below 2.14 mmol/L (8.5 mg/dL) without outward paralysis, the cow suffers from subclinical hypocalcemia (SCH)¹. Because calcium is an essential second messenger for immune cell activation, SCH severely impairs the phagocytic capabilities of polymorphonuclear leukocytes (PMNs)¹. Epidemiological data demonstrate that cows with SCH are 4.3 times more likely to develop metritis¹.

Uterine Involution and Endometritis Thresholds

The puerperium is dedicated to the physical involution of the massively hypertrophied uterus, which shrinks rapidly via myometrial contractions and the avascular necrosis of the maternal caruncles¹. Sloughing of these caruncular stalks generates physiological lochia. However, impaired immunity often leads to subclinical endometritis (SE), an insidious, localized inflammation of the endometrium drastically depressing conception rates¹. SE is diagnosed via endometrial cytology, calculating the ratio of inflammatory PMNs to normal epithelial cells.

Diagnostic thresholds dynamically shift from $> 18\%$ PMNs at 20-33 days postpartum to $> 5\%$ PMNs post-involution (>40 days)¹.

Obstetrical Biomechanics and Neonatal Immunity

Dystocia, affecting up to 5.6% of calvings, demands precise biomechanical intervention to prevent fetal hypoxia and severe maternal nerve damage¹. Traction must be applied using stainless steel obstetrical chains secured with a "double half-hitch" configuration to distribute tension and prevent metacarpal fractures¹. Instrumented studies utilizing the Calving Assistance Force Logger demonstrate that safe manual delivery generates a mean peak force of 95.4 kg, whereas mechanical extractors routinely exceed 188.6 kg, risking calving paralysis via the crushing of the obturator or sciatic nerves¹. In cases of fetal death and disproportion, a fetotomy utilizing a Thygesen's fetotome and obstetrical wire saw is systematically performed to amputate and extract the fetus, preventing catastrophic uterine rupture¹.

Following delivery, the survival of the neonate hinges entirely on the immediate transfer of passive immunity via colostrum¹. Due to the synepitheliochorial structure of the bovine placenta, calves are born entirely agammaglobulinemic¹. The neonatal intestine possesses a transient physiological window allowing enterocytes to absorb intact Immunoglobulin G (IgG) macromolecules via pinocytosis¹. This absorptive capacity severely blunts by 12 hours and ends in absolute "gut closure" by 24 hours¹. Calves must ingest a minimum of 3 to 4 liters of

high-quality colostrum ($> 50 \text{ g/L}$ IgG) within the first two hours, augmented by micronutrients like 3.0 ppm of selenium, which rapidly activates enterocyte pinocytosis to maximize IgG absorption¹.

Pre-Harvest Hygiene and Hand-Milking Sanitary Protocols

With the physiological foundations of lactation secured, the extraction of milk introduces the highest risk of microbiological contamination. The external bovine udder is continuously exposed to environmental pathogens, including coliforms, *Bacillus* spores, and environmental streptococci². Furthermore, the mammary tissue is vulnerable to contagious mastitis pathogens, most notably *Staphylococcus aureus* and *Streptococcus agalactiae*, which are mechanically transferred between animals during the milking process³. Preventing their entry into the raw milk supply necessitates a rigorous sanitary protocol.

Udder Preparation and Pre-Milking Disinfection

The physiological mechanism of milk let-down requires tactile stimulation seamlessly integrated into the hygiene routine. Following the removal of gross visible soils, a pre-milking teat disinfectant (pre-dip) is mandatory to rapidly oxidize environmental bacterial cell structures on the keratinized skin of the teat². Formulations utilizing sodium chlorite (e.g., 0.49 g/L activated base) or iodophor-based compounds are highly effective².

The pre-dip must completely coat the full length of the teat and remain in undisturbed contact for a minimum of 15 to 30 seconds to penetrate microscopic epidermal fissures⁶. Subsequently, the solubilized debris and disinfectant must be meticulously removed using a single-use, disposable paper towel or a dedicated cloth towel for each individual cow². The reuse of cloth towels across

multiple animals is a primary vector for disseminating contagious mastitis². The teats must be rendered entirely dry; milking wet teats facilitates the capillary transport of bacteria-laden moisture directly into the milk receptacle and drastically increases the frequency of liner slips in machine systems².

Foremilk Inspection and Subclinical Mastitis Detection

Prior to full-volume extraction, the technician must execute a foremilk procedure, manually expressing the first streams of milk into a specialized dark-surfaced strip cup⁶. Foremilking flushes the teat canal of the keratin plug and the highest concentration of resident bacteria². Diagnostically, the dark background allows the immediate visual identification of flaking, clotting, purulence, blood, or watery consistency⁴.

In subclinical cases lacking visual abnormalities, early-stage inflammation is detected via the California Mastitis Test (CMT). The CMT utilizes an alkyl aryl sulfonate reagent that lyses somatic cells (primarily PMNs) in the milk, precipitating their exposed DNA to form a gelatinous matrix⁹. The viscosity of this gel correlates directly to the somatic cell count (SCC); scores

indicating $> 500,000 \text{ cells/mL}$ mandate immediate segregation of the milk¹⁰. Identifying mastitic milk is vital, as it introduces massive bacterial loads and exhibits elevated plasmin activity that severely degrades subsequent dairy processing yields⁴.

Biomechanics and Hygiene of Hand-Milking

Hand-milking relies entirely on positive mechanical pressure. To prevent trauma to the highly vascularized internal mucosal lining (the rosette of Fürstenberg), the milker must employ the "full-hand" technique⁸. The milk is trapped in the teat cistern by closing the index finger and thumb at the base, followed by a sequential, downward squeezing motion with the remaining fingers to expel the fluid⁸. Stripping by pulling downward with the thumb and forefinger inflicts severe micro-tearing on epithelial tissues, predisposing the animal to opportunistic pathogen colonization⁸.

The milker's hands represent a highly mobile vector for cross-contamination³. The Food and Agriculture Organization (FAO) mandates that hands must be washed, thoroughly dried, and remain entirely dry throughout the procedure⁸. The historical practice of lubricating teats with milk, water, or saliva is a catastrophic hygienic failure⁸. To establish an absolute barrier—particularly against the transfer of human-derived *Staphylococcus aureus*—the use of clean, disposable latex or nitrile gloves is strongly advised³.

Post-Milking Disinfection and Sphincter Closure

Following milking, the teat sphincter muscle remains dilated and mechanically relaxed for up to 30 minutes, leaving the canal entirely unprotected³. The immediate application of a post-milking barrier teat dip is the single most critical intervention for preventing new contagious intramammary infections². Post-dips contain higher concentrations of germicides combined with emollient skin conditioners³. The solution coats the residual milk film, neutralizing lingering pathogens, while emollients promote epidermal healing and prevent microscopic cracking⁷. The cow should be encouraged to remain standing until the teat sphincter physiologically contracts⁸.

Particulate Exclusion and Raw Milk Regulatory Frameworks

Milk collected via hand-milking into open pails is highly susceptible to the incorporation of airborne dust, aerosolized manure, shed hair, and bedding particulates⁴. Immediate physical filtration is required using sterile, single-use, non-woven synthetic filter disks or finely meshed stainless-steel strainers.

However, it is crucial to recognize the microbiological limitations of physical filtration; while a filter removes visible organic matter, it does not remove the microscopic bacterial load suspended in the aqueous phase of the milk⁴. Filtration must be viewed strictly as a clarification step, not a microbiological intervention. To comply with stringent raw milk handling regulations—such as those outlined in the German Tier-LMHV (Lebensmittelhygiene-Verordnung) regarding direct-to-consumer raw milk (*Milch-ab-Hof*) and *Vorzugsmilch*—the milk must be protected at all stages¹². According to Annex 4 of the Tier-LMHV, *Vorzugsmilch* requires a total aerobic mesophilic plate count strictly capped at limits typically below 100,000 CFU/mL¹⁷. To guarantee broader public safety beyond these highly controlled, raw-certified outputs, the vast majority of milk must be subjected to stringent thermal treatments.

Thermal Pasteurization Kinetics in Wood-Fired Kettles

To render raw milk microbiologically safe for human consumption while preserving its nutritional and biochemical integrity, it undergoes thermal pasteurization. In artisanal settings, this is frequently executed as a Low-Temperature Long-Time (LTLT) batch process utilizing direct-fire mechanisms such as wood-fired copper kettles. This operation demands a profound understanding of microbiological destruction kinetics and fluid thermodynamics.

Microbiological Targets and Inactivation Kinetics

The stringency of modern milk pasteurization parameters is fundamentally based on the thermal resistance profile of *Coxiella burnetii*¹⁹. Historically, *Mycobacterium tuberculosis* was considered the most heat-resistant non-spore-forming pathogen in milk²⁰. However, following the identification of *C. burnetii* as the etiological agent of Q fever, extensive studies demonstrated that this obligate intracellular bacterium possesses superior thermal resilience²¹.

C. burnetii exists in two distinct morphological phases: the metabolically active Large Cell Variant (LCV) and the highly resistant, spore-like Small Cell Variant (SCV)²³. The SCV form exhibits profound resistance to osmotic stress, desiccation, ultraviolet radiation, and heat, allowing it to survive in raw milk and environmental aerosols²³. Consequently, *C. burnetii* was universally established by bodies such as the Codex Alimentarius as the index organism for pasteurization efficacy²¹.

The thermal death of microorganisms follows first-order logarithmic kinetics, defined by the D-value (Decimal Reduction Time) and the z-value (temperature increase required to reduce the D-value by a factor of 10)²². Modern kinetic data confirms these stringent requirements.

Kinetic Parameter	Value for <i>Coxiella burnetii</i>
Reference Temperature (T_{ref})	63°C (145°F)
D-value (D_{63})	~3.73 - 4.0 minutes
z-value	4.34°C - 6.4°C
Target Log Reduction	$\geq 5.0 \log_{10}$

Table 2: Thermal resistance parameters of *Coxiella burnetii* during milk pasteurization.
[cite: 19, 21, 23, 28, 29]

To achieve the universally mandated ≥ 5 -log reduction of *C. burnetii*, LTLT batch protocols dictate that every individual particle of milk must be heated to a minimum of 63°C and held continuously at or above that temperature for 30 minutes²⁰. Contemporary validations utilizing highly advanced Integrated Cell Culture-PCR (ICC-PCR) methodologies on Vero cell lines have superseded the necessity of historical in vivo guinea pig infection models, consistently proving that this time-temperature threshold provides a massive safety margin, often resulting in theoretical reductions exceeding 8 logs²¹.

Thermodynamics and Heat Transfer in Copper Kettles

Achieving precise, uniform temperatures in a wood-fired batch kettle presents extreme thermodynamic challenges. A wood fire generates intense, highly localized heat fluxes at the base of the copper kettle. The overall heat transfer relies on conduction through the highly conductive copper wall and forced convection into the bulk fluid³¹.

Milk is a complex biological fluid characterized by a specific heat capacity (C_p) of approximately 3.93 kJ/(kg · K) and a density (ρ) of roughly 1042 kg/m³ ³². Elevating the temperature of a large mass of milk requires massive energy inputs according to the thermodynamic equation $Q = m \cdot C_p \cdot \Delta T$ ³³.

When intense heat is applied to the copper boundary, a thermal boundary layer forms in the fluid immediately adjacent to the metal³². In a stagnant or poorly mixed vessel, heat transfers through this boundary layer primarily via conduction, which is highly inefficient due to the relatively low thermal conductivity of milk³². The fluid at the kettle wall can reach boiling temperatures—inducing severe Maillard browning, the denaturation of whey proteins, and the creation of scorched flavor profiles—while the geometric center of the vessel remains

dangerously below the 63°C lethal threshold³².

To destroy the thermal boundary layer and ensure uniform heat distribution, continuous and aggressive mechanical agitation is absolutely mandatory³². Agitation shifts the system from natural to forced convection. The efficiency of this convective heat transfer is characterized by

the Nusselt number (Nu), which relates the convective heat transfer to the conductive heat transfer across the boundary³². The Nusselt number is fundamentally dependent on the flow regime (Reynolds number, Re) and fluid properties (Prandtl number, Pr)³².

$$Nu = C \cdot Re^a \cdot Pr^b$$

By utilizing mechanical agitators, the processor dramatically increases the Reynolds number, creating turbulent vortices that constantly sweep the superheated milk away from the copper interface and fold it into the cooler bulk fluid³². This guarantees that no individual milk particle avoids the 63°C/30-minute requirement, securing microbiological safety without compromising organoleptic quality.

Material Science of Copper: Lipid Oxidation and Protein Degradation

The use of raw, uncoated copper kettles is a hallmark of certain traditional alpine dairy processes. Copper boasts extraordinary thermal conductivity, allowing for rapid and highly responsive heat transfer. However, copper is a highly reactive transition metal. When acidic or biologically complex fluids like milk come into direct contact with bare copper at elevated

temperatures, trace amounts of Cu^{2+} ions invariably migrate into the food matrix³⁸. This migration triggers a cascade of profound biochemical reactions that rapidly deteriorate the quality of the dairy product.

Lipid Auto-Oxidation and Hexanal Evolution

Milk fat consists largely of triglycerides containing polyunsaturated fatty acids (PUFAs), such as linoleic and linolenic acid⁴⁰. These PUFAs are inherently unstable and highly susceptible to lipid auto-oxidation—a free radical chain reaction consisting of initiation, propagation, and termination phases⁴¹.

The initiation phase requires the abstraction of a hydrogen atom from an allylic methylene group, a process that is typically slow at ambient temperatures⁴¹. However, Cu^{2+} acts as an extremely potent pro-oxidant catalyst⁴¹. Through Fenton-like redox cycling, trace copper rapidly decomposes existing lipid hydroperoxides into highly reactive alkoxyl and peroxy radicals, bypassing the high activation energy barrier of auto-oxidation⁴¹.

Once generated, these radicals abstract hydrogen from adjacent lipid molecules, propagating a violent chain reaction⁴¹. The alkoxyl radicals ultimately undergo homolytic β -scission, cleaving

the carbon chain to form secondary oxidation products, predominantly short-chain aldehydes and ketones⁴¹. The most prominent and widely tracked biomarker of this copper-induced degradation is **hexanal**⁴⁰.

Hexanal, alongside other volatile aldehydes like pentanal and heptanal, imparts highly undesirable organoleptic defects to the milk, universally described by sensory panels as "painty," "metallic," "cardboard," or "oxidized"⁴⁰. Analytical chemistry studies utilizing proton transfer reaction mass spectrometry (PTR-MS) demonstrate that spiking pasteurized milk with even trace concentrations of copper exponentially accelerates hexanal evolution; the mass fragment m/z 83 evolves with a sigmoidal trend exhibiting a maximum rate directly proportional to the Cu^{2+} content⁵².

While raw milk contains endogenous antioxidants—most notably alpha-tocopherol (Vitamin E) and ascorbic acid (Vitamin C)—these compounds are rapidly exhausted attempting to scavenge the flood of copper-induced free radicals, rendering the fluid defenseless against progressive rancidity⁴⁸. Furthermore, copper acts as a catalyst for protein oxidation. Cu^{2+} ions actively promote the oxidation of free sulfhydryl (-SH) groups in milk proteins, forcing the generation of undesirable intra- and inter-molecular disulfide bonds, which severely alter protein functionality, rheology, and flavor development⁴⁷.

Microbiology of Food Contact Surfaces: Biofilms and the Copper Paradox

Conversely to its negative impact on lipid stability, copper's profound chemical reactivity grants it exceptional antimicrobial properties, which heavily influence the colonization dynamics of dairy processing equipment. In typical dairy settings, residual milk proteins, fats, and lactose provide an ideal nutritional substrate for bacterial adhesion⁵⁷. Bacteria such as *Staphylococcus aureus*, *Bacillus* spp., and *Pseudomonas aeruginosa* rapidly adhere to surfaces, initiating a complex lifecycle that culminates in the excretion of an Extracellular Polymeric Substance (EPS)⁵⁷.

The EPS is a highly resilient, self-produced hydrogel matrix composed primarily of polysaccharides, structural proteins, and extracellular DNA (eDNA)⁵⁹. This three-dimensional architecture shields the embedded microbial micro-colonies from thermal stress, desiccation, and chemical disinfectants⁵⁹. Conventional sanitizers like sodium hypochlorite and peracetic acid struggle to penetrate a mature EPS matrix, rendering established biofilms nearly impossible to eradicate through standard Clean-In-Place (CIP) protocols⁶⁰. These biofilms act as persistent reservoirs, undergoing continuous cycles of maturation and dispersion, shedding massive pathogen loads into successive batches of milk and directly fueling high somatic cell counts and mastitis outbreaks if the contamination originates from the milking equipment⁵. Uncoated copper surfaces exhibit a profound oligodynamic effect against planktonic bacteria and early-stage biofilm attachment⁶⁷. Contact with elemental copper induces massive cellular stress in microbes; it compromises membrane integrity, causes an overwhelming influx of toxic ions, displaces native metals in metalloproteins, and generates lethal intracellular oxidative stress via Fenton reactions⁶⁸. Consequently, *S. aureus* and other prevalent dairy pathogens

demonstrate significantly reduced capacities to form mature biofilms on copper surfaces compared to standard AISI 304 stainless steel or polypropylene substrates⁶⁸.

However, this protection is not absolute. Recent microbiological research reveals that certain multi-species communities can develop a highly sophisticated "copper economy"⁷⁰. In mixed fungal-bacterial biofilms (e.g., *Candida albicans* alongside *S. aureus*), the organisms coordinate the up-regulation of copper-efflux pumps and copper-binding proteins, utilizing trace copper to actually strengthen the architectural integrity of the EPS matrix⁷⁰. Thus, while bare copper severely retards initial mono-species colonization, relying entirely on the metal's inherent toxicity in lieu of rigorous mechanical sanitation is a fundamental biological miscalculation.

Regulatory Compliance, Migration Testing, and Hot-Dip Tinning

Due to the toxicological risks associated with excessive heavy metal ingestion—and the severe organoleptic degradation caused by hexanal evolution—the use of bare copper in contact with fluid milk is heavily scrutinized under modern food safety legislation.

EU Frameworks and Specific Release Limits

In the European Union, all materials utilized in dairy processing are governed by the overriding Framework Regulation (EC) No 1935/2004⁷¹. Article 3 of this regulation dictates a strict principle of inertness: Food Contact Materials (FCMs) must be manufactured under Good Manufacturing Practices (GMP) such that, under normal or foreseeable conditions, they do not release constituents into food at levels that endanger human health, induce an unacceptable change in the food's composition, or deteriorate its organoleptic properties (smell and taste)⁷¹. Because bare copper demonstrably alters the organoleptic profile of milk via volatile aldehyde generation, its unrestricted use is fundamentally incompatible with the organoleptic clauses of 1935/2004. While Regulation (EU) No 10/2011 provides exhaustive Specific Migration Limits (SMLs) for plastics, harmonized EU legislation for metals and alloys remains absent³⁸. To address this legislative void, the Council of Europe issued Resolution CM/Res(2013)9, accompanied by a comprehensive technical guide³⁸. This resolution establishes Specific Release Limits (SRLs) for metallic ions migrating into food, tested utilizing rigorous simulants under DIN EN 13130 protocols⁸⁵.

Under CM/Res(2013)9, the maximum permitted SRL for copper migrating into food is strictly capped at **4.0 mg/kg**⁸⁰. In acidic environments or under high thermal loads (such as boiling dairy processing), bare copper can easily exceed this specific release limit, necessitating immediate material intervention³⁸.

Hot-Dip Tinning and Chemical Sanitation

To bridge the gap between the superior thermodynamic properties of copper kettles and the stringent inertness requirements of modern FCM regulations, artisanal equipment must undergo physical passivation. The globally recognized metallurgical solution is the application of a protective tin barrier.

Tin (Sn) is a highly ductile, non-toxic, and chemically stable metal⁹⁰. Applying a layer of pure tin over the copper substrate effectively isolates the reactive copper from the milk, completely

arresting copper-induced lipid auto-oxidation and eliminating the risk of exceeding the 4.0 mg/kg copper SRL. The most robust application method for heavy dairy processing equipment is hot-dip tinning⁹⁰.

In this process, the copper surface undergoes aggressive chemical pickling (using dilute hydrochloric or sulfuric acid) to remove existing copper oxides and biological slimes⁹⁵. The cleaned vessel is subsequently pre-heated, fluxed, and immersed in, or wiped with, molten tin⁹¹. The operating temperature of the tinning bath is strictly maintained between 260°C and 280°C⁹¹. At this elevated temperature interface, the molten tin reacts with the copper to form an

intermetallic compound (IMC) layer (primarily Cu_3Sn and Cu_6Sn_5) that metallurgically bonds the pure tin coating to the copper substrate, ensuring extreme mechanical durability against the friction of agitators⁹¹.

For food contact applications, the purity of the tin is of absolute paramount importance. Regulatory compliance dictates the exclusive use of "Food-Grade" pure tin, designated as **Sn99.9** (minimum 99.9% purity)⁹⁰. The use of lower-grade alloys is categorically prohibited, as they frequently contain high concentrations of lead (Pb), which is highly toxic and strictly regulated⁹⁶. CM/Res(2013)9 permits a generous Specific Release Limit of **100 mg/kg** for tin, reflecting its highly inert and non-hazardous nature when ingested in trace amounts⁸⁰.

Mitigating Amphoteric Corrosion

While the tin lining protects the milk, it introduces profound complexities into post-processing sanitation protocols. Pure tin is an amphoteric metal, rendering it highly susceptible to rapid corrosion when exposed to the aggressive chemical extremes typically used in dairy CIP systems (e.g., highly concentrated sodium hydroxide)⁹⁷.

Cleaning milk residues—which consist of baked-on fats and denatured proteins—requires an alkaline environment to achieve saponification and peptization. To safely clean tinned copper, formulators utilize highly buffered alkaline builders, most notably **sodium metasilicate** (

Na_2SiO_3)⁹⁷. Sodium metasilicate provides the necessary high pH and buffering capacity to emulsify milk fats, while possessing a unique chemical property: it actively deposits a microscopic, sacrificial silicate film across the surface of the tin and exposed copper⁹⁸. This silicate layer acts as a potent anodic corrosion inhibitor, preventing the alkaline hydroxide ions from aggressively dissolving the soft tin matrix⁹⁸. However, even with silicate inhibitors, prolonged contact times or allowing the metasilicate solution to dry will inevitably lead to the chemical etching of the tin lining, ultimately re-exposing the reactive copper substrate beneath⁹⁷.

Verification of Thermal Efficacy: Alkaline Phosphatase Assays

Once the batch pasteurization process has been concluded, the processor must empirically verify that the thermal application was sufficient to guarantee the destruction of *Coxiella burnetii*. Because culturing *C. burnetii* is notoriously difficult, time-consuming, and highly hazardous (requiring Biosafety Level 3 facilities), direct pathogen testing is useless for real-time production

release²⁴.

Instead, the global dairy industry relies on a biochemical proxy: the denaturation of an endogenous milk enzyme known as Alkaline Phosphatase (ALP)¹⁷. ALP is naturally present in high concentrations in the raw milk of mammals¹⁰². The immense utility of ALP lies in its precise thermal inactivation kinetics; the enzyme exhibits a heat resistance profile that is marginally higher than that of *Coxiella burnetii*¹⁰⁰. Therefore, if a thermal treatment successfully and completely denatures ALP, it serves as an absolute, fail-safe guarantee that the target bacterial pathogen population has been destroyed¹⁰¹.

Historically, crude colorimetric tests such as the Scharer Rapid Phosphatase Test (measuring indophenol formation) or the Aschaffenburg and Mullen Test (measuring nitrophenol) were utilized¹⁰⁰. However, these methods suffered from poor sensitivity thresholds, frequently failing to detect low-level raw milk contamination (0.1 to 0.2%)¹⁰⁰.

Today, the internationally recognized gold standard for quantifying residual ALP activity is the **ISO 11816-1** continuous fluorimetric direct kinetic assay (often commercialized as the Fluorophos Assay)¹⁷. In this protocol, a non-fluorescent aromatic monophosphoric ester substrate (specifically, 2'-[2-benzothiazolyl]-6'-hydroxybenzothiazole phosphate) is introduced into the pasteurized milk sample¹⁰⁶. If active ALP remains in the fluid—indicating either insufficient heating due to thermal boundary layer failures or post-pasteurization contamination with raw milk—the enzyme rapidly hydrolyzes the phosphate radical from the substrate¹⁰⁰. This cleavage yields a highly fluorescent product¹⁰⁶. The sample is incubated at 38°C, and a fluorimeter performs multiple kinetic readings over a 3-minute period to measure the rate of fluorescence emission, which is directly proportional to the enzymatic activity up to 7,000 mU/L¹⁰⁶.

Dairy Matrix	ALP Regulatory Limit for Successful Pasteurization	Detection Methodology
Bovine (Cow) Milk	≤ 350 mU/L	ISO 11816-1 (Fluorimetric)
Caprine (Goat) Milk	Assumed ≤ 300 mU/L (Lower native baseline)	ISO 11816-1 (Fluorimetric)
Ovine (Sheep) Milk	Assumed ≤ 500 mU/L (Higher native baseline)	ISO 11816-1 (Fluorimetric)

Table 3: Regulatory Alkaline Phosphatase (ALP) thresholds for thermal pasteurization verification across species.

[cite: 17, 102, 103]

Under ISO 11816-1 and overriding European Union directives, a bovine milk sample is legally classified as successfully pasteurized if the residual ALP activity is measured at ≤ 350 **milliunits per liter (mU/L)**¹⁷. Proper LTLT pasteurization at 63°C for 30 minutes in a well-agitated kettle will typically yield residual activities well below 50 mU/L¹⁰². It is critical to note that native ALP baseline levels vary significantly across species; caprine milk exhibits inherently lower levels, while ovine and asinine (donkey) milk levels are significantly higher, with raw donkey milk native ALP ranging wildly from 1758 to 6086 mU/L depending on days in milk (DIM)¹⁰². Nonetheless, for multi-species operations, utilizing the highly sensitive fluorimetric threshold prevents the catastrophic release of sub-lethally processed dairy into the public food chain¹⁰³.

Conclusion

The successful execution of artisanal dairy processing demands a rigorous, systems-level approach that harmonizes reproductive veterinary science, advanced nutritional biochemistry, and precise thermal engineering. The vulnerability of the raw milk supply necessitates stringent pre-harvest protocols deeply rooted in managing the physiological transition of the dairy cow, protecting the herd from metabolic cascades like subclinical hypocalcemia, and exploiting the narrow biological window of intestinal pinocytosis for neonatal immunity.

During extraction, minimizing the introduction of environmental and contagious pathogens (*Staphylococcus aureus*) requires unyielding adherence to udder preparation, utilizing pre-and post-milking disinfection with iodophor or sodium chlorite-based germicides, the detection of somatic cell counts via CMT, and the absolute avoidance of wet hand-milking. While physical filtration successfully excludes macroscopic particulates, it cannot remediate bacteriological loads, forcing a total reliance on meticulously validated thermal pasteurization.

Operating a wood-fired copper kettle necessitates profound mechanical intervention. Relying merely on external temperature application is insufficient; continuous mechanical agitation is required to increase the Reynolds and Nusselt numbers, stripping away the inefficient thermal boundary layer and driving forced convection. This guarantees that the bulk fluid satisfies the

strict LTLT parameters (63°C for 30 minutes) necessary to achieve a > 5 -log reduction of the highly heat-resistant intracellular pathogen *Coxiella burnetii*.

Crucially, the utilization of bare copper processing vessels poses severe biological and legal risks. While copper imparts an oligodynamic effect that initially retards the formation of Extracellular Polymeric Substance (EPS) biofilms, it simultaneously catalyzes massive lipid auto-oxidation via radical Fenton chemistry, producing sensory-destroying compounds like hexanal and directly violating the inertness principles of EU Regulation 1935/2004. Reconciling the thermal efficiency of copper with legislative safety requires hot-dip tinning utilizing high-purity, food-grade Sn99.9. This metallurgical barrier satisfies the specific release limits outlined in CM/Res(2013)9 while necessitating carefully calibrated sanitation protocols utilizing silicate-buffered alkaline cleaners to prevent amphoteric corrosion. Finally, the total integration of these myriad biological and mechanical systems is validated exclusively through the continuous fluorimetric ISO 11816-1 assay, confirming the total denaturation of Alkaline

Phosphatase and securing the ultimate biological safety of the product.

Works cited

1. 44 Bovine Reproduction And Herd Management.pdf
2. Milking Procedures on U.S. Dairy Operations, 2007 - USDA-Aphis, https://www.aphis.usda.gov/sites/default/files/dairy07_is_milkingproc.pdf
3. Milking Procedures on U.S. Dairy Operations - USDA-Aphis, https://www.aphis.usda.gov/animal_health/nahms/dairy/downloads/dairy02/Dairy02_is_MilkingProc.pdf
4. 3. milk hygiene - Milking, milk production hygiene and udder health, <https://www.fao.org/4/t0218e/T0218E03.htm>
5. Biofilm-Producing Ability of Staphylococcus aureus Obtained from, <https://www.mdpi.com/2306-7381/10/6/386>
6. END GAME SANITIZING PRE-MILKING AND POST MILKING TEAT, <https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=2767cd11-74cc-41f5-9031-8d1af559a58b&type=display>
7. Pre- and Post-Milking Teat Disinfection, http://www.reseaumammite.org/tactic/wp-content/uploads/2020/02/EN_teat_disinfection.pdf
8. How to Milk a Cow by Hand: A Hygienic Step-by-Step Guide, <https://www.zubairkhalid.com/knowledge/animal-farming/dairy-cattle/how-to-milk-a-cow>
9. Hygienic Milking Practices to Prevent Mastitis - CGSpace, <https://cgspace.cgiar.org/bitstreams/8eb4d9e7-b062-48cf-9c7d-0b9859d04cfd/download>
10. Using the California Mastitis Test (CMT) after calving - Teagasc, <https://teagasc.ie/news--events/daily/using-the-california-mastitis-test-cmt-after-calving/>
11. 5. systems of milking, <https://www.fao.org/4/t0218e/t0218e05.htm>
12. § 17 Tier-LMHV - Einzelnorm - Gesetze im Internet, https://www.gesetze-im-internet.de/tier-lmhv/_17.html
13. 17 Tier-LMHV – Abgabe von Rohmilch oder Rohrahm an Verbraucher, <https://lxgesetze.de/tier-lmhv/17>
14. Ab-Hof-Milch - Wikipedia, <https://de.wikipedia.org/wiki/Ab-Hof-Milch>
15. - 1 - 2. SCHRIFTTUM 2.1 Direktvermarktung von Rohmilch und, https://refubium.fu-berlin.de/bitstream/handle/fub188/8264/03_KAP2.PDF?sequence=4&isAllowed=y
16. Tier-LMHV - Verordnung über Anforderungen an die Hygiene beim, <https://www.gesetze-im-internet.de/tier-lmhv/BJNR182800007.html>
17. The use of alkaline phosphatase and possible alternative testing to, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8085980/>
18. Rohmilch in der Gastronomie, https://www.ua-bw.de/uploaddoc/cvuafr/GAP_2013_Poster_Rohmilch_in_der_Gastronomie_CVUA_Freiburg.pdf

19. Coxiella Burnetii DNA in Milk, Milk Products, and Fermented Dairy,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8775727/>
20. Can anybody explain the target organism for pasteurization?,
https://www.researchgate.net/post/Can_anybody_explain_the_target_organism_for_pasteurization
21. Coxiella burnetii and milk pasteurization: an early application of the,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC2870484/>
22. The Science Behind Milk Pasteurization - Agriculture Institute,
<https://agriculture.institute/milk-processing-packaging/science-behind-milk-pasteurization/>
23. Inactivation Kinetics of Coxiella burnetii During High-Temperature,
<https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2021.753871/full>
24. Quantitation of viable Coxiella burnetii in milk using an ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/26143937/>
25. Assessing the evidence for effective biosafety risk management in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12345007/>
26. Prevalence of Coxiella burnetii in unpasteurized dairy products in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11836485/>
27. Ultraviolet C inactivation of Coxiella burnetii for production of ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10993581/>
28. Inactivation Kinetics of Coxiella burnetii During High-Temperature,
<https://pubmed.ncbi.nlm.nih.gov/35069465/>
29. D-and z-values of Coxiella isolates M, WDK299, and WDK1188.,
https://www.researchgate.net/figure/D-and-z-values-of-Coxiella-isolates-M-WDK299-and-WDK1188_tbl2_357677707
30. Milk pasteurization with Plevnik devices,
<https://www.plevnik.eu/milk-pasteurization-with-plevnik-devices/>
31. 5.1 Heat transfer coefficient: exercise 2 (example) - TU Delft OCW,
<https://ocw.tudelft.nl/course-lectures/5-1-heat-transfer-coefficient-exercise-2/>
32. Determination of forced convection heat transfer coefficients ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6033811/>
33. Thermal Energy Consumption Assessment in a Fluid Milk Plant,
<https://www.scipublications.com/journal/ojes/article/thermal-energy-consumption-assessment-in-a-fluid-milk-plant-392>
34. Heat capacity - Medienportal Siemens Stiftung,
<https://medienportal.siemens-stiftung.org/view/104558>
35. THERMAL PROCESSING OF MILK - ResearchGate,
https://www.researchgate.net/profile/Anubhav-Pratap-Singh/publication/334326868_Thermal_Processing_of_Milk/links/60fd2e721e95fe241a88658b/Thermal-Processing-of-Milk.pdf
36. Design a VAT milk pasteurizer in Tanzania - Crimson Publishers,
<https://crimsonpublishers.com/aes/fulltext/AES.000525.php>
37. What is the Nusselt Number (Nu)? Formula & Guide - SimScale,
<https://www.simscale.com/docs/simwiki/numerics-background/what-is-nusselt-num>

- [ber/](#)
38. Metal kitchen utensils: Do substances transfer into food? - BfR,
<https://www.bfr.bund.de/en/opinions/metal-kitchen-utensils-do-substances-transfer-into-food/>
 39. Produkte: Bedarfsgegenstände mit Lebensmittelkontakt,
https://www.lgl.bayern.de/produkte/bedarfsgegenstaende/bg_lebensmittelkontakt/index.htm
 40. Correlating Volatile Lipid Oxidation Compounds with Consumer,
<https://www.mdpi.com/2076-3921/9/4/338>
 41. Oxidative Quality of Dairy Powders - Encyclopedia.pub,
<https://encyclopedia.pub/entry/15099>
 42. The antioxidant components of milk and their role in processing,
<https://www.veterinaryworld.org/Vol.12/January-2019/3.pdf>
 43. Oxidative Quality of Dairy Powders: Influencing Factors and Analysis,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8534860/>
 44. (PDF) Lipid Oxidation - ResearchGate,
https://www.researchgate.net/publication/247768996_Lipid_Oxidation
 45. OXIDATION OF LIPIDS IN FOOD Erwin W'sowicz¹, Anna Gramza¹,
<https://journal.pan.olsztyn.pl/pdf-98621-30276?filename=30276.pdf>
 46. Analytical Methods for Lipid Oxidation and Antioxidant Capacity in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8533275/>
 47. Light and copper catalysed oxidized flavours in stored milk,
https://www.researchgate.net/publication/230294500_Light_and_copper_catalysed_oxidized_flavours_in_stored_milk
 48. Protection against Oxidation of Fish-Oil-Enriched Milk Emulsions,
<https://pubs.acs.org/doi/10.1021/jf047960f>
 49. Influence of Oxygen and Copper Concentration on Lipid Oxidation in,
<http://lib3.dss.go.th/fulltext/Journal/J.AOCS/J.AOCS/1998/no.8/1998.vol75.no8.p1041-1046.pdf>
 50. (PDF) Oxidation of lipids in food - ResearchGate,
https://www.researchgate.net/publication/281317898_Oxidation_of_lipids_in_food
 51. Quantification of hexanal as an index of lipid oxidation in human,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3208008/>
 52. Hexanal as biomarker for milk oxidative stress induced by copper ions,
<https://pubmed.ncbi.nlm.nih.gov/28088419/>
 53. Antioxidant properties of Milk and dairy products, <https://d-nb.info/1180193040/34>
 54. Quantification of hexanal as an index of lipid oxidation in human,
https://www.researchgate.net/publication/51840945_Quantification_of_hexanal_as_an_index_of_lipid_oxidation_in_human_milk_and_association_with_antioxidant_components
 55. Excess copper catalyzes protein disulfide bond formation ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/36756756/>
 56. Excess copper catalyzes protein disulfide bond formation in ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10155707/>
 57. Preventing Biofilm Formation by Dairy-Associated Bacteria Using,

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC6608603/>
58. A] Understanding Biofilm In Dairy Processing Industry - BioDtex,
<https://biodtex.com/case-study/biofilms-in-dairy-processing-industry-understanding-prevention-and-control/>
 59. Temporal Profiling of Extracellular Polymeric Substances Associated,
<https://microbiologyjournal.org/temporal-profiling-of-extracellular-polymeric-substances-associated-with-biofilm-formation-in-bacteria-isolated-from-wound-infection/>
 60. (PDF) Biofilm formation by Staphylococcus aureus from food contact,
https://www.researchgate.net/publication/263322281_Biofilm_formation_by_Staphylococcus_aureus_from_food_contact_surfaces_in_a_meat-based_broth_and_sensitivity_to_sanitizers
 61. Microbial Biofilms in the Food Industry—A Comprehensive Review,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7922197/>
 62. Extracellular polymeric substances, a key element in understanding,
<https://www.aimspress.com/article/doi/10.3934/microbiol.2018.2.274?viewType=HTML>
 63. Biofilm - A threat to dairy industry - ResearchGate,
https://www.researchgate.net/profile/Vk-Ch/post/I_need_papers_about_the_effect_of_probiotics_on_biofilm_formation_on_stainless_steel_equipment_please/attachment/59d64459c49f478072ead06d/AS%3A273788673953792%401442287714405/download/39317-109146-1-PB%281%29.pdf
 64. Biosynthesis of Extracellular Polymeric Substances (EPS) and Its,
<https://www.pjoes.com/pdf-88051-21910?filename=Biosynthesis%20of.pdf>
 65. Biofilms in Milking Equipment | Dairy Farms | QualiTru,
<https://qualitru.com/biofilms-in-milking-equipment/>
 66. [PDF] Formation of biofilms on dairy equipment and the influence of,
<https://www.semanticscholar.org/paper/Formation-of-biofilms-on-dairy-equipment-and-the-of-Kukhtyn-Berhilevych/08ad12da17c42e8d814fb2d07d0462bad6f80fb8>
 67. Antimicrobial activity of copper surfaces against biofilm formation by,
<https://pubmed.ncbi.nlm.nih.gov/33279070/>
 68. Antimicrobial activity of copper surfaces against biofilm formation by,
https://www.researchgate.net/publication/346581070_Antimicrobial_activity_of_copper_surfaces_against_biofilm_formation_by_Salmonella_Enteritidis_and_its_potential_application_in_the_poultry_industry
 69. Silver and Copper Nanoparticles Inhibit Biofilm Formation by ... - MDPI,
<https://www.mdpi.com/2076-2615/11/7/1884>
 70. A 'copper economy' helps fungi and bacteria build better biofilms,
<https://microbiologysociety.org/resource/a-copper-economy-helps-fungi-and-bacteria-build-better-biofilms.html>
 71. EU Food Contact Products regulation - Trivium Packaging,
<https://www.triviumpackaging.com/resources/knowledge-articles-and-faqs/knowledge-articles/eu-food-contact-products-regulation>
 72. Legislation - Food Safety - European Commission,
https://food.ec.europa.eu/food-safety/chemical-safety/food-contact-materials/legislation_en

73. List of Products Covered by the EU FCM Framework Regulation (No, <https://www.compliancegate.com/eu-food-contact-material-framework-regulation-product-list/>)
74. Regulation (EC) No 1935/2004 on Materials and Articles Intended to, <https://powerrubber.com/en/blog/regulation-no-1935-2004-materials-and-articles-intended-to-come-into-contact-with-food>
75. BfR Recommendations on Food Contact Materials, <https://www.bfr.bund.de/en/service/databases/bfr-recommendations-on-food-contact-materials/>
76. BfR-Empfehlungen zu Materialien für den Lebensmittelkontakt, <https://www.bfr.bund.de/service/datenbanken/bfr-empfehlungen-zu-materialien-fuer-den-lebensmittelkontakt/>
77. Detailed explanation: BfR recommendations on food contact materials, <https://www.bfr.bund.de/en/service/frequently-asked-questions/topic/detailed-explanation-bfr-recommendations-on-food-contact-materials/>
78. Food contact materials - AGES, <https://www.ages.at/en/human/nutrition-food/food-information/food-contact-materials>
79. EU Food Contact Materials - EC 1935/2004 and EU 10/2011 - Intertek, <https://www.intertek.com/housewares-home-decor/food-contact-articles-testing/eu-food-contact-materials/>
80. Belgium Regulates Food Contact Metals and Alloys - SGS, <https://www.sgs.com/en-gb/news/2021/03/safeguards-03821-belgium-regulates-food-contact-metals-and-alloys>
81. Metals and alloys - used in food contact materials and articles, https://www.icoval.org/files/documentacion/documentos/metalls_alloys.pdf
82. Metals and alloys as a food contact material - Ruokavirasto, <https://www.ruokavirasto.fi/en/foodstuffs/food-sector/packages-and-other-food-contact-materials/question-and-answers/metals-and-alloys/>
83. CM/Res(2013)9 - CoE Search - The Council of Europe, <https://search.coe.int/cm?i=09000016805c8094>
84. Resolutions of the Council of Europe and Belgian Legislation, <https://chemtrust.org/wp-content/uploads/belgium-reach-fcm.pdf>
85. Migration Testing for Food Contact Materials (FCMs) - Intertek, <https://www.intertek.com/packaging/testing/migration-for-food-contact-materials/>
86. Product specific requirements: 101. Food contact article (plastic), https://manor.a.bigcontent.io/v1/static/101.Food_Contact_Materials_Plastics_V3.0
87. Compliance and migration tests on food packaging - Fraunhofer IVV, <https://www.ivv.fraunhofer.de/en/packaging/migration-tests.html>
88. WMF Group, https://dam.groupeseb.com/m/218b56ba217f78b8/original/18_9295_6030_KE_EN_3201018428.pdf?timestamp=20260312135239
89. View Source, <https://faq.edqm.eu/plugins/viewsource/viewpagesrc.action?pagelId=1376777>
90. Tin, tin ingot 99.9% pure tin Sn99.9% food grade - metallos.de,

- <https://www.metallos.de/en/p/tin-tin-ingot-99-9-pure-tin-sn99-9-food-grade>
91. Optimal Hot-Dipped Tinning Process Routine for the Fabrication of,
<https://www.mdpi.com/1996-1944/13/5/1191>
 92. Hot-dip tinning - Sofiamed, <https://sofiamed.com/product-sofiamed/hot-dip-tinning/>
 93. Tin Plating - Aurubis,
<https://www.aurubis.com/en/stolberg/products/rolled-products/tin-plating>
 94. (PDF) Optimal Hot-Dipped Tinning Process Routine for the,
https://www.researchgate.net/publication/339767223_Optimal_Hot-Dipped_Tinning_Process_Routine_for_the_Fabrication_of_Solderable_Sn_Coatings_on_Circuit_Lead_Frames
 95. Tinning process: Hot dipping (Corrosion control) - YouTube,
<https://www.youtube.com/watch?v=-K8Xb2tUWbY>
 96. Pure Tin Sn99.9 Granules - Zinn Gießerei Göhler,
<https://zinn-giesserei-goehler.com/en/produkt/pure-tin-sn99-9-granules/>
 97. Sodium Metasilicate | Solvents | Hydrite.com,
<https://www.hydrite.com/Product/Industrial/Sodium-Metasilicate.htm>
 98. Sodium Metasilicate In Cleaning | Uses, Benefits & Safety,
<https://evolvingelements.com.au/sodium-metasilicate-cleaning-uses-benefits/>
 99. Sodium metasilicate cleaner: strong, useful, but risky - Finishing.com,
<https://www.finishing.com/155/96.shtml>
 100. (PDF) The Application of Alkaline Phosphatase Assays for the,
https://www.researchgate.net/publication/49625937_The_Application_of_Alkaline_Phosphatase_Assays_for_the_Validation_of_Milk_Product_Pasteurization
 101. Validation of Milk Product Pasteurization by Alkaline Phosphatase,
<https://lupinepublishers.com/dairy-veterinary-science-journal/fulltext/validation-of-milk-product-pasteurization-by-alkaline-phosphatase-activity.ID.000113.php>
 102. Challenges of Using the ISO 11816-1 ALP Method for Pasteurization,
<https://www.mdpi.com/2304-8158/15/4/677>
 103. The use of alkaline phosphatase and possible alternative testing to,
<https://researchportal.ukhsa.gov.uk/en/publications/the-use-of-alkaline-phosphatase-and-possible-alternative-testing-/>
 104. Blog 4 Fluorophos FLM Pasteurisation Verification ALP Limits,
<https://www.qclscientific.com/blog-4-pasteurisation-verification-alp-limits/>
 105. ISO 11816-1 | IDF 155-1: 2024 - Determination of alkaline,
<https://shop.fil-idf.org/products/iso-11816-1-idf-155-1-2024-determination-of-alkaline-phosphatase-activity-part-1-fluorimetric-method-for-milk-and-milk-based-drinks>
 106. DIN EN ISO 11816-1 - 2014-03,
<https://www.dinmedia.de/en/standard/din-en-iso-11816-1/188208674>