

The Physics, Rheology, and Biochemistry of Traditional Gravity Cream Separation

Introduction to Polyphasic Dairy Emulsions and Historical Context

Bovine milk is an extraordinary biological fluid characterized by a highly complex, polyphasic architecture. Fundamentally, milk functions simultaneously as a colloidal dispersion of casein micelles and an oil-in-water emulsion consisting of liquid triglyceride droplets suspended within an aqueous plasma matrix¹. The inherent thermodynamic instability of this lipid emulsion is the foundational driver of cream separation. Left undisturbed in a quiescent state, the dispersed lipid phase naturally seeks to minimize its free energy by rising through the denser aqueous continuous phase, a macroscopic phenomenon that has been exploited by human dairy processors since antiquity¹.

Prior to the advent of continuous mechanical processing, the extraction of milk fat relied exclusively on manipulating this natural phase separation utilizing gravity, geometric containment, and ambient temperature controls⁴. Historical dairy practices, particularly those institutionalized during the 19th century through specific engineering methodologies like the Swartz and Cooley systems, utilized shallow setting pans and cold water or ice baths to deliberately accelerate the creaming process⁴. The literature from 19th-century dairy pioneers, such as Wilhelm Fleischmann in his seminal text *Lehrbuch der Milchwirtschaft* and Gustav Martiny, exhaustively detailed these early attempts to optimize the geometry of the "Flachschalen" (shallow pans) to maximize fat yield and minimize spoilage⁴.

While the contemporary global dairy industry universally employs high-speed centrifugal separators—originally pioneered in 1878 by the Swedish engineer Gustaf de Laval—to achieve near-instantaneous and highly efficient continuous fat extraction, gravity separation remains a critical subject of advanced study². Modern research into gravity separation heavily informs our understanding of emulsion rheology, food physical chemistry, phase fractionation, and the production of artisanal dairy products².

Understanding traditional gravity separation requires a rigorous, multidisciplinary synthesis. The macroscopic separation of cream from skim milk is governed strictly by the classical fluid dynamics described by Stokes' Law¹⁵. However, predicting the actual velocity of this separation in cold, raw milk introduces a profound physical paradox: the increasing viscosity of cold milk plasma should theoretically retard the rise of fat globules, yet historical methodologies intentionally utilized extremely cold temperatures to dramatically accelerate the process¹⁷. The resolution to this paradox is rooted not in physics, but in immunology. It lies in the biological phenomenon of cryoagglutination, wherein native immunoglobulins mediate the flocculation of microscopic fat globules into massive, highly buoyant clusters³. This report provides an exhaustive physical, biochemical, and operational analysis of traditional gravity cream separation, detailing the mechanics of the milk fat globule membrane (MFGM), the rheology of

milk plasma, the fractionating effects on fatty acids, and the severe microbiological implications inherent to shallow pan setting.

The Physical Chemistry and Fluid Dynamics of Gravity Separation

Stokesian Mechanics and Terminal Velocity Derivations

The rate at which a discrete, spherical particle rises or falls through a viscous, continuous fluid is not arbitrary; it is a highly predictable vector defined by the interplay of buoyant forces and frictional drag. In the context of dairy physical chemistry, the creaming of milk is fundamentally and mathematically described by Stokes' Law². The physical equation for the terminal rising

velocity (v) of a single, non-flocculated milk fat globule is expressed mathematically as:

$$v = \frac{2r^2(\rho_p - \rho_f)g}{9\eta}$$

Where:

- v represents the terminal upward velocity of the rising fat globule.
- r represents the geometric radius of the fat globule.
- ρ_p is the specific density of the continuous phase (the milk plasma or skim milk).
- ρ_f is the specific density of the dispersed phase (the milk fat).
- g is the acceleration constant due to gravity (9.81 m/s^2).
- η is the dynamic viscosity of the milk plasma surrounding the globule.

In raw bovine milk, the parameters governing this thermodynamic equation are highly specific and inherently variable based on composition and temperature. The mean density of milk fat (ρ_f) is approximately 0.93 kg/L (930 kg/m^3) at a standard temperature of 16°C ,

whereas the density of the surrounding skim milk plasma (ρ_p) is roughly 1.036 kg/L (1036 kg/m^3)²². Because the plasma density (ρ_p) is greater than the fat density (ρ_f), the term $(\rho_p - \rho_f)$ yields a positive integer, indicating that the buoyant force exceeds the downward gravitational pull, resulting in a positive upward velocity vector².

The most critical and heavily weighted variable within the Stokes' equation is the radius (r) of the fat globule, as the resulting terminal velocity scales directly with the square of the radius (r^2)⁶. Bovine milk fat is synthesized and secreted by the mammary epithelial cells as discrete,

spherical droplets that range wildly in diameter from $0.1\ \mu m$ to $15\ \mu m$, with the volumetric majority of the lipid mass concentrated in globules measuring between $4.0\ \mu m$ and $5.1\ \mu m$ ³. Human milk (HM) fat globules exhibit a similar primary distribution, generally measuring around $4.2\ \mu m$ to $5.1\ \mu m$ ²⁶. However, the presence of exceptionally large globules in bovine milk—some reaching up to $8.7\ \mu m$ or larger—profoundly alters the overall separation kinetics²⁷. Due to the exponential influence of the radius in the numerator of the equation, a large fat globule measuring $10\ \mu m$ in diameter will rise through the plasma exactly 100 times faster than a small globule measuring only $1\ \mu m$ in diameter⁶. This size-dependent velocity is the foundational reason why mechanical homogenization is utilized in modern liquid milk production. By forcing the raw milk through microscopic valves under extreme pressure, the homogenization process mechanically shears the fat globules, reducing their average diameter to approximately $0.5\ \mu m$ ¹⁵. At this highly reduced radius, the resultant Stokes velocity becomes so infinitesimally slow that the upward buoyant force is entirely overcome by the random, thermal Brownian motion of the fluid matrix, effectively and permanently preventing the formation of a cream layer¹⁵.

Rheological Properties and the Temperature-Viscosity Relationship

The denominator of the Stokes equation (9η) represents the frictional drag forces opposing the upward rise of the fat globule, and it is governed exclusively by the dynamic viscosity (η) of the plasma²⁸. The viscosity of a complex biological fluid like skim milk is highly temperature-dependent, a phenomenon that is critical to understanding historical processing methodologies¹⁸. As the thermal kinetic energy of the fluid decreases through cooling, intermolecular friction within the plasma increases exponentially, leading to a much higher dynamic viscosity¹⁹.

Furthermore, the viscosity is influenced by the total dry matter of the fluid, particularly the concentration and voluminosity of casein micelles and whey proteins¹⁹. Mathematical models, such as the Einstein and Eilers equations, are frequently utilized by rheologists to approximate the intrinsic viscosity based on these suspended solids, though they often overestimate the voluminosity of casein micelles at ambient temperatures³³. The actual dynamic viscosity of milk and cream exhibits distinct Newtonian behavior at high temperatures and low fat concentrations, transitioning to thixotropic (shear-thinning) behavior at low temperatures and high fat concentrations³⁴.

Fluid Matrix	Temperature (°C)	Dynamic Viscosity	Rheological
--------------	------------------	-------------------	-------------

		(mPa·s / cP)	Behavior
Skim Milk	50°C	1.05	Newtonian
Skim Milk	20°C	1.50	Newtonian
Skim Milk	5°C	2.40	Newtonian
Cream (18% fat)	20°C	6.00	Mildly Thixotropic
Cream (18% fat)	5°C	9.60	Thixotropic

Table 1: Viscosity of milk fractions as a function of temperature and fat content. The data demonstrates a strict inverse relationship between temperature and dynamic viscosity. [cite: 31, 32, 34, 35, 36, 37]

According to standard Stokesian fluid dynamics, chilling raw milk from ambient conditions down to 5°C increases the skim milk plasma viscosity to $2.4 \text{ mPa} \cdot \text{s}$, representing a 60% increase in drag forces compared to the $1.5 \text{ mPa} \cdot \text{s}$ viscosity observed at 20°C³⁵. Pure physical modelling using Stokes' Law would therefore predict that gravity creaming at 4°C should be dramatically slower than creaming at body temperature, as the increased viscosity would severely retard the terminal velocity of the rising globules²⁹.

However, historical dairy observations and modern empirical data definitively prove the exact opposite: fresh, raw bovine milk separated in cold, shallow setting pans creams exponentially faster, and much more completely, than warm milk²⁰. The actual velocity of the cream layer rising in cold, raw milk vastly exceeds the theoretical limits predicted by the Stokes equation for individual, discrete globules²⁰. The failure of Stokes' Law to accurately predict the cold-separation velocity of raw milk is not a failure of physics, but rather an omission of biology. Stokes' Law assumes that the fat globules act as independent, non-interacting spheres; in cold raw milk, this assumption is fundamentally violated²⁰.

The Biological Catalyst: Cryoagglutination and Immunoglobulin Interactions

The resolution to the apparent paradox of rapid cold-creaming lies deeply rooted in protein biochemistry and veterinary immunology. The highly accelerated creaming velocity observed in cold raw milk is driven almost entirely by a reversible biological phenomenon known as cold agglutination, or cryoagglutination³.

Immunoglobulin M (IgM) and Macroscopic Flocculation

Cryoagglutination is a strict temperature-dependent immunological reaction wherein specific innate proteins physically cluster individual, microscopic milk fat globules into massive, macroscopic aggregates³. Because Stokes' Law dictates that terminal velocity scales with the square of the particle's radius, fusing tens of thousands of microscopic fat globules into a

single, massive cluster instantly and radically transforms the effective radius (r) utilized in the equation. This massive, exponential increase in the buoyant force of the cluster heavily overrides the linear penalty introduced by the increased viscosity of the cold plasma, resulting in rapid vertical migration³.

The primary biological agent responsible for this clustering is Immunoglobulin M (IgM), a massive, pentameric antibody complex naturally synthesized by the bovine immune system and secreted into the raw milk matrix³. In mammalian biochemistry, IgM uniquely functions as a cryoglobulin—a protein complex that undergoes profound conformational changes and exhibits precipitation behavior at low temperatures³. The structural heavy chain of the IgM molecule folds into multiple distinct domains, allowing it to anchor and interact extensively with surrounding biological structures⁴³.

At temperatures approaching 4°C to 5°C, these large IgM cryoglobulins exhibit an extremely high binding affinity for specific structural glycoproteins anchored on the surface of the fat droplets²⁰. Advanced contemporary research utilizing confocal laser scanning microscopy (CLSM) and detailed proteomic analysis has confirmed that poly-immunoglobulin G receptors located on the outer membrane of the fat globule play a critical, highly conserved role in facilitating these binding sites⁴².

When the raw milk is rapidly chilled in a shallow pan, the pentameric IgM complexes act as structural bridges, physically cross-linking adjacent fat globules³. Furthermore, these immunoglobulins complex with native lipoproteins to strengthen the resulting matrix³⁸. The resulting agglutinated clusters can easily measure hundreds of micrometers in diameter. These massive, irregular clusters experience tremendous buoyant force and rapidly sweep upward through the dense, cold plasma, accelerating the macroscopic creaming process and leaving a distinct, sharply defined cream line at the surface⁴⁰.

Thermal Sensitivities, Reversibility, and Denaturation Kinetics

The binding affinity of the IgM complex to the poly-IgG receptors on the fat globule membrane is extraordinarily sensitive to the thermal environment of the fluid²⁰. The cryoagglutination process reaches its absolute maximum kinetic efficiency at temperatures of 4°C to 5°C⁴⁴. This specific biological temperature optimum aligns perfectly with the historical dairy processing protocols that utilized flowing spring water, deep-setting wells, or harvested ice to rapidly chill setting pans⁷.

However, if the ambient temperature of the milk is raised to 15°C, the agglutination behavior is markedly reduced, leading to smaller cluster sizes and a slower creaming rate⁴⁴. Elevating the temperature further to 20°C, or returning the milk to human or bovine body temperature (37°C), completely cancels any detectable interaction between the IgM complexes and the fat globules⁴⁴. At these warmer temperatures, the poly-immunoglobulin G receptors and the

pentameric IgM physically dissociate, the massive clusters disintegrate back into individual, discrete globules, and the creaming rate instantly plummets to align with the standard, slow parameters of Stokes' Law for independent particles⁴⁶.

Crucially, this low-temperature dissociation is an entirely reversible non-covalent interaction⁴⁶. If the milk is warmed to 37°C and then subsequently chilled back to 4°C, cryoagglutination will reoccur. However, the application of extreme heat introduces permanent structural damage. If raw milk is subjected to thermal pasteurization or heated above 60°C, the native tertiary and quaternary structures of the IgM proteins and the complementary membrane glycoproteins undergo irreversible thermal denaturation²⁵.

The decimal reduction time (D -value) for the heat denaturation of bovine immunoglobulins demonstrates that IgM is inherently more heat-sensitive than IgG or IgA⁵¹. Heating above this critical 60°C threshold permanently destroys the cryoglobulin mechanism; consequently, conventionally pasteurized or homogenized milk will never exhibit cold agglutination, fundamentally altering its long-term phase stability and preventing natural creaming during refrigeration²⁰.

Storage / Process Temperature	IgM-Receptor Interaction Status	Resulting Macroscopic Creaming Velocity
4°C – 5°C	Maximal cross-linking / Massive Agglutination	Highly Accelerated (Greatly exceeds Stokes' Law for single globules)
15°C	Markedly reduced binding	Moderate Velocity
20°C – 37°C	Complete dissociation	Slow (Strict adherence to Stokes' Law for discrete globules)
> 60°C	Irreversible protein denaturation	Permanently impaired cryoagglutination

Table 2: The profound thermal sensitivity of Immunoglobulin M (IgM) mediated cryoagglutination and its subsequent impact on milk fat creaming velocities.
[cite: 25, 44, 45, 46, 48, 51]

The Structural Architecture of the Milk Fat Globule Membrane (MFGM)

To fully comprehend the biochemical differences between cream harvested via traditional

gravity separation and cream extracted via mechanical centrifugation, one must perform an exhaustive analysis of the Milk Fat Globule Membrane (MFGM). The MFGM is the delicate, bioactive interface that separates the hydrophobic triglyceride core of the fat droplet from the hydrophilic aqueous plasma⁴¹.

Biogenesis and the Trilayered Phospholipid Architecture

The MFGM is a highly complex, 5 to 50 nanometer thick trilayered structure that coats every successfully secreted milk fat globule⁵⁵. Its unique architecture is a direct result of the biological mechanism of lipid secretion within the mammary epithelial cell²⁶.

During intracellular biogenesis, lipid droplets originate in the endoplasmic reticulum (ER). As they bud off from the ER, they are coated in an inner monolayer of specific polar lipids and associated proteins⁵⁸. These droplets then migrate to the apical pole of the epithelial cell. Upon secretion into the alveolar lumen, the droplets are pushed through the apical plasma membrane, enveloping the droplet in a true, outer cellular bilayer⁵⁵. Consequently, the intact MFGM consists of a highly specific inner polar lipid monolayer and an outer phospholipid bilayer, creating a robust trilayered architecture that physically stabilizes the emulsion and prevents spontaneous droplet coalescence in raw milk⁵³.

Highly Conserved Membrane Glycoproteins and Enzymes

Embedded deeply within this intricate phospholipid matrix is a vast array of highly specific, immune-active, and nutritional trans-membrane glycoproteins, peripheral proteins, and enzymes⁵⁶. Proteomic analysis has revealed that genes encoding for these MFGM proteins are highly conserved across mammalian species, indicating their profound evolutionary importance for neonatal nutrition and immunological defense²⁶.

The most prominent and biologically significant proteins anchored in the bovine MFGM include:

- **Xanthine Oxidase (XO) / Xanthine Oxidoreductase (XOR):** A massive 155 kDa enzyme that plays a critical dual role in neonatal antimicrobial defense (producing reactive oxygen species) and acting as a structural scaffolding protein for the membrane²⁶.
- **Butyrophilin (BTN):** A 67 kDa transmembrane glycoprotein that is an absolute prerequisite for successful lipid droplet secretion⁵⁹. Butyrophilin forms a tight structural complex with Xanthine Oxidase, often linked via intermolecular disulfide bonds, creating a rigid lattice that maintains the integrity of the MFGM⁶⁷.
- **Lactadherin (PAS 6/7):** A peripheral membrane glycoprotein critical for preventing viral infections in the neonate and facilitating the clearance of apoptotic cells⁶⁹.
- **Adipophilin and Fatty Acid-Binding Protein (FABP):** Proteins primarily located on the inner monolayer involved in intracellular lipid droplet assembly and fatty acid transport⁶³.

Prominent MFGM Protein	Molecular Weight (kDa)	Primary Biological/Structural Function
Xanthine Oxidase (XO)	155 kDa	Structural scaffolding; antimicrobial reactive

		oxygen generation.
Butyrophilin (BTN)	67 kDa	Transmembrane glycoprotein; facilitates droplet secretion; complexes with XO.
Lactadherin (PAS 6/7)	~47-53 kDa	Antiviral activity; apoptotic cell clearance.
Adipophilin	50 kDa	Intracellular lipid droplet assembly and stability.

Table 3: Primary protein constituents of the intact bovine Milk Fat Globule Membrane (MFGM) and their respective biological roles.

[cite: 59, 63, 64, 66, 69, 70]

Centrifugal Hydrodynamic Shear vs. Gravitational Preservation

The integrity of this delicate, trilayered MFGM is the primary casualty of modern mechanical processing. Continuous centrifugal separators operate at exceptionally high rotational

velocities, subjecting the milk to forces up to 10,000 times greater than standard gravity (g)⁷². The mechanical act of pumping the fluid and forcing it through the tight disc stacks of the separator centrifuge induces massive, violent hydrodynamic shear stress⁵³.

This intense mechanical shear inflicts profound physical damage upon the MFGM. Studies analyzing the zeta-potential and proteomic profile of centrifugally separated cream demonstrate that the mechanical forces forcibly strip away up to 24% of the native polar lipids from the membrane⁵³. Furthermore, the shear stress severs the delicate intermolecular disulfide bonds and dissociates critical peripheral and integral proteins—most notably butyrophilin and xanthine oxidase—removing them from the membrane architecture entirely and depositing them into the skim milk phase⁵³. The destruction of the MFGM exposes the naked, hydrophobic triglyceride core to indigenous milk lipases and dissolved oxygen, radically accelerating the rate of lipid auto-oxidation, premature free fatty acid release, and the subsequent rapid development of rancid off-flavors during storage⁶⁴.

Conversely, the passive, highly quiescent environment of a traditional gravity setting pan exerts virtually zero mechanical shear on the lipid emulsion⁴¹. The fat globules rise gently under the exclusive influence of natural buoyancy and cryoagglutination, leaving the trilayered MFGM entirely intact⁴¹. Creams harvested via traditional gravity separation exhibit vastly superior oxidative stability and maintain the full nutritional and immunological profile of the native butyrophilin and xanthine oxidase complexes, rendering the final artisanal product

fundamentally and biochemically distinct from its industrially centrifuged counterpart⁵⁴.

Traditional Methodologies: Shallow Pans, Deep Setting, and Devonshire Variations

Before the widespread adoption of centrifugal force in the late 19th and early 20th centuries, dairy practitioners systematically optimized the parameters of Stokes' Law and cryoagglutination through specific geometric and environmental interventions⁴. The primary historical methodology utilized globally for artisanal butter and cream production was the shallow pan setting system⁸.

Geometric Optimization and the Swartz/Cooley Methods

The architectural design of the historical setting pan was governed exclusively by the physical requirement to minimize the vertical distance the fat globules had to travel, thereby maximizing the total yield within a 24-hour period before widespread bacterial spoilage could render the milk unfit for consumption⁷⁹. Traditional setting pans were meticulously constructed as wide, flat-bottomed vessels, typically measuring 45 to 60 centimeters in diameter, with a shallow depth strictly limited to 2 to 4 inches (approximately 10 centimeters)⁴.

Fresh, warm milk, procured immediately following the hand-milking process, was poured into these pans to maximize the surface area-to-volume ratio⁷⁸. By forcibly restricting the depth to 10 centimeters, even the smaller fat globules located at the very bottom of the vessel required significantly less time to overcome the plasma drag and reach the surface¹⁶.

To fully exploit the rapid creaming velocities induced by IgM cold agglutination, these shallow pans were subjected to rapid cooling regimens¹⁶. In the late 19th century, this concept was formally commercialized in Germany, Scandinavia, and North America via the Swartz Method (utilizing shallow pans) and the Cooley System (utilizing submerged deep-setting cans), both of which mandated the submergence of the milk vessels into flowing spring water or harvested ice-water baths⁴.

By rapidly crashing the milk temperature down from 37°C to the optimal 4°C to 7°C window, the processor forcefully engaged massive IgM flocculation, allowing the bulk of the cream to separate and solidify at the surface within 12 to 24 hours¹⁶. After this requisite resting period, a specialized, perforated skimmer tool was utilized to manually lift and remove the dense, agglomerated lipid layer from the aqueous skim milk below⁴.

Organoleptic and Structural Variants: The Devonshire Protocol

While traditional pan separation relied heavily on cold ambient environments to trigger cryoagglutination, a distinctly different regional variation evolved in the United Kingdom, specifically in the counties of Devonshire and Cornwall, resulting in the production of highly prized clotted cream⁸¹.

The physics and thermal chemistry of the Devonshire protocol deviate significantly from standard Swartz-style gravity separation. Instead of cold setting, full-cream cow's milk is placed in traditional shallow pans and subjected to indirect, sustained heating⁸¹. The pans are floated over a bain-marie (water bath) or placed upon a steam grate, and the milk is slowly brought to a high scalding temperature, specifically avoiding allowing the fluid matrix to

achieve a rolling boil⁸³.

This deliberate application of high heat drastically alters the fluid dynamics and protein chemistry of the milk. The extreme elevation in temperature sharply reduces the dynamic

viscosity of the plasma to roughly $1.05 \text{ mPa} \cdot \text{s}$, increasing the un-agglutinated terminal velocity of the fat globules according to Stokes' Law, even while completely destroying the IgM cryoagglutination mechanism².

Simultaneously, the sustained heat denatures the primary whey proteins dissolved in the

plasma, particularly β -lactoglobulin⁶⁹. As these specific whey proteins denature, they unfold and chemically co-precipitate with the rising fat droplets and casein micelles, creating a dense, highly stable, interwoven matrix at the fluid surface⁸¹. The pans are subsequently transferred to a cold environment to cool slowly over many hours, permanently solidifying this complex matrix into a thick, buttery, and characteristically nutty crust⁸¹. Unlike raw, cold-separated cream, which relies purely on intact fat globule clustering, Devonshire cream is a structurally modified composite of cooked fat, denatured whey proteins, and casein, yielding a fundamentally different rheological and sensory profile⁸⁷.

Distribution Kinetics, Phase Fractionation, and Skimming Efficiency

Skimming Inefficiency and Residual Lipid Fractions

The rapid, global transition from gravity separation to centrifugal separation in the modern dairy era was driven almost entirely by the comparative skimming efficiency of the two methods⁸⁸. Efficiency in commercial cream separation is strictly inversely proportional to the amount of residual fat left behind in the skim milk matrix⁹¹.

Gravity separation acts as a slow, natural particle size fractionator. Because the upward velocity is dictated by the square of the globule radius, the largest, most buoyant fat globules ($> 5 \mu\text{m}$) rapidly migrate to the surface⁶. Consequently, the top layer (the harvested cream) is heavily concentrated with large-diameter globules, exhibiting a unimodal size distribution skewed heavily toward the upper geometric limits²⁷.

However, the smallest fat globules—those measuring less than $1 \mu\text{m}$ in diameter—possess such a minute radius that their intrinsic buoyant force is nearly negligible⁸⁶. Furthermore, these micro-globules possess too small of a surface area to effectively participate in massive IgM agglutination networks, allowing them to remain permanently suspended in the bottom fraction due to the overriding force of random Brownian motion³.

As a direct result of these physics, even under optimal cold-setting conditions utilizing ice baths for 24 to 48 hours, gravity separation invariably leaves a highly significant residual fat content in the skim milk, typically measuring between 0.1% and 0.5% (or higher, depending on the breed and initial globule size distribution of the herd)⁸⁹. In traditional, PDO-protected cheese-making processes that still legally mandate gravity separation, such as the production

of Grana Padano and Parmigiano-Reggiano, this residual fat is not considered a loss, but rather an acceptable and strictly necessary compositional component of the raw material⁹⁸.

In stark contrast, a continuously operating mechanical centrifugal separator bypasses the limitations of Brownian motion entirely. The extreme application of centripetal force overwhelmingly supersedes the small radius of the micro-globules, driving almost the entirety of the lipid content toward the rotational axis with extreme prejudice¹⁰⁰. A well-calibrated, modern centrifugal separator will leave only 0.01% to 0.07% residual fat in the skimmed milk, recovering nearly 100% of the available commercial butterfat²⁴.

Separation Methodology	Primary Motive Force	Required Processing Time	Typical Residual Fat in Skim Milk
Traditional Shallow Pan	Standard Gravity ($1g$) + IgM Agglutination	12 to 48 hours	0.10% – 0.50%+
Centrifugal Separator	Centripetal Acceleration ($> 5000g$)	Continuous Flow / Minutes	0.01% – 0.07%

Table 4: Empirical comparison of skimming efficiency and operational parameters between traditional gravity setting and continuous centrifugal separation.

[cite: 16, 24, 73, 88, 89, 95, 100]

Fractional Sorting of Specific Fatty Acids

Because gravity separation partitions the milk strictly based on globule size, it inherently alters the biochemical profile of the resulting fractions. Smaller fat globules possess a much higher surface-area-to-volume ratio than large globules. Because the MFGM contains high concentrations of polar lipids, and because certain fatty acids are preferentially located within or near the membrane, the segregation of small globules into the bottom skim fraction alters the lipidomics of the system⁶⁹.

Extensive gas chromatography analysis of gravity-separated fractions demonstrates that the bottom fraction (the skim milk) retains disproportionately higher concentrations of highly desirable, bioactive polyunsaturated fatty acids⁴⁷. Specifically, the micro-globules left behind in gravity-separated skim milk exhibit significantly elevated levels of oleic acid, linolenic acid, arachidonic acid, and Conjugated Linoleic Acid (CLA) compared to the macroscopic globules harvested in the top cream⁴⁷. The preservation of these specific fatty acids within the skim milk fraction contributes heavily to the unique nutritional and ripening profiles of raw-milk cheeses produced via traditional gravity methods.

Microbiological Vulnerabilities and Somatic Cell Co-Creaming

While the physical chemistry of the shallow pan method successfully isolates the lipid fraction while perfectly preserving the delicate MFGM, the prolonged 12 to 48-hour setting time required introduces severe, highly complex microbiological vulnerabilities. The raw milk matrix is a nutrient-dense, highly hospitable biological medium, and its microbiological profile undergoes dynamic, threatening shifts over the course of the gravity separation period⁷⁸. The microbiological outcome of the final cream is inextricably linked to the initial baseline somatic and hygienic quality of the herd, deeply rooted in the transition nutrition and obstetrical management of the dairy cow prior to milking.

The Co-creaming Phenomenon: Somatic Cells and Bacterial Flora

As the massive, IgM-mediated fat clusters rise slowly through the milk plasma, they do not travel through an empty void; they act as a massive, biological filtration net, mechanically sweeping up suspended biological particulates in the fluid. Advanced research into the gravitational separation of raw milk reveals a high degree of co-creaming, a phenomenon where indigenous bacteria, bacterial spores, and somatic cells (predominantly leukocytes, neutrophils, and shed epithelial cells from the udder) become physically entangled within or directly agglutinate to the rising fat droplets²⁰.

Microbiological and cytological observations confirm that up to 50% to 80% of the total baseline bacterial load and somatic cell count is physically relocated from the bulk plasma into the top cream fraction during the gravity separation process¹⁰⁵. While this mechanism acts as a rudimentary clarifier for the resulting skim milk—effectively reducing its somatic cell and bacterial count, rendering it superior for certain cheese-making applications—it concurrently concentrates extreme, potentially hazardous microbiological loads directly into the cream⁹⁹. In modern centrifugal systems, this physical reality is managed efficiently by the extreme centripetal forces. Centrifuges push the denser biological particles (like heavy somatic cells, spores, and dirt) outward to the extreme periphery of the bowl wall, discharging them as dense "separator sludge," effectively clarifying both the cream and the skim milk simultaneously¹. In historical gravity systems, however, this dense bacterial and somatic load remains inextricably trapped within the harvested cream, severely limiting its shelf life and complicating downstream processing into butter.

Psychrotrophic Spoilage and Thermostable Lipolytic Degradation

The absolute reliance on cold temperatures (4°C to 7°C) to optimize Swartz-method separation creates a highly selective microbiological environment within the shallow pan¹⁶. While rapid chilling effectively arrests the proliferation of standard mesophilic lactic acid bacteria (such as *Lactococcus* species, which would otherwise rapidly sour the milk), it actively and dangerously selects for psychrotrophic (cold-thriving) microorganisms¹⁰³.

If raw milk is allowed to stand in a shallow pan for 24 to 48 hours, psychrotrophic bacteria—predominantly gram-negative rods from the environmental genera *Pseudomonas* (especially *P. fluorescens*) and *Acinetobacter*, alongside potentially pathogenic strains like

Yersinia enterocolitica—will undergo rapid, exponential growth¹⁰⁹. While the vegetative cells of these bacteria themselves may ultimately be destroyed by subsequent thermal pasteurization or cooking, their prolific metabolic byproducts represent a catastrophic threat to dairy quality⁶⁴.

During their extended cold-incubation period in the setting pan, psychrotrophic bacteria synthesize and excrete massive volumes of extracellular lipases and proteases into the plasma⁶⁴. Unlike native, endogenous milk lipase (LPL), these psychrotrophic enzymes are uniquely and severely thermo-stable; they easily survive standard commercial pasteurization parameters that destroy the bacteria that produced them⁶⁴. Once the cream is harvested, these highly resilient, residual enzymes immediately begin to hydrolyze the ester bonds of the triglycerides, releasing short-chain free fatty acids (such as butyric, caproic, and caprylic acid) that impart highly objectionable rancid, soapy, and bitter organoleptic defects to the final cream or churned butter⁶⁴.

Therefore, the successful execution of traditional gravity separation requires significantly more than just precise thermal management to induce cryoagglutination. It demands an uncompromising, absolute standard of pre-harvest veterinary hygiene and strict sanitary milking protocols to minimize the initial environmental psychrotrophic inoculation of the raw milk before it ever enters the setting pan¹¹³.

Conclusion

The traditional extraction of cream via gravity separation in cold, shallow pans represents a profound, highly nuanced intersection of fluid mechanics, protein biochemistry, and applied thermodynamics. While the fundamental upward trajectory of the milk fat globule is dictated entirely by the precise density differentials and plasma viscosity matrices outlined mathematically in Stokes' Law, purely physical models inherently fail to account for the astonishing speed of separation achieved in historical dairy practices. The methodology functions efficiently only because it brilliantly exploits the biological mechanism of cryoagglutination. By rapidly chilling the milk to 4°C, 19th-century dairy processors utilizing the Swartz and Cooley methods inadvertently activated native Immunoglobulin M (IgM), which binds aggressively to the poly-IgG receptors on the Milk Fat Globule Membrane (MFGM), flocculating individual microscopic droplets into massive, buoyant clusters that easily overcome the viscous drag of the cold plasma.

The engineered geometry of the shallow pan minimized the required transit distance, maximizing yield over a 24-hour setting period. Although high-speed centrifugal separation has entirely superseded this methodology in industrial applications due to its unparalleled ability to recover fat down to 0.01% residual levels, the intense hydrodynamic shear forces of modern processing cause severe, irreversible structural damage to the MFGM, forcefully stripping vital glycoproteins like butyrophilin and xanthine oxidase from the lipid interface. Gravity separation, by contrast, relies entirely on quiescent buoyant forces, perfectly preserving the complex trilayered architecture of the fat globule and fractionally preserving highly desirable polyunsaturated fatty acids in the skim milk.

However, this gentle, prolonged cold-setting methodology inherently concentrates somatic

cells and exposes the nutrient-dense matrix to the severe, highly destructive threat of psychrotrophic bacterial spoilage and thermostable lipolysis. Thus, traditional gravity separation remains a chemically superior but microbiologically vulnerable process, requiring exacting environmental control, exhaustive understanding of emulsion rheology, and exceptional raw milk hygiene to execute successfully.

Works cited

1. Cream Separation Techniques in Dairy Production | PDF - Scribd, <https://de.scribd.com/document/467141341/dairy-re>
2. Centrifugal separators and milk standardization, <http://dairyprocessinghandbook.tetrapak.com/chapter/centrifugal-separators-and-milk-standardization>
3. Cold agglutination: mechanism and role - indian farmer, <https://indianfarmer.net/uploads/cold%20agglutinatn.pdf>
4. Workings of Different Types of Small Cream Separators | Slavic Beauty, <https://before.slavicbeauty.net/dairy/workings-of-different-types-of-small-cream-separators/>
5. Equipment History and Function | Conococheague Stainless LLC, <https://www.cstainlessllc.com/equipment-history-and-function>
6. Methods and Factors in Milk Separation - Agriculture Institute, <https://agriculture.institute/milk-processing-packaging/milk-separation-methods-factors/>
7. History of the dairy industry - Internet Archive, <https://archive.org/download/historyofdairyin00pirt/historyofdairyin00pirt.pdf>
8. The Book of the Farm: Detailing the Labours of the Farmer, Farm, <https://dokumen.pub/the-book-of-the-farm-detailing-the-labours-of-the-farmer-farm-steward-ploughman-shepherd-hedger-farm-labourer-field-worker-and-cattle-man-4-4nbsped.html>
9. Tin cream skimmer - McLeod County Historical Society, <https://mcleodhistory.pastperfectonline.com/webobject/A6D0588C-76D0-4389-BC93-842219161780>
10. Nummer 02/15 7 januari 2015 - RVO, https://www.rvo.nl/sites/default/files/octrooiportal/2015/01/IE_02_2015.pdf
11. Lehrbuch der Milchwirtschaft - Wilhelm Fleischmann - Google Books, https://books.google.com/books/about/Lehrbuch_der_Milchwirtschaft.html?id=9lA10gEACAAJ
12. Lehrbuch der Milchwirtschaft - Wilhelm Fleischmann - Google Play, https://play.google.com/store/books/details/Lehrbuch_der_milchwirtschaft?id=mLrF44BQmUC&hl=is
13. Catalog Record: Lehrbuch der Milchwirtschaft, <https://catalog.hathitrust.org/Record/006911327>
14. The Beginnings of Modern Milk Age in Germany, <https://d-nb.info/1140118900/34>
15. Stokes' Law: Milk Fat Separation Physics - Watson Dairy Consulting, <https://dairyconsultant.co.uk/si-stokeslaw.php>

16. The Principle of Cream Separation: Understanding Gravity and,
<https://agriculture.institute/dairy-products-i/principle-cream-separation-gravity-mechanical/>
17. What Temperature Should Milk Be for a Cream Separator?,
<https://wildoaktrail.com/blogs/off-grid-living/milk-temperature-cream-separation>
18. Dynamic Viscosity (mPas) for Liquids, Oils & Chemicals,
<https://www.behaelter-kg.de/en/insights/what-is-the-viscosity-of-my-medium.html>
19. What is the Viscosity of Milk? - Q-Pumps,
<https://www.q-pumps.com/noticias/what-is-the-viscosity-of-milk>
20. Effect of temperature on the microstructure of fat globules and the,
https://www.researchgate.net/publication/322911115_Effect_of_temperature_on_the_microstructure_of_fat_globules_and_the_immunoglobulin-mediated_interactions_between_fat_and_bacteria_in_natural_raw_milk_creaming
21. The Effect of Milkfat Fraction Melting Properties on Physical,
<https://vtechworks.lib.vt.edu/bitstreams/c0b511ea-2d6d-4006-98ec-eafb8086d5e9/download>
22. Principle of Separation - Dairy Technology,
<http://dairy-technology.blogspot.com/2014/01/principle-of-separation.html>
23. Cream Separation Calculator | DairyCalc,
<https://dairycalc.in/calculators/milk/cream-separation-calculator/>
24. Understanding and Minimizing Fat Losses in Skim Milk During,
<https://agriculture.institute/dairy-products-i/minimize-fat-loss-skim-milk-cream-separation/>
25. Advancement on Milk Fat Globule Membrane: Separation ... - Frontiers,
<https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2021.807284/full>
26. Milk Fat Globules: 2024 Updates - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11521418/>
27. microstructure, rheological and geometrical properties of fat globules,
<https://journal.pan.olsztyn.pl/pdf-97931-30527?filename=30527.pdf>
28. cream separation from milk - GeoCities.ws,
<http://www.geocities.ws/talha/files/CREAM%20SEPARATION%20FROM%20MILK.pdf>
29. Factors affecting the creaming of human milk - CORA,
<https://cora.ucc.ie/bitstreams/dfb2f827-063c-444e-8b5b-be05708904a3/download>
30. Factors affecting the creaming of human milk - ResearchGate,
https://www.researchgate.net/publication/341182997_Factors_affecting_the_creaming_of_human_milk
31. Overview viscosities of liquids - INDAG Maschinenbau GmbH,
<https://indag.com/en/news/news-detail/uebersicht-viskositäten-von-flüssigkeit-en>
32. Viscosity of liquids - buerke.de,
https://www.buerke.de/assets-original/wissenswertes/viscosity_en.pdf

33. Effect of Temperature on the Viscosity Of Skimmilk - SciSpace,
<https://scispace.com/papers/effect-of-temperature-on-the-viscosity-of-skimmilk-3tzbrx523v>
34. APPROXIMATE VISCOSITIES OF SOME COMMON LIQUIDS,
<https://quillinox.com/docs/tabla-viscosidad.pdf>
35. Viscosity in the Food & Beverage Industry | AxFlow,
<https://www.axflow.com/support/engineering-toolbox/viscosity-in-the-food--beverage-industry/>
36. Relationships between Viscosity and the Contents of,
<https://www.hrpub.org/download/20160930/FST1-11107328.pdf>
37. regression models for density and viscosity of ultrafiltration milk,
<https://www.agrojournal.org/14/06-04-08.pdf>
38. Milk Composition and Nutritional Value | PDF | Milk | Fat - Scribd,
<https://www.scribd.com/document/985489440/Composition-Nutritive-Value-of-Milk>
39. Cold Agglutination: The Real Reason Cream Rises Fast in Raw Milk,
<https://getrawmilk.com/content/cold-agglutination-raw-milk-creaming>
40. Milk Lipids- Chemical Properties; Physical Properties; Structure and,
<https://books.lib.uoguelph.ca/dairyscienceandtechnologyebook/chapter/milk-lipids-chemical-properties-physical-properties-structure-and-fat-globules-function-al-properties/>
41. Effect of fat globule size on functionality and stability of dairy emulsions,
https://espace.library.uq.edu.au/view/UQ:bd80bb7/s4401287_phd_finalthesis.pdf
42. Thermal effects on IgM-milk fat globule-mediated agglutination,
<https://www.cambridge.org/core/journals/journal-of-dairy-research/article/thermal-effects-on-igm-milk-fat-globule-mediated-agglutination/566ED4F4A0A2F03093DB729FB2815F8A>
43. The structure of membrane bound and secreted immunoglobulin,
https://www.researchgate.net/figure/The-structure-of-membrane-bound-and-secreted-immunoglobulin-M-slgM-A-A-monomer_fig2_345148511
44. Thermal effects on IgM-milk fat globule-mediated agglutination,
https://www.researchgate.net/publication/329452086_Thermal_effects_on_IgM-milk_fat_globule-mediated_agglutination
45. Thermal effects on IgM-milk fat globule-mediated agglutination.,
<https://scispace.com/papers/thermal-effects-on-igm-milk-fat-globule-mediated-1z0sadgqtu>
46. Thermal effects on IgM-milk fat globule-mediated agglutination,
<https://pubmed.ncbi.nlm.nih.gov/30520391/>
47. Fatty acid composition of the bovine milk fat globules obtained by,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20173175974>
48. Thermal effects on IgM-milk fat globule-mediated agglutination - Pure,
<https://pure.au.dk/portal/en/publications/thermal-effects-on-igm-milk-fat-globule-mediated-agglutination/>
49. Skimming cream hi-res stock photography and images - Alamy,
<https://www.alamy.com/stock-photo/skimming-cream.html>

50. Abstract #T56, <https://m.adsa.org/2019/abs/t/78173>
51. Kinetic and Thermodynamic Parameters for Heat Denaturation of, https://www.researchgate.net/profile/Houda_Kawas/post/ask_for_a_standard_He_at_denatured_IgG_protocol/attachment/59d6420979197b807799ddc6/AS:437330336063488@1481279085264/download/56dff1b708aee77a15fde49d.pdf
52. Effect of temperature on the microstructure of fat globules ... - PubMed, <https://pubmed.ncbi.nlm.nih.gov/29398025/>
53. Bacterial Removal Separation with a Reduced Throughput Adapted, https://serviss.bib.hs-hannover.de/files/3455/wente_etal2024-bacterial_removal_separation.pdf
54. Bacterial Removal Separation with a Reduced Throughput Adapted, <https://www.mdpi.com/2624-862X/5/4/48>
55. Structural and physical-chemical properties of milk fat globules, https://nano.ku.dk/english/research/structural-food-physics-and-soft-matter-self-assembly/pdfs/jjkk_107.pdf
56. Milk fat globule membrane in early-life nutrition - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13323632/>
57. Stability of milk fat globule membrane proteins toward ... - PubMed, <https://pubmed.ncbi.nlm.nih.gov/22541459/>
58. Composition, Properties and Nutritional Aspects of Milk Fat Globule, <https://journal.pan.olsztyn.pl/pdf-98286-31039?filename=31039.pdf>
59. Investigations of health benefits of milk fat globule membrane lipid, https://doras.dcu.ie/16641/1/Anna_Kuchta_thesis.pdf
60. Emulsion Structural Remodeling in Milk and Its Gelling Products - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11507225/>
61. The Immunological Role of Milk Fat Globule Membrane - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9655658/>
62. Milk Fat Globule Proteins Are Relevant Bovine Milk Allergens in, <https://iris.unito.it/retrieve/33c5420e-ca1f-44f9-afa8-4bcbcd7b2746/Milk%20Fat%20Globule%20Proteins%20Are%20Relevant%20Bovine%20Milk%20Allergens%20in%20Patients%20with%20CE%20B1%E2%80%90Gal%20Syndrome.pdf>
63. Molecular Remodeling of Milk Fat Globules Induced by Centrifugation, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12732456/>
64. Dairy Fat Products and Functionality: Fundamental Science and, <https://dokumen.pub/dairy-fat-products-and-functionality-fundamental-science-and-technology-1st-ed-9783030416607-9783030416614.html>
65. Lactoferrin effects on immune response in critically ill - SciSpace, <https://scispace.com/pdf/lactoferrin-effects-on-immune-response-in-critically-ill-tq30zs77fm.pdf>
66. Intermolecular Interactions in Plant Based vs. Animal Milks, <https://research-management.mq.edu.au/ws/portalfiles/portal/445740459/445099664.pdf>
67. Characterization of protein components of natural and heat-treated, https://www.researchgate.net/publication/223206667_Characterization_of_protein_components_of_natural_and_heat-treated_milk_fat_globule_membranes

68. xanthine dehydrogenase knock: Topics by Science.gov,
<https://www.science.gov/topicpages/x/xanthine+dehydrogenase+knock>
69. (A) SDS-PAGE profile of milk fat globule membrane (MFGM) isolated,
https://www.researchgate.net/figure/A-SDS-PAGE-profile-of-milk-fat-globule-membrane-MFGM-isolated-from-different-sized_fig4_376505671
70. Milk Fat Globule Membrane (MFGM) - Encyclopedia.pub,
<https://encyclopedia.pub/entry/2185>
71. Molecular Remodeling of Milk Fat Globules Induced by Centrifugation,
https://www.researchgate.net/publication/398543845_Molecular_Remodeling_of_Milk_Fat_Globules_Induced_by_Centrifugation_Insights_from_Deep_Learning-Based_Detection_of_Milk_Adulteration
72. cream separation.pptx,
<https://pt.slideshare.net/slideshow/cream-separationpptx/254602925>
73. Dairy product - Separation, Processing, Nutrition | Britannica,
<https://www.britannica.com/topic/dairy-product/Separation>
74. Increasing the Retention of Lipid-Soluble Components in a Curd,
<https://digitalcommons.usu.edu/cgi/viewcontent.cgi?referer=&httpsredir=1&article=2166&context=etd>
75. Changes in milk fat globules and membrane lipids under the shear,
https://www.academia.edu/90129638/Changes_in_milk_fat_globules_and_membrane_lipids_under_the_shear_fields_of_microfiltration_and_centrifugation
76. Emulsion-based delivery systems as a strategy to improve,
<https://www.tdx.cat/bitstream/10803/669633/5/Tagf1de1.pdf>
77. Cream Separation Methods Explained | PDF | Cream | Milk - Scribd,
<https://www.scribd.com/document/581096867/Cream-Part-I>
78. ILRI Training Manual 1: Rural Dairy Technology - CGSpace,
<https://cgspace.cgiar.org/bitstreams/2004e43d-67bd-44ad-96b1-a5d2f437eead/download>
79. Skimmer, Dairy ; Unknown manufacturer; 1920-1940; WY.1983.34,
<https://ehive.com/collections/3102/objects/1024138/skimmer-dairy>
80. Fat globule size distribution of 2% milk prepared by gravity,
https://www.researchgate.net/figure/Fat-globule-size-distribution-of-2-milk-prepared-by-gravity-separation-at-4-C-for-24-h_fig8_12338442
81. Clotted Cream - British Food: A History,
<https://britishfoodhistory.com/2019/10/11/clotted-cream/>
82. The Cream of England: What is Clotted Cream? - Fine Dining Lovers,
<https://www.finedininglovers.com/explore/articles/cream-england-what-clotted-cream>
83. Clotted cream - Wikipedia, https://en.wikipedia.org/wiki/Clotted_cream
84. Clotted Cream: A Cream-Tea Essential | TeaTime Magazine,
<https://teatimemagazine.com/clotted-cream-a-cream-tea-essential/>
85. Mock Devonshire Cream Recipe - What's Cooking America,
<https://whatscookingamerica.net/devonshirecream.htm>
86. fat globule size distribution and fat content of milk fractions - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/10984147/>

87. The composition of Devonshire cream - SciSpace,
<https://scispace.com/pdf/the-composition-of-devonshire-cream-2re373ldoo.pdf>
88. How does a milk separator work? - Milky Day Blog,
<https://milkyday.com/blog/2018/09/27/milk-separator-how-it-works/>
89. dairy science abstracts - CABI Digital Library,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20063034255>
90. Factors that Impact the Efficiency of Cream Separator Machine for,
<https://scholar.its.ac.id/en/publications/factors-that-impact-the-efficiency-of-cream-separator-machine-for/>
91. Separation of Milk - Dairy Technology,
<http://dairy-technology.blogspot.com/2014/01/separation-of-milk.html>
92. Milk fat globule size distribution of the bottom fraction, F1, after,
https://www.researchgate.net/figure/Milk-fat-globule-size-distribution-of-the-bottom-fraction-F1-after-gravity-separation_fig3_12338442
93. Comparing composition and structure of cream and skimmed milk,
<https://researchprofiles.ku.dk/da/publications/comparing-composition-and-structure-of-cream-and-skimmed-milk-profile>
94. Gravity Separation of Raw Bovine Milk: Fat Globule Size Distribution,
https://www.researchgate.net/publication/12338442_Gravity_Separation_of_Raw_Bovine_Milk_Fat_Globule_Size_Distribution_and_Fat_Content_of_Milk_Fractions
95. (PDF) Cream: Types of cream - ResearchGate,
https://www.researchgate.net/publication/294787584_Cream_Types_of_cream
96. Milk Reception and Storage in Dairies | PDF | Filtration | Pump - Scribd,
<https://www.scribd.com/document/690221045/6-Common-dairy-operations>
97. Cream Separation Techniques Explained | PDF | Centrifuge - Scribd,
<https://www.scribd.com/presentation/514806756/3-CREAM-SEPARATION-UNIT-II-23-3-2020>
98. (PDF) Effects of the Cooling Temperature at the Farm on Milk,
https://www.researchgate.net/publication/354913787_Effects_of_the_Cooling_Temperature_at_the_Farm_on_Milk_Maturation_and_Cheesemaking_Process_in_the_Manufacture_of_Parmigiano_Reggiano_PDO_Cheese
99. The Gravity Separation Of Fat, Somatic Cells, Bacteria, And Spores,
<https://ecommons.cornell.edu/entities/publication/46ae4e8d-465e-4836-b336-82825297cc0a>
100. Milk Separator: How to Calculate Efficiency and Fat Losses - Viravix,
<https://viravix.com/blog/milk-separator-efficiency-fat-losses/>
101. Impact of physico-chemical and operational factors on the process of,
https://www.researchgate.net/publication/367670925_Impact_of_physico-chemical_and_operational_factors_on_the_process_of_milk_skimming
102. Comparative lipidomics analysis of different-sized fat globules in,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10776417/>
103. (PDF) Unveiling the significance of psychrotrophic bacteria in milk,
https://www.researchgate.net/publication/377348737_Unveiling_the_significance_of_psychrotrophic_bacteria_in_milk_and_milk_product_spoilage_-_A_review
104. Recent Development in Detection and Control of Psychrotrophic,

- <https://www.mdpi.com/2304-8158/13/18/2908>
105. Gravity separation of fat, somatic cells, and bacteria in raw ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/23415516/>
 106. Influence of Somatic Cell Removal on Milk Quality and Yield - MDPI,
<https://www.mdpi.com/2306-5710/10/1/5>
 107. Spoilage potential of psychrotrophic bacteria isolated from raw milk,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6102184/>
 108. Microbiology of the Dairy Animal.pdf,
<https://s90289ad4ac6e0ddc.jimcontent.com/download/version/1395308938/module/9342846783/name/Microbiology%20of%20the%20Dairy%20Animal.pdf>
 109. Psychrotrophic bacteria and milk and dairy products quality,
<https://hrcak.srce.hr/file/124020>
 110. Psychrotrophic Bacteria Equipped with Virulence and Colonization,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10467336/>
 111. Chapter 1. Introduction, <https://eopcw.com/find/downloadLectureNote/1159>
 112. Faculty of Agriculture Ph.D. in Food Science and Biotechnology XXII,
https://amsdottorato.unibo.it/id/eprint/2865/1/Franciosi_Elena_Tesi.pdf
 113. (PDF) The main spoilage-related psychrotrophic bacteria included in,
https://www.researchgate.net/publication/395661649_The_main_spoilage-related_psychrotrophic_bacteria_included_in_the_industrial_slicing_of_mozzarella_cheese_under_sanitation_standard_operating_procedures
 114. Psychrotrophic bacteria in milk and their negative effects on dairy,
<https://dspace.univ-mila.dz/jspui/bitstream/123456789/3889/1/Psychrotrophic%20bacteria%20in%20milk%20and%20their%20negative%20effects%20on%20dairy%20products%20quality%20A%20review.pdf>