

The Biochemical and Physical Chemistry of Cultured Butter and Ghee Production: Phase Inversion, Clarification, and Oxidative Stability

The transformation of raw bovine milk into cultured butter, and its subsequent thermal clarification into anhydrous milk fat (ghee), represents one of the most intricate series of biophysical and biochemical operations in dairy science. Bovine milk is fundamentally a polyphasic biological fluid, existing simultaneously as a colloidal dispersion of casein micelles and an oil-in-water emulsion in which liquid triglyceride droplets are suspended within an aqueous plasma matrix¹. The thermodynamic instability of this emulsion is the foundational driver of phase separation, but the conversion of this emulsion into a water-in-oil matrix (butter) and eventually into a highly stable, 99.5% pure lipid phase (ghee) requires precisely controlled mechanical shear, microbial fermentation, and extreme thermal kinetics².

This report provides an exhaustive physical, biochemical, and operational analysis of this transformation. It delineates the microbial metabolism driving cream ripening, the rheological phenomena governing macroscopic phase inversion, the protective denaturation kinetics of whey proteins during clarification, the complex Maillard reaction cascade that generates the organoleptic profile of ghee, and the thermodynamic kinetics of lipid and cholesterol auto-oxidation during extended storage.

Emulsion Stability and the Milk Fat Globule Membrane (MFGM)

To comprehend the mechanics of churning, one must first rigorously analyze the structural barrier that prevents spontaneous phase inversion in raw milk: the Milk Fat Globule Membrane (MFGM). The MFGM is a highly complex, 5 to 50-nanometer-thick trilayered structure that coats every successfully secreted milk fat globule, physically isolating the hydrophobic triglyceride core from the hydrophilic aqueous plasma¹.

During intracellular biogenesis, lipid droplets originate within the endoplasmic reticulum. As they bud off, they are coated in an inner monolayer of specific polar lipids and intracellular proteins¹. These droplets subsequently migrate to the apical pole of the mammary epithelial cell and are pushed through the apical plasma membrane, enveloping the droplet in a true, outer cellular bilayer¹. This trilayered architecture creates a robust physical barrier possessing a highly negative zeta potential, which induces electrostatic repulsion and prevents the spontaneous coalescence of the droplets while they remain suspended in the milk plasma².

Embedded deeply within this intricate phospholipid matrix is a vast array of highly conserved transmembrane glycoproteins, peripheral proteins, and numerous active enzymes. Advanced proteomic analyses reveal the presence of at least 28 different enzymes within the MFGM,

including 5'-nucleotidase, gamma-glutamyl transpeptidase, plasmin, and aldolase, which collectively dictate the biological and physical resilience of the globule⁵.

MFGM Protein	Molecular Weight (kDa)	Biochemical and Structural Function
Xanthine Oxidase (XO)	155 kDa	Structural scaffolding; generates antimicrobial reactive oxygen species. Complexation with BTN maintains membrane rigidity.
Butyrophilin (BTN)	67 kDa	Transmembrane glycoprotein essential for lipid droplet secretion. Forms a robust structural lattice via disulfide bonds with XO.
Lactadherin (PAS 6/7)	~47–53 kDa	Peripheral glycoprotein providing antiviral activity and facilitating apoptotic cell clearance.
Adipophilin	50 kDa	Associated primarily with the inner monolayer; critical for intracellular lipid droplet assembly and structural stability.

Table 1: Primary protein constituents of the intact bovine Milk Fat Globule Membrane (MFGM) and their respective functions¹.
The physical integrity of this membrane is the primary obstacle to butter production. The mechanical disruption of the Butyrophilin-Xanthine Oxidase structural lattice, alongside the reduction of the membrane's electrostatic repulsion, is the fundamental biochemical prerequisite for phase inversion¹.

Microbial Fermentation and the Biochemistry of Cream Ripening

While sweet cream can be mechanically churned directly into butter, the production of traditional, highly prized cultured butter requires a carefully calibrated microbial fermentation phase. This ripening process serves a critical dual purpose: it lowers the pH of the plasma to biochemically alter the electrostatic stability of the MFGM, and it generates the vital volatile carbonyl compounds responsible for the characteristic aroma of cultured butter and downstream ghee¹¹.

Lactic Acid Glycolysis and Zeta Potential Modification

Fresh, pasteurized cream is inoculated with a mesophilic lactic starter culture, typically a precisely controlled biological consortium of *Lactococcus lactis* subsp. *cremoris*, *Lactococcus lactis* subsp. *lactis* biovar. *diacetylactis*, and *Leuconostoc mesenteroides*¹¹. As these obligate fermenters proliferate at an optimized incubation temperature of approximately 20°C, they metabolize available lactose into lactic acid via the homofermentative glycolytic pathway¹². The continuous accumulation of lactic acid rapidly depresses the pH of the cream from a native physiological baseline of 6.7 down to an acidic optimum of 4.4 to 5.0¹¹. This acidification profoundly impacts the physical chemistry of the emulsion. As the pH approaches the isoelectric point of the integral MFGM glycoproteins, the highly protective negative zeta potential of the fat globules is effectively neutralized⁷. The electrostatic repulsion that previously maintained the dispersion collapses, allowing the fat globules to flocculate and aggregate. This renders them vastly more susceptible to the mechanical shear forces that will be applied during subsequent churning¹⁷.

Citrate Metabolism and Diacetyl Synthesis Kinetics

The signature organoleptic profile of cultured butter—and by extension, the complex flavor profile of the resulting ghee—is driven heavily by the microbial synthesis of diacetyl (2,3-butanedione)¹¹. Crucially, diacetyl is not a direct byproduct of lactose fermentation; rather, it is synthesized via the citrate metabolism pathway uniquely possessed by specific strains, such as *Lactococcus lactis* biovar. *diacetylactis*¹⁵.

Bovine milk naturally contains a small concentration of citric acid. Citrate-positive bacterial strains utilize a specialized, plasmid-encoded citrate permease to transport citrate from the plasma into the intracellular matrix, where it is enzymatically cleaved into acetate and oxaloacetate¹⁵. Through subsequent decarboxylation, oxaloacetate is converted to pyruvate,

which is further metabolized into α -acetolactate. The spontaneous, non-enzymatic oxidative

decarboxylation of this α -acetolactate yields the highly aromatic compound diacetyl¹⁵.

The kinetic efficiency of this specific metabolic pathway is exquisitely sensitive to the thermal and chemical environment of the fermentation vat. Advanced research indicates that the absolute maximum diacetyl yield (achieving concentrations up to 105 mg/L) is realized at a cultivation temperature of exactly 20°C, particularly when the substrate is supplemented with

trace metal catalysts¹⁵. Specifically, the addition of Fe^{2+} at 0.001% significantly upregulates

α -acetolactate decarboxylase activity, forcefully driving the reaction toward diacetyl

accumulation¹⁵. Conversely, if the ripening temperature is inadvertently elevated to 30°C, the activity of diacetyl reductase increases exponentially, rapidly reducing the highly desirable diacetyl into a relatively flavorless compound known as acetoin, thereby permanently impairing the sensory quality and commercial value of the butter¹⁵.

Thermal Tempering and Triglyceride Crystallization

Concurrent with the microbial fermentation phase, the cream must undergo precise thermal tempering to induce partial fat crystallization. Bovine milk fat is an exceptionally complex and heterogeneous lipid matrix, comprising well over 3,000 distinct triacylglycerol (TAG) species²⁰. Because of this immense molecular diversity, milk fat does not possess a single, discrete melting point; rather, it melts over a broad thermodynamic continuum ranging from -40°C to +40°C²¹.

Solid Fat Content (SFC) Optimization

For the mechanical process of churning to succeed, the liquid cream must be engineered to possess a highly specific Solid Fat Content (SFC)—typically stabilized between 30% and 40% prior to entering the churn¹¹. If the SFC is too low, the fat remains excessively liquid and fails to retain the air necessary for foam formation, resulting in massive fat losses into the buttermilk¹¹. Conversely, if the SFC is too high due to over-chilling, the fat globules become rigid, brittle spheres that refuse to coalesce under mechanical shear, yielding a crumbly, unworkable product¹¹.

Crystal Polymorphism and Sub-cell Packing

Milk fat exhibits profound crystal polymorphism, meaning the constituent triglycerides can pack into several entirely different crystalline lattices depending strictly on the thermal history and specific cooling rate applied to the fluid²². Detailed analyses utilizing Differential Scanning Calorimetry (DSC) and Real-Time X-Ray Powder Diffraction (XRD) have confirmed that the three primary polymorphic forms of milk fat, classified in order of increasing thermodynamic stability and crystalline density, are the α , β' , and β modifications²².

Polymorphic Modification	Sub-cell Lattice Packing	Thermodynamic Stability	Primary Triglyceride Drivers and Formation Conditions
α (Alpha)	Hexagonal	Lowest (Metastable)	Formed during rapid quenching or extreme cooling rates. Driven by high concentrations of saturated TAGs with carbon

			numbers (CN) 34-38.
β' (Beta-prime)	Orthorhombic	Intermediate	Formed during controlled, gradual tempering. Essential for optimal butter texture, spreadability, and structural puncturing of the MFGM.
β (Beta)	Triclinic	Highest (Most Stable)	Formed during prolonged storage or slow maturation. Favored by unsaturated TAGs with carbon numbers (CN) 52-54.

Table 2: Polymorphic crystal modifications of bovine milk fat and their corresponding thermodynamic properties²⁰.

During the precise tempering protocol, the pasteurized, inoculated cream is rapidly cooled to approximately 8°C. This rapid thermal drop forces the spontaneous nucleation of the unstable α -crystals²⁴. Following this initial shock, the temperature is gently raised to roughly 20°C to melt the lowest-melting fractions, and then slowly cooled again to 15°C²⁷. This specific temperature cycling deliberately forces a polymorphic transition, driving the fat from the metastable α -form into the highly desired, intermediate β' -form²³.

The resulting β' crystals form a delicate, needle-like internal network that protrudes outward toward the periphery of the fat globule. During the subsequent churning process, these sharp, dense β' crystals physically puncture the MFGM from the inside out. This internal structural failure provides a critical mechanical catalyst for membrane rupture, allowing the entrapped, sticky liquid fat to exude from the globule and cement adjacent droplets together¹¹.

The Fluid Dynamics and Mechanics of Phase

Inversion

The physical transformation of fluid cream into solid butter represents the macroscopic manifestation of extreme microscopic emulsion destabilization. Contemporary physical chemistry relies on a multidisciplinary synthesis of several historical models to explain the exact mechanics of this phase inversion, moving past overly simplistic models to embrace complex fluid dynamics¹⁷.

Historical and Contemporary Churning Theories

Early scientific attempts to explain the churning process, such as the theory proposed by Fisher and Hooker, posited a simple, direct reversal. They argued that cream is naturally an oil-in-water (O/W) emulsion and that the mechanical agitation of churning simply reverses this arrangement entirely, converting it into a water-in-oil (W/O) emulsion¹⁸. However, advanced microscopic studies demonstrated that finished butter still contains millions of fully intact fat globules suspended within the free fat matrix, proving that the phase reversal is not absolute, rendering the Fisher and Hooker model incomplete¹⁸.

Subsequently, Rahn proposed the Foam Theory, which argued that agitation creates a massive foam structure by incorporating air into the fluid. Due to intense surface tension dynamics, the discrete milk fat globules rapidly migrate to the air-water interfaces of these newly formed air bubbles, concentrating the fat and forcing the globules into proximity¹⁸.

Contemporary dairy science relies on King's Modern Theory, which builds upon Rahn's foundation by incorporating the mechanical destruction of the MFGM. According to King, as the agitation persists, the fat globules concentrated in the foam are forced into violent, high-velocity collisions¹⁷.

Churning Theory	Primary Proponents	Core Mechanistic Argument	Limitations in Modern Context
Phase Reversal	Fisher & Hooker	Agitation directly forces an absolute flip from an O/W emulsion to a W/O emulsion.	Fails to account for the millions of intact fat globules proven to remain suspended in finished butter.
Foam Theory	Rahn	Air incorporation creates a foam; surface tension draws fat globules to the air-plasma interface where they crowd and merge.	Incomplete; does not fully explain the precise mechanism of membrane rupture required for coalescence.

Modern Synthesis	King	Foam concentrates globules, but intense mechanical shear and friction actively tear the MFGM. Liquid fat leaks out, acts as a foam depressant, and cements the mass together.	Currently accepted as the most accurate biophysical model of phase inversion.
------------------	------	---	---

Table 3: The evolution of biophysical theories explaining the mechanics of cream churning¹⁷.

Macroscopic Coalescence and Foam Collapse

The hydrodynamic shear stress generated by the churn's dashers, combined with the extreme pressure exerted at the air-plasma interface and the internal puncturing action of the β' fat crystals, subjects the MFGM to catastrophic physical strain². The delicate intermolecular disulfide bonds holding the Butyrophilin and Xanthine Oxidase complex together are mechanically severed, and the phospholipid bilayer is physically ripped apart, stripping the protective membrane away from the hydrophobic core¹.

Upon the complete rupture of the MFGM, the liquid triglyceride fraction—which was carefully preserved through meticulous temperature control during the tempering phase—leaks out of the globule. This highly viscous, sticky liquid fat spreads as a thin film over the surface of the air bubbles¹⁸. In fluid dynamics, this free lipid acts as a profound chemical foam depressant¹⁸. As the free fat rapidly extinguishes the surface tension supporting the foam, the air bubbles burst, and the foam structure undergoes a sudden, violent, and macroscopic collapse. The localized fat clusters, now completely lacking electrostatic repulsion and heavily coated in adhesive liquid fat, instantly coalesce into massive, macroscopic agglomerations known universally as butter grains².

This precise, visually distinct moment represents the phase inversion. The system transitions instantaneously from an oil-in-water emulsion (fat suspended in plasma) to a true water-in-oil emulsion (aqueous droplets permanently trapped within a continuous matrix of solid and liquid fat)². The expelled aqueous phase, containing the sheared remnants of the MFGM (including high concentrations of polar lipids and glycoproteins), unfermented lactose, and whey proteins, drains away as buttermilk⁵.

Buttermilk Washing and Hydrolytic Rancidity

Mitigation

Following phase inversion, the residual buttermilk entrapped within the newly formed butter grains must be systematically expelled through a careful process of washing and mechanical

working. Chilled break water—strictly calculated to generally not exceed 25% of the total cream volume—is introduced into the churn to rapidly harden the butter grains and flush out the residual aqueous plasma¹⁸.

The biochemical imperative for thorough washing is the absolute removal of residual lactose and non-fat proteins. The continued presence of these hydrophilic, nutrient-dense compounds within the final butter matrix creates a highly hospitable microenvironment for aggressive bacterial spoilage³⁰.

The Threat of Psychrotrophic Lipolysis

The most critical and devastating microbiological threat to cultured butter is the proliferation of *Pseudomonas fluorescens* and allied gram-negative psychrotrophs (such as *Alcaligenes* and *Flavobacterium* species)³¹. While the vegetative cells of these specific bacteria are typically destroyed by initial pasteurization, post-processing contamination or the prolonged survival of their metabolic byproducts poses a catastrophic risk to lipid stability during storage³⁰.

Pseudomonas species that have adapted to cold environments possess a dangerous evolutionary advantage: they synthesize massive quantities of extracellular lipases and proteases at refrigeration temperatures³¹. Crucially, the specific lipases produced by *P. fluorescens* are exceptionally heat-stable³⁰. These extracellular enzymes easily survive standard pasteurization and even ultra-high-temperature (UHT) protocols that obliterate the bacteria that produced them³⁰.

If these resilient lipases are allowed to persist in the aqueous pockets of the butter, they will aggressively hydrolyze the ester bonds of the triglycerides at the oil-water interface, preferentially cleaving short-chain fatty acids from the glycerol backbone³³. The resulting liberation of free butyric, caproic, and caprylic acids imparts severe hydrolytic rancidity to the butter, characterized by intensely soapy, bitter, and vomit-like organoleptic defects that render the fat entirely unfit for human consumption or downstream conversion into ghee⁸.

Consequently, meticulous washing and aggressive mechanical working of the butter is required to minimize the total volume of the aqueous phase and distribute it as microscopic droplets

(typically less than 5 μm in diameter). This microscopic compartmentalization physically restricts bacterial mobility and severely limits enzymatic access to the triglyceride interface, extending the shelf life of the butter²⁷.

Thermal Clarification: The Conversion of Butter into Ghee

While cultured butter is a stabilized emulsion, it remains inherently susceptible to microbiological spoilage and enzymatic degradation due to its ~16% moisture content. The conversion of butter into ghee (anhydrous milk fat) via thermal clarification removes this moisture entirely, halting all microbial activity and initiating a cascade of complex, heat-driven biochemical reactions that radically extend the shelf life and alter the flavor profile of the lipid³.

Thermodynamics of Moisture Evaporation and Protein Coagulation

The clarification process requires heating the cultured butter in a specialized, steam-jacketed

boiler to temperatures reaching between 110°C and 120°C³⁷. As the thermal energy of the system increases, the entrapped water reaches its boiling point, generating intense physical effervescence and foaming at the surface. Simultaneously, the residual solids-not-fat (SNF)—comprising primarily casein proteins, native whey proteins, and unfermented lactose—begin to chemically coagulate under the extreme heat³⁹.

As the temperature rapidly breaches 100°C, the water is forcefully driven off as steam. The violent boiling of the dispersed aqueous phase forces the coagulated SNF particles to agglomerate, eventually settling at the bottom of the vessel as a dense, dark mass known universally in the industry as ghee residue³⁷. The precise endpoint of clarification is dictated visually and audibly by the complete cessation of effervescence and the profound browning of the curd particles, signalling that the moisture content of the lipid has safely fallen below 0.3%³⁹.

Whey Protein Denaturation and Sulfhydryl Release

A critical, yet historically underappreciated, biochemical phenomenon occurring during

clarification is the severe thermal denaturation of residual whey proteins, most notably β

-lactoglobulin⁴¹. β -lactoglobulin is a globular protein containing buried cysteine residues that possess highly reactive free sulfhydryl (-SH) groups⁴³.

As the clarification temperature exceeds 60°C, the tertiary structure of the β -lactoglobulin molecule begins to unfold⁴². By the time the overall temperature of the fat reaches 110°C, the protein undergoes massive, irreversible thermal denaturation. This unfolding physically exposes the previously shielded sulfhydryl groups directly to the surrounding hot lipid matrix⁴².

The liberation of these sulfhydryl compounds is profoundly beneficial to the chemical longevity of the ghee. These exposed -SH groups function as potent, natural, sacrificial antioxidants. By preferentially reacting with dissolved molecular oxygen and rapidly sequestering free radicals

within the fluid, the sulfhydryl compounds derived from the denatured β -lactoglobulin drastically retard the initiation phase of lipid auto-oxidation, granting high-heat clarified ghee its legendary, extended oxidative stability⁴¹.

The Maillard Reaction Cascade in Clarified Butter

The signature nutty, caramelized, and intensely savory flavor profile of traditional ghee is not derived from the pure lipids themselves, but rather from the severe thermal degradation of the trace carbohydrates and proteins trapped in the sinking ghee residue. This entire organoleptic transformation is governed strictly by the Maillard reaction, a highly complex, multi-stage non-enzymatic browning cascade⁴⁴.

The Early and Intermediate Stages: Amadori Rearrangement and Enolization

The Maillard reaction initiates rapidly when the electrophilic carbonyl group of a reducing sugar (typically residual lactose, or its cleaved monomers glucose and galactose) undergoes

nucleophilic attack by the free ϵ -amino group of a protein-bound amino acid, most notably

lysine⁴⁷. This condensation reaction immediately forms an unstable Schiff base, which quickly undergoes an Amadori rearrangement to form a stable ketoamine⁵⁰. In dairy systems subjected to heat, the primary Amadori product generated is lactulosyl-lysine (or fructosyl-lysine)⁴⁸. Because the Amadori compounds themselves are entirely colorless and flavorless, they must degrade further to generate the desirable organoleptic properties of ghee⁴⁶. The specific degradation pathway these compounds follow is dictated entirely by the pH of the surrounding microenvironment⁵⁰.

Because the butter was cultured and heavily acidified by *Lactococcus lactis* prior to churning, the pH during clarification is relatively low (acidic). Under acidic conditions, the Amadori products are forced to undergo **1,2-enolization**⁵⁰. This specific dehydration pathway rapidly yields 3-deoxyosones, which are subsequently converted into highly reactive intermediates such as 5-hydroxymethylfurfural (HMF) and furfural⁵⁰. (Conversely, had sweet, uncultured cream been utilized, the neutral pH would have heavily favored 2,3-enolization, leading to a vastly different profile of 1-deoxyosones and reductones, ultimately resulting in an inferior flavor complexity⁵¹).

Maillard Stage	Chemical Mechanism	Primary Intermediate/Analytical Marker	Visual and Organoleptic Impact
Early	Condensation & Amadori Rearrangement	Lactulosyl-lysine; Furosine (Quantified via HPLC)	Odorless, colorless. Reduces the nutritional bioavailability of lysine.
Intermediate	1,2-Enolization (Dictated by the acidic pH of cultured butter)	3-Deoxyglucosone; 5-Hydroxymethylfurfural (HMF)	Precursors to aroma formation; slight yellowing of the residue.
Advanced	Strecker Degradation, Aldol Condensation, Polymerization	Carboxymethyl-lysine (CML); Melanoidins; Pyrazines	Intense browning; generation of roasted, nutty, and highly savory aromas.

Table 4: The sequential Maillard reaction cascade occurring during the thermal clarification of cultured butter into ghee⁴⁶.

The Advanced Stage: Melanoidins, AGEs, and Flavor Chemistry

As clarification continues and the temperature holds at 120°C, the reaction enters the advanced stage. Here, the intermediate dicarbonyls undergo Strecker degradation by reacting with free amino acids, releasing volatile Strecker aldehydes and generating a massive, highly aromatic array of heterocyclic volatile compounds⁴⁷.

Gas Chromatography-Mass Spectrometry (GC-MS) profiling of high-quality ghee reveals an incredibly dense concentration of pyrazines (yielding roasted, nutty notes), furans (yielding sweet, caramel notes), and various highly specific lactones⁵⁴. Specifically, the synthesis of δ

-decalactone and γ -dodecalactone during this phase imparts the heavy, fatty, and characteristic peach-like sweet notes that define premium artisanal ghee⁵⁸.

Simultaneously, the chemical reaction terminates in the widespread polymerization of melanoidins—large, nitrogen-rich brown polymers that settle heavily into the ghee residue and impart the rich golden hue to the liquid fat above⁴⁷. Advanced Glycation End-products (AGEs),

such as N- ϵ -(carboxymethyl)lysine (CML) and N- ϵ -(carboxyethyl)lysine (CEL), are also formed via the oxidative cleavage of the early Amadori products⁵¹.

The exact extent of this thermal damage is closely monitored in modern dairy science. The early-stage marker, lactulosyl-lysine, is indirectly quantified by measuring its acid-hydrolysis product, **Furosine**, using reversed-phase High-Performance Liquid Chromatography (HPLC) coupled with fluorescence detection⁴⁸. Concurrently, CML serves as an excellent analytical marker for advanced thermal processing severity⁴⁸. While a controlled Maillard reaction is essential for flavor, extreme accumulation of AGEs resulting from excessive temperature abuse (>150°C) can severely diminish the nutritional quality of the fat and block lysine bioavailability⁴⁸.

Kinetics of Lipid Auto-Oxidation and Cholesterol

Degradation

Despite the complete expulsion of moisture and the generation of highly protective sulfhydryl groups and antioxidant melanoidins during clarification, ghee remains a lipid and is therefore susceptible to long-term chemical degradation during ambient storage. The primary mechanisms of shelf-life failure in anhydrous milk fat are lipid auto-oxidation and cholesterol oxidation³.

Arrhenius Kinetics of Lipid Auto-Oxidation

The thermodynamic degradation of the polyunsaturated and monounsaturated fatty acids within the ghee matrix follows a classical free-radical chain reaction, comprising distinct initiation, propagation, and termination phases⁶⁵.

This auto-oxidation process is mathematically modeled using Arrhenius reaction kinetics, where

the reaction rate constant (k) scales exponentially with the absolute temperature of the environment. During the highly destructive propagation phase, lipid radicals ($R^\bullet$) react rapidly with dissolved molecular oxygen to form peroxy radicals ($ROO^\bullet$). These peroxy radicals

then aggressively abstract hydrogen from adjacent, intact lipids to form hydroperoxides (

$ROOH$)⁶⁵. Hydroperoxides are the primary, yet chemically unstable, products of oxidation, and their accumulation is typically quantified via the Peroxide Value (PV) iodometric assay⁸. As the oxidation reaction progresses over months of storage, these hydroperoxides undergo homolytic cleavage to generate secondary, highly volatile oxidation products. The most notable of these is hexanal, a short-chain aldehyde which imparts the characteristic rancid, cardboard-like off-flavors to degraded ghee⁶⁵. Advanced studies characterizing the

thermodynamic parameters of milk fat oxidation report activation energies (E_a) ranging widely from 80 to 154 kJ/mol⁶⁷. The exact activation energy depends heavily on the ratio of saturated to unsaturated fatty acids in the specific batch of milk, as well as the initial concentration of inherent antioxidants (such as the denatured β -lactoglobulin sulfhydryls and naturally occurring α -tocopherol)⁴⁵. To artificially elevate this activation energy and extend the Oxidative Stability Index (OSI), contemporary research has successfully explored the addition of natural polyphenol-rich herb extracts, such as orange peel powder or *Aegle marmelos*, which forcefully scavenge the DPPH radicals before they can propagate the chain reaction⁹.

Cholesterol Oxidation Products (COPs)

Bovine milk fat naturally contains significant quantities of cholesterol, an unsaturated sterol integral to animal cell membranes. When ghee is subjected to extreme thermal stress (such as during deep-fat frying) or prolonged aerobic storage, the cholesterol fraction undergoes severe oxidative degradation, yielding highly toxic Cholesterol Oxidation Products (COPs)⁷¹.

The auto-oxidation of cholesterol initiates primarily at the C-7 position via targeted hydrogen abstraction, followed immediately by oxygen addition to form 7 α -hydroperoxycholesterol and 7 β -hydroperoxycholesterol⁷². As heating continues, these volatile hydroperoxides readily undergo dehydration to form **7-ketocholesterol**, which serves as the predominant and most heavily monitored COP marker in thermally abused dairy fats⁷².

Crucially, the kinetics of COP formation are heavily temperature-dependent. Clarifying ghee at the traditional optimum of 110°C to 120°C generally minimizes cholesterol oxidation⁷³. However, elevating the clarification or frying temperature beyond the critical threshold of 150°C triggers a

massive, exponential surge in the formation of 7-ketocholesterol, α -epoxides, and 25-hydroxycholesterol⁷⁵. Because COPs are heavily implicated in advanced atherogenesis, cytotoxicity, and neurodegenerative pathologies, precise thermal management during the final boiling stage of ghee is an absolute necessity to ensure both organoleptic quality and human food safety⁷².

Industrial vs. Artisanal Methodologies: Process Engineering

The biochemical and biophysical principles outlined above are universally applicable to the transformation of milk into ghee. However, the specific engineering methodologies utilized to

extract the fat and apply the heat profoundly alter the thermodynamic efficiency, yield, and final chemical profile of the product. The global dairy industry recognizes several distinct processing methodologies, ranging from ancient artisanal techniques to modern, highly optimized continuous fluid dynamics systems⁴⁰.

The Indigenous (Bilona/Desi) Method

The traditional, artisanal method involves the fermentation of whole milk (or raw, clotted cream) into a solid curd (*dahi*), which is subsequently hand-churned in a wooden vessel (*bilona*)⁴⁰. The resulting butter (*makkhan*) is then heated over an open, smoky fire.

While highly inefficient from an industrial standpoint—suffering from severe fat loss into the massive volume of buttermilk generated (recovering only 80-85% of available lipids)—the Bilona method is deeply revered for yielding a product with an unmatched sensory profile⁴⁰. The prolonged fermentation of the entire whole milk matrix maximizes diacetyl and short-chain fatty acid development, while the unique inclusion of massive amounts of lactic acid bacteria directly into the boiling vessel vastly enhances the complexity of the final Maillard reaction⁴⁰.

Furthermore, proponents correctly note that the gentle, low-shear traditional method fractionally preserves slightly higher concentrations of bioactive conjugated linoleic acid (CLA) and essential fatty acids compared to the violent hydrodynamic forces of centrifugal extraction⁸⁰.

Pre-Stratification and Creamery Butter Methods

To achieve massive industrial scale, modern manufacturers typically utilize the **Creamery Butter Method**. Here, cream is centrifugally separated from the milk, pasteurized, culturally ripened, mechanically churned in continuous high-speed Fritz-style churns, and then melted and pumped into massive steam-jacketed kettles for boiling¹⁷. While highly scalable and yielding a respectable 88-92% fat recovery, the extensive frothing during the boiling of the 16% moisture butter requires careful thermal management to prevent boil-over⁴⁰.

To drastically reduce fuel consumption and limit the exposure of the fat to prolonged heat, dairy engineers developed the **Pre-Stratification Method**. In this highly efficient protocol, the butter is melted at roughly 80°C and held completely undisturbed in a specialized settling tank⁷⁸. Over several hours, the density differentials cause the emulsion to physically stratify into three distinct layers: denatured proteins float to the top, pure liquid fat settles in the middle, and the heavy, acidic buttermilk serum sinks to the bottom⁸². By mechanically draining the bottom serum layer prior to boiling, up to 50% of the moisture is removed without the application of high heat. The remaining fat is then boiled briefly at 110°C to induce the required Maillard browning⁴⁰. This minimizes the formation of Free Fatty Acids (FFAs), significantly extending the oxidative shelf life of the resulting ghee while saving up to 45% in energy costs⁴⁰.

Continuous Ghee Manufacture and Scraped Surface Heat Exchangers (SSHE)

For ultra-high-volume industrial operations, batch boiling in open kettles has been superseded by fully continuous, closed-loop systems designed to maximize thermodynamic efficiency⁷⁷. In the modern continuous method, raw cream is subjected to extreme centrifugal force in a clarifixer or centrifixer, which mechanically obliterates the MFGM and breaks the emulsion

without a traditional churning step, concentrating the fat directly to 95-99%⁷⁷. The nearly pure lipid is then pumped directly into a **Scraped Surface Heat Exchanger (SSHE)**. The SSHE consists of a steam-jacketed cylinder housing a rapidly rotating central shaft fitted with razor-sharp scraper blades⁸³. As the highly viscous fat is pumped through the cylinder, the blades continuously wipe the heated inner wall⁸⁵.

Ghee Production Method	Fat Recovery Efficiency	Energy Profile & Thermodynamics	Unique Chemical/Sensory Implications
Indigenous (Bilona)	Lowest (80–85%)	Highly inefficient; requires prolonged boiling of massive volumes of water.	Superior flavor complexity due to heavy lactic acid bacteria inclusion in the Maillard reaction. High CLA retention.
Creamery Butter	High (88–92%)	Moderate energy use; scalable batch process.	Standardized flavor; highly consistent output for commercial retail.
Pre-Stratification	High (~92%)	High efficiency; saves 35-50% thermal energy by gravitationally draining serum prior to boiling.	Low FFA development yields exceptional oxidative shelf life. Flavor is notably milder.
Continuous (SSHE)	Highest (>95%)	Maximum efficiency; lowest specific exergy destruction.	Prevents burn-on entirely. Excellent preservation of heat-labile vitamins due to rapid residence time.

Table 5: Comparison of operational efficiencies and chemical outcomes across ghee production methodologies⁴⁰.

This continuous mechanical scraping disrupts the static thermal boundary layer that would

normally insulate the fluid, ensuring an exceptionally high and uniform overall heat transfer coefficient (achieving U -values up to $1600 \text{ W/m}^2 \text{ K}$)⁸⁵. The fat is flash-heated to 115°C in a matter of seconds, inducing the requisite Maillard browning and moisture evaporation instantly⁷⁷. Because the residence time at high heat is reduced from hours (in an open pan) to mere seconds, the SSHE completely eliminates burn-on, totally prevents the formation of toxic cholesterol oxidation products (COPs), and highly preserves the delicate, heat-labile fat-soluble vitamins (A, D, E, and K) while achieving an overall exergy efficiency of 26.8% ⁷⁷.

Conclusion

The production of cultured butter and shelf-stable ghee is a profound triumph of applied biophysics, fluid mechanics, and food chemistry. The process requires a multidisciplinary orchestration of microscopic variables: the precise biological manipulation of the *Lactococcus citrate* metabolism to generate diacetyl, the exact thermodynamic tempering of complex

triglyceride polymorphism to yield the critical β' crystals, and the violent hydrodynamic shear required to rupture the highly conserved Butyrophilin-Xanthine Oxidase architecture of the MFGM.

Once macroscopic phase inversion is achieved, meticulous washing is required to starve out psychrotrophic *Pseudomonas* lipases, ensuring the triglyceride ester bonds remain intact. Finally, the thermal clarification of the butter into ghee perfectly balances the destructive and protective forces of extreme heat. By carefully driving the temperature to 115°C , manufacturers deliberately induce the 1,2-enolization Maillard cascade to build complex pyrazine and lactone

flavor profiles, while simultaneously denaturing β -lactoglobulin to release highly protective, antioxidant sulfhydryl groups into the matrix. However, strict engineering controls—best realized through the highly efficient thermodynamics of modern Scraped Surface Heat Exchangers—must be maintained to prevent the reaction from proceeding into severe cholesterol oxidation. The resulting anhydrous milk fat stands as a highly stable, microbiologically secure, and biochemically complex lipid matrix, perfectly engineered for long-term ambient stability.

Works cited

1. 81 Gravity Cream Separation Physics.pdf
2. Traditional Butter Making: The Physics of Phase Inversion in Cream, <https://milkiry.pages.dev/traditional-butter-making-physics-phase-inversion/>
3. Food Technology (2023), 47(2), 124-139 ORIGINAL RESEARCH P, <https://www.gup.ugat.ro/ugatjournals/index.php/food/article/download/6436/5560>
4. Ghee Processing - NIFTEM, https://niftem.ac.in/newsite/pmfme/wp-content/uploads/2022/08/ppt_ghee-processing.pdf
5. Milk fat globule membrane and buttermilks: from composition to, <https://popups.uliege.be/1780-4507/index.php?id=5828>
6. Oxidative Stability of Ghee Incorporated with Clove Extracts and,

- <https://www.tandfonline.com/doi/full/10.1080/10942912.2012.752382>
7. Science Animation - MFGM: Milk Fat Globule Membrane - YouTube,
<https://www.youtube.com/watch?v=Iz9dIJ1Zbdo>
 8. Comparative Study on Keeping Quality of Market Ghee Samples,
<https://www.ijcmas.com/9-9-2020/Niranjan.%20et%20al.pdf>
 9. Retardation of oxidative rancidity in ghee adding orange peel ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9813280/>
 10. Research of milk fat oxidation processes during storage of butter,
<https://potravinarstvo.com/journal1/index.php/potravinarstvo/article/view/1283/1411>
 11. Butter Science 101 - Center for Dairy Research,
<https://cdr.wisc.edu/butter-science-101>
 12. Influence of temperature regimes of ripening and fermentation,
https://www.researchgate.net/publication/328786055_Influence_of_temperature_regimes_of_ripening_and_fermentation_stages_on_the_physical_and_chemical_properties_of_cream_and_sour-cream_butter_quality_indicators
 13. Physicochemical changes during cream fermentation and quality,
https://www.researchgate.net/publication/371654485_Physicochemical_changes_during_cream_fermentation_and_quality_characteristics_of_lactic_butter_produced_using_cultured_creams_with_different_fermentation_levels
 14. Lactococcus lactis in Dairy Fermentation—Health-Promoting ... - MDPI,
<https://www.mdpi.com/2311-5637/10/1/16>
 15. Isolation of Lactococcus lactis ssp. cremoris LRCC5306 and ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8112315/>
 16. Screening and selection of cultures based on techno-functional,
<https://epubs.icar.org.in/index.php/IJDS/article/view/81033>
 17. Buttermaking and the churning of blended fat emulsions,
https://www.researchgate.net/publication/228056869_Buttermaking_and_the_churning_of_blended_fat_emulsions
 18. Theories Behind the Churning Process in Butter Production,
<https://agriculture.institute/dairy-products-i/theories-churning-process-butter-production/>
 19. Effect of citrate on growth of Lactococcus lactis subsp. lactis in milk,
<https://pubmed.ncbi.nlm.nih.gov/9299783/>
 20. Milk fat crystallization: From molecules to networks - Research@WUR,
<https://research.wur.nl/en/publications/milk-fat-crystallization-from-molecules-to-networks/>
 21. Modelling the butterfat crystallisation process - ORBi,
<https://orbi.uliege.be/bitstream/2268/74784/1/butter.pdf>
 22. Polymorphism of milk fat studied by differential scanning calorimetry,
https://www.researchgate.net/publication/227325756_Polymorphism_of_milk_fat_studied_by_differential_scanning_calorimetry_and_real-time_X-ray_powder_diffraction
 23. POLYMORPHISM IN MILK FAT - CABI Digital Library,
<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20063034165>

24. Fat Crystallisation:,
<https://www.soci.org/-/media/files/conference-downloads/2007/sci-cairo-conference-mar-07/ralphtimms.ashx>
25. Polymorphism, microstructure and rheology of butter. Effects of,
https://nano.ku.dk/english/research/structural-food-physics-and-soft-matter-self-assembly/pdfs/ronholt_2012.pdf
26. Formation of β Polymorphs in Milk Fats with Large Differences in,
<https://pubmed.ncbi.nlm.nih.gov/27121766/>
27. Standardisation of cultured butter processing for smallscale production,
<https://mro.massey.ac.nz/bitstreams/cc475c31-94f1-4d7c-a5e7-b1223a4b34f3/download>
28. Chemistry: Inverting the original cream emulsion : Butter - Webexhibits,
<https://www.webexhibits.org/butter/churning.html>
29. Technology of Butter making – Batch and Continuous Processes Introd,
https://epgp.inflibnet.ac.in/epgpdata/uploads/epgp_content/S000015FT/P000357/M002914/ET/1455793585ET09.pdf
30. The lipolytic and proteolytic activity of bacteria isolated from blended,
https://www.researchgate.net/publication/230109593_The_lipolytic_and_proteolytic_activity_of_bacteria_isolated_from_blended_raw_milk
31. Dairy Spoilage and Pseudomonas spp. - Encyclopedia.pub,
<https://encyclopedia.pub/entry/20197>
32. Microbial lipolysis at low temperatures - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC291280/>
33. Microbial lipolysis at low temperatures - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/6766702/>
34. The Biodiversity of the Microbiota Producing Heat-Resistant,
<https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2017.00302/full>
35. Purification and Kinetics of Lipase of Pseudomonas fluorescens,
<https://scialert.net/fulltext/?doi=jbs.2021.29.37>
36. impacts on dairy product quality and biotechnological potential,
<https://rsdjournal.org/rsd/article/view/21213>
37. Recent innovations in functionality and shelf life enhancement of,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9813307/>
38. Flavour profile of desi cow's milk ghee prepared with bael pulp extract,
<https://epubs.icar.org.in/index.php/IJDS/article/download/170104/65868/525272>
39. Different Types of Ghee Making Process - Urbana Superfoods,
<https://urbanasuperfoods.com/blogs/nutrition/different-types-of-ghee-making-process>
40. Comparing Ghee Production Methods: Which One is Right for You?,
<https://agriculture.institute/dairy-products-i/comparing-ghee-production-methods/>
41. (PDF) Ghee Residue: Processing and Utilization in Food and Non,
https://www.researchgate.net/publication/344602736_Ghee_Residue_Processing_and_Utilization_in_Food_and_Non-Food_Application
42. Heat Induced Changes in Milk and Traditional Milk Products - AWS,

https://gph-prh.s3.ap-south-1.amazonaws.com/2024/Apr/20-Apr-24/2024_ARJOCS_1505/Ms_ARJOCS_1505.pdf

43. Quantification of heat-induced casein–whey protein interactions in, https://www.researchgate.net/publication/222048043_Quantification_of_heat-induced_casein-whey_protein_interactions_in_milk_and_its_relation_to_gelation_kinetics
44. Effect of Maillard reaction on the quality of clarified butter, ghee, https://www.researchgate.net/publication/365090534_Effect_of_Maillard_reaction_on_the_quality_of_clarified_butter_ghee
45. Effect of Maillard reaction on the quality of clarified butter, ghee, <https://pubmed.ncbi.nlm.nih.gov/36324007/>
46. Maillard Reaction: Mechanism, Influencing Parameters, Advantages, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12154226/>
47. Maillard reaction - Wikipedia, https://en.wikipedia.org/wiki/Maillard_reaction
48. (PDF) Blocked Lysine in Dairy Products: Formation, Occurrence, https://www.researchgate.net/publication/284136220_Blocked_Lysine_in_Dairy_Products_Formation_Occurrence_Analysis_and_Nutritional_Implications
49. Dietary glycation compounds – implications for human health? - DFG, <https://www.dfg.de/resource/blob/174038/191206-roundtable-conference-material.pdf>
50. Maillard reaction in food technologies, <https://nuft.edu.ua/doi/doc/ujfs/2023/2/4.pdf>
51. Antioxidant Activity of Maillard Reaction Products in Dairy Products, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12841591/>
52. Maillard Reaction in Flour Product Processing: Mechanism, Impact, <https://www.mdpi.com/2304-8158/14/15/2721>
53. Qualitative and quantitative analysis of furosine in fresh ... - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5766691/>
54. Science | Maillard Reaction - The Brewers Journal, <https://www.brewersjournal.info/science-maillard-reaction/>
55. The role of the Maillard reaction in the formation of flavour, https://www.researchgate.net/publication/230797196_The_role_of_the_Maillard_reaction_in_the_formation_of_flavour_compounds_in_dairy_products_-_Not_only_a_deleterious_reaction_but_also_a_rich_source_of_flavour_compounds
56. (PDF) How Maillard Reaction Influences Sensorial Properties (Color, https://www.researchgate.net/publication/332174644_How_Maillard_Reaction_Influences_Sensorial_Properties_Color_Flavor_and_Texture_of_Food_Products
57. (PDF) Flavour Profile of Ghee by Gas Chromatography Mass, https://www.researchgate.net/publication/392553698_Flavour_Profile_of_Ghee_by_Gas_Chromatography_Mass_Spectrometry_Gc-Ms_A_Mini_Review
58. chemical composition and aroma description of some volatiles, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20153030687>
59. Part 11 (γ -decalactone, γ -dodecalactone, exaltolide, ambrettolide, 5, <https://www.flavorist.com/gamma-decalactone-gamma-dodecalactone-1-15-pentadecalactone-or-exaltolide-ambrettolide-5-methylfurfural-2-acetylfurfural-theaspirane-2-methyl-tetrahydro-furan-3-one-2-furan-methane-thiol/>

60. Concentrations of δ -decalactone and γ -dodecalactone in, https://www.researchgate.net/figure/Concentrations-of-d-decalactone-and-g-dodecalactone-in-biotransformation-mixtures-after_fig1_398708354
61. Current Developments in Analytical Methods for Advanced Glycation, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12566224/>
62. Validation of a Simple HPLC-Based Method for Lysine ... - MDPI, <https://www.mdpi.com/1420-3049/26/14/4173>
63. Formation of furosine and N-carboxymethyl-lysine (CML) from the, https://www.researchgate.net/figure/Formation-of-furosine-and-N-carboxymethyl-lysine-CML-from-the-Amadori-Maillard-reaction_fig1_306381031
64. Cholesterol oxides: their occurrence and methods to prevent their, https://www.researchgate.net/publication/227603465_Cholesterol_oxides_their_occurrence_and_methods_to_prevent_their_generation_in_foods
65. Kinetic Simulation of Lipid Oxidation and Lycopene Antioxidant, <https://www.mdpi.com/2304-8158/15/9/1522>
66. OXIDATION OF LIPIDS IN FOOD Erwin W'sowicz¹, Anna Gramza¹, <https://journal.pan.olsztyn.pl/pdf-98621-30276?filename=30276.pdf>
67. Kinetic Model of Lipid Oxidation in Foods - ResearchGate, https://www.researchgate.net/publication/229574638_Kinetic_Model_of_Lipid_Oxidation_in_Foods
68. rancimat test - CABI Digital Library, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20163088795>
69. (PDF) Analysis of thermoxidation kinetics of milk fat - ResearchGate, https://www.researchgate.net/publication/323399138_Analysis_of_thermoxidation_kinetics_of_milk_fat
70. (PDF) Oxidative stability of ghee as affected by natural antioxidants, https://www.researchgate.net/publication/273511071_Oxidative_stability_of_ghee_as_affected_by_natural_antioxidants_extracted_from_food_processing_wastes
71. Cholesterol and Its Oxidation Derivatives Content in Market Dairy, <https://www.mdpi.com/2072-6643/16/9/1371>
72. Cholesterol oxidation products – analytical methods and levels in, https://stud.epsilon.slu.se/11840/1/ubhayasekera_k_171016.pdf
73. Analysis, Formation and Inhibition of Cholesterol Oxidation Products, <https://www.jfda-online.com/cgi/viewcontent.cgi?article=2860&context=journal>
74. Impact of cooking techniques on the formation of 7-ketocholesterol, <https://pubmed.ncbi.nlm.nih.gov/41825720/>
75. Monitoring the formation of cholesterol oxidation products in model, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4512089/>
76. Cholesterol and its oxidation products - Riviste UNIMI, <https://riviste.unimi.it/index.php/haf/article/view/9324/9310>
77. Exploring Different Methods for Ghee Production - Agriculture Institute, <https://agriculture.institute/dairy-products-i/exploring-ghee-production-methods/>
78. SOP for Ghee Production Methods | PDF | Butter | Emulsion - Scribd, <https://www.scribd.com/document/519948271/Principles-and-Methods-of-Manufacture-of-Ghee-and-Butter-Oil>

79. Taste of Tradition: Bilona vs Processed Ghee Comparison,
<https://nuclearfarm.com/blogs/news/taste-of-tradition-bilona-vs-processed-ghee-comparison>
80. Changes in Milk lard reaction products in ghee during storage,
https://www.researchgate.net/publication/230809474_Changes_in_Milk_lard_reaction_products_in_ghee_during_storage
81. Gir Cow Ghee: India's Most Revered A2 Ghee | Premium Quality,
<https://www.cowdignity.com/blog/blog-07-gir-cow-ghee>
82. CA3127416A1 - Method for producing ghee - Google Patents,
<https://patents.google.com/patent/CA3127416A1/en>
83. (PDF) Technological and Biochemical Aspects of Ghee (Butter Oil),
https://www.researchgate.net/publication/336121477_Technological_and_Biochemical_Aspects_of_Ghee_Butter_Oil
84. Scrape Surface Heat Exchanger for Food Processing | Shrijee Nut Co.,
<https://www.shrijeenut.com/scrape-surface-heat-exchanger>
85. Developments and Future Perspectives in Scraped Surface Heat,
<https://www.intechopen.com/online-first/1251185>
86. INVESTIGATION OF HEAT TRANSFER CONDITIONS IN SCRAPED,
<https://pp.bme.hu/me/article/download/5520/4625/9278>
87. Application of Scraped Surface Heat Exchanger in Manufacturing of,
https://www.researchgate.net/publication/345543420_Application_of_Scraped_Surface_Heat_Exchanger_in_Manufacturing_of_Dairy_Products_A_Review
88. (PDF) Performance Evaluation of Three-Stage SSHE for Continuous,
https://www.researchgate.net/publication/337459002_Performance_Evaluation_of_Three-Stage_SSHE_for_Continuous_Manufacturing_of_Burfi
89. What is a Scraped Surface Heat Exchanger and What Products,
https://www.fttherm.com/news/What_is_a_Scraped_Surface_Heat_Exchanger_and_What_Products_Does_It_Do-en.html