

The Physical Chemistry and Rheology of Mascarpone Manufacture: A Comprehensive Analysis of Thermo-Acid Coagulation in High-Fat Dairy Emulsions

Introduction to Polyphasic Dairy Emulsions and Mascarpone

Mascarpone is a highly specialized, unripened soft cheese that occupies a fundamentally unique position in the taxonomy of global dairy products. Originating from the Lombardy region of Italy, it is characterized by an exceptionally high fat content, a delicate, slightly sweet flavor profile, and a smooth, glossy, spreadable texture that closely resembles heavy cream or soft butter¹. Unlike the vast majority of traditional cheeses—which rely upon the enzymatic cleavage

of κ -casein via the enzyme rennet (chymosin) combined with mesophilic or thermophilic bacterial fermentation—Mascarpone is manufactured exclusively through the direct thermo-acidic coagulation of a strictly standardized, fat-enriched cream emulsion⁴.

The biochemical, rheological, and physical mechanics governing Mascarpone production represent a profound intersection of fluid dynamics, thermal protein denaturation kinetics, and precise acid-base titration. Under international and regional dairy legislation, such as the German Käseverordnung, Mascarpone is classified within the "Doppelrahmstufe" (double-cream) category, legally mandating a minimum fat in dry matter (FDM) content of 60%, frequently reaching up to 80% to 87% FDM⁶. To achieve these absolute final parameters, the defining phases of its manufacture include the standardization of the starting cream to a lipid content of exactly 20% to 26%, the application of sustained high heat (85°C to 95°C), and the highly controlled volumetric titration of an organic acid—typically citric or tartaric acid—to induce micellar precipitation¹.

Understanding the absolute necessity of each of these highly specific interventions requires an exhaustive, multidisciplinary analysis of dairy physical chemistry. The enrichment of the fat phase fundamentally alters the microstructural geometry and the fat-to-protein ratio, physically modifying the viscoelasticity of the resulting casein network to ensure a soft, spreadable matrix rather than a rigid, rubbery gel¹¹. The extreme thermal treatment intentionally denatures soluble serum whey proteins, forcing their co-precipitation with casein micelles to drastically augment product yield, moisture retention, and structural smoothness⁴. Finally, the precise titration of the acidulant modulates the electrostatic repulsion (zeta potential) of the casein matrix; improper acid selection, or overly rapid administration, inevitably results in localized isoelectric shock,

structural graininess, and catastrophic fat loss¹.

Lipid Extraction and the Physics of Cream Standardization

The initial phase of Mascarpone production requires the extraction and concentration of bovine milk fat to achieve the critical 20-25% lipid threshold. Bovine milk functions simultaneously as a colloidal dispersion of casein micelles and an oil-in-water emulsion consisting of liquid triglyceride droplets encapsulated within a complex phospholipid Milk Fat Globule Membrane (MFGM)¹⁴.

Historical Gravity Separation and Cryoagglutination

Prior to the advent of high-speed centrifugal separators, the extraction of cream for artisanal Mascarpone relied exclusively on manipulating natural phase separation utilizing gravity and ambient thermodynamics⁴. The macroscopic separation of cream from skim milk plasma is strictly governed by classical fluid dynamics, explicitly described by Stokes' Law, which models the terminal upward velocity of a rising spherical fat globule based on the density differential between the lipid and aqueous phases, the dynamic viscosity of the plasma, and the square of the globule's radius¹⁴.

While Stokesian mechanics dictate that chilling the milk should theoretically retard separation by exponentially increasing the dynamic viscosity of the plasma, historical dairy methodologies deliberately utilized extreme cold (4°C to 7°C) in shallow setting pans to dramatically accelerate the creaming process¹². This profound physical paradox is resolved through immunology. Rapid cold-setting activates native Immunoglobulin M (IgM), a massive pentameric cryoglobulin naturally secreted into raw milk¹. At cold temperatures, IgM exhibits an extraordinarily high binding affinity for specific poly-immunoglobulin G receptors located on the outer bilayer of the MFGM¹⁴.

These pentameric IgM complexes act as structural bridges, physically cross-linking adjacent, microscopic fat globules into massive, macroscopic clusters measuring hundreds of micrometers in diameter¹. Because Stokes' Law dictates that buoyant velocity scales with the square of the particle radius, these massive agglutinated clusters easily overcome the increased viscous drag of the cold plasma, resulting in rapid vertical migration and the swift formation of a concentrated cream layer¹. This passive, highly quiescent gravity separation exerts virtually zero mechanical shear on the lipid emulsion, leaving the delicate, trilayered MFGM entirely intact and preserving critical highly conserved membrane glycoproteins, notably butyrophilin and xanthine oxidase, which are essential for emulsion stability¹⁴.

Modern Centrifugal Separation and Hydrodynamic Shear

While traditional gravity separation perfectly preserves the MFGM, the prolonged 12 to 48-hour cold-setting period introduces severe microbiological vulnerabilities. The cold environment selectively encourages the proliferation of psychrotrophic bacteria (e.g., *Pseudomonas* species), which synthesize massive volumes of highly thermostable extracellular lipases¹⁴. These enzymes easily survive subsequent pasteurization, aggressively hydrolyzing the triglycerides in the cream to release short-chain free fatty acids that impart highly objectionable rancid, soapy,

and bitter organoleptic defects¹¹. Furthermore, gravity separation acts as a particle size fractionator, leaving a highly significant residual fat content (0.1% to 0.5%) in the skim milk due to the overriding force of Brownian motion on micro-globules¹³. Consequently, the contemporary global dairy industry, including industrial Mascarpone production, universally employs continuous, high-speed centrifugal separators to achieve the 20-25% target¹⁴. By subjecting the fluid to centripetal acceleration thousands of times greater than standard gravity, these continuous systems bypass the limitations of Brownian motion entirely, recovering nearly 100% of the available commercial butterfat and reducing processing times from days to minutes¹⁴. However, the violent hydrodynamic shear stress generated within the tight disc stacks of a continuous centrifuge inflicts profound physical damage upon the MFGM. This mechanical shear forcibly strips away up to 24% of the native polar lipids and severs the delicate intermolecular disulfide bonds anchoring butyrophilin and xanthine oxidase, depositing them into the skim milk phase¹¹. The partial destruction of the MFGM exposes the naked, hydrophobic triglyceride core, fundamentally altering the interfacial tension of the cream and necessitating downstream interventions, such as high-pressure homogenization, to re-stabilize the emulsion before thermal processing¹⁴.

Separation Methodology	Primary Motive Force	Processing Time	Residual Fat in Skim	MFGM Structural Integrity	Microbiological Risk
Traditional Gravity Setting	1g + IgM Cryoagglutination	12 to 48 hours	0.10% - 0.50%+	Entirely intact; glycoproteins preserved	High risk of psychrotrophic thermostable lipolysis
Modern Centrifugal Separation	Centripetal Acceleration (>5000g)	Continuous / Minutes	0.01% - 0.07%	Severely damaged; severe polar lipid stripping	Negligible due to immediate processing

Table 1: Physical, biochemical, and operational comparison between traditional gravity setting and continuous centrifugal separation for the extraction of cream.
[cite: 11, 14, 19, 28, 30, 31, 33]

Emulsion Physics and the 20-25% Lipid Threshold

Once the cream is extracted, it must be meticulously standardized to a fat content of 20.0% to 26.0% prior to any application of heat or acid¹. The fundamental necessity of this precise

enrichment threshold is rooted in the targeted macroscopic composition of the final product and the exacting microstructural requirements of the cheese matrix.

Phase Volume and Matrix Interruption

The rheology and macroscopic texture of any cheese are dictated entirely by the continuous protein network, which functions as the structural scaffolding of the gel. The dispersed lipid droplets act as inactive, stress-concentrating filler particles that physically interrupt the cross-linking of the casein micelles³⁴. In standard cheesemaking utilizing low-fat whole milk (3.5% fat), the casein micelles exist in close spatial proximity. Upon coagulation, these micelles form extensive, highly interlinked networks dominated by strong electrostatic forces, hydrophobic interactions, and internal hydrogen bonding¹¹. This dense, highly cross-linked protein architecture manifests as a firm, rubbery, and highly elastic texture, which is desirable in hard cheeses but represents a catastrophic structural defect in soft-spread products¹¹. By artificially enriching the starting cream to 20-25% fat, the processor radically alters the spatial geometry of the emulsion. The massive influx of fat globules vastly increases the total dispersed phase volume, physically forcing the casein micelles further apart within the aqueous plasma¹¹. When the proteins are subsequently precipitated via acid titration, the resulting matrix is heavily discontinuous; the vast quantity of interspersed fat globules physically obstructs widespread casein-casein aggregation¹¹.

Advanced microstructural analyses utilizing Confocal Laser Scanning Microscopy (CLSM) and Cryogenic Scanning Electron Microscopy (Cryo-SEM) visually confirm this phenomenon³⁶. These techniques, which allow hydrated samples to be examined without chemical dehydration artifacts, reveal that high-fat cream cheeses exhibit a highly open, porous protein network heavily disrupted by massive lipid aggregates¹¹. This structurally compromised network yields easily under shear stress, delivering the characteristic "melt-in-the-mouth" spreadability and high oral lubrication expected of Mascarpone¹¹.

Homogenization Pressures and Recombinant Membranes

In modern industrial settings, the standardized 20-25% cream is frequently subjected to high-pressure homogenization prior to thermal treatment²⁹. This process ensures emulsion stability and prevents spontaneous phase separation (free oil release) during the intense heating phase. The cream is forced through microscopic annular valves under pressures typically ranging from 10 to 30 MPa (100 to 300 bar)³².

This extreme mechanical pressure shears the massive native fat globules into uniform, sub-micron droplets, exponentially increasing the total surface area of the lipid phase³². Because the native MFGM was largely destroyed during centrifugal separation, the emulsion is now in a critically unstable state. To lower the massive interfacial tension and re-stabilize these naked, newly sheared lipid droplets, casein micelles and soluble whey proteins are rapidly and forcefully adsorbed from the plasma onto the lipid-water interface, creating a "recombined" artificial membrane³².

This forced adsorption profoundly alters the subsequent coagulation kinetics. The protein-coated fat globules cease to act as entirely passive fillers; they essentially behave as massive pseudo-protein particles that actively cross-link and participate in the subsequent

acid-induced gelation network³². This increases the yield stress, dynamic storage modulus (G'), and overall structural firmness of the resulting Mascarpone, which can be measured empirically using small-amplitude oscillatory rheometry³². However, precise control is required; over-homogenization, or the homogenization of creams with fat contents significantly exceeding 25%, leads to the formation of massive homogenization clusters. This occurs due to a critical absolute shortage of available plasma protein to coat the exponentially increased lipid surface area, resulting in severe textural graininess and a loss of spreadability³⁹.

Rheology, Viscoelasticity, and Yield Stress Mechanics

Following the coagulation of the 20-25% standardized cream, the fluid physically transitions from a liquid emulsion into a weak, viscoelastic particulate gel¹³. The exact rheological behavior of this newly formed coagulum is highly complex and heavily non-Newtonian.

The Herschel-Bulkley Fluid Dynamics Model

The flow dynamics of the Mascarpone matrix are best characterized mathematically by the Herschel-Bulkley model, a non-linear approach to modeling complex, shear-thinning viscoelastic fluids⁴³. The mathematical equation is defined as:

$$\tau = \tau_0 + K(\dot{\gamma})^n$$

Where:

- τ represents the applied shear stress.
- τ_0 represents the yield stress, the absolute minimum threshold of force required to initiate flow.
- K represents the consistency coefficient.
- n represents the flow behavior index.
- $\dot{\gamma}$ represents the shear rate.

The formation of the casein-whey-fat matrix generates a highly distinct yield stress (τ_0).

Because the flow behavior index (n) is less than 1, the product exhibits profound shear-thinning behavior; it acts as an anisotropic, viscoelastic solid and maintains its structural shape at rest, but flows smoothly and spreads easily when the applied force of a knife or the shear of mastication exceeds the yield stress threshold⁴¹. If the fat content were lower than the 20-25% threshold, the increased casein-casein cross-linking would exponentially increase the yield stress and the dynamic storage modulus (G'), resulting in a tough, rubbery texture that would stubbornly resist spreading¹¹.

Rheological Defects: Syneresis, Storage, and Hydrocolloid Interventions

The preservation of this specific viscoelastic state is highly dependent on storage parameters. Extended frozen storage (-18°C to -20°C) is highly detrimental to the functionality of Mascarpone and high-fat cream cheeses²³. The thermodynamics of ice crystallization during freezing cause severe localized dehydration of the casein micelles, creating extreme solute concentration gradients within the matrix²³. This cryoconcentration causes irreversible physical destabilization of the proteins, leading to a loss of internal cohesion, massive whey syneresis upon thawing, and a profound deterioration in the product's fundamental spreadability and lubrication properties²³.

To mitigate these risks or to simulate the mouthfeel of 50% fat Mascarpone in reduced-fat variants, dairy technologists frequently utilize exogenous hydrocolloids and fat replacers¹¹. The

incorporation of κ -carrageenan and ι -carrageenan, for example, fundamentally alters the rheology. Carrageenans possess highly sulfated groups that engage in strong electrostatic interactions with the casein micelles, forming double-helical structures that increase the complex modulus (G^*) and gel strength⁵⁰. Furthermore, these hydrocolloids swell by binding massive volumes of water, which reduces ice recrystallization, improves freeze-thaw stability, and mimics the high-lubrication mouth-coating sensation naturally provided by high phase-volume lipids¹¹.

Similarly, the incorporation of microparticulated β -glucan and emulsified phytosterols acts to mechanically break casein-casein interactions, softening the texture of reduced-fat matrices to mirror the specific dominance rates of creaminess observed in traditional high-fat Mascarpone¹¹.

Thermal Processing: Denaturation Kinetics and Matrix Integration

Following lipid standardization and potential homogenization, the 20-25% cream is subjected to a severe, sustained thermal treatment, strictly controlled between 85°C and 95°C. This extreme heat is not applied merely for standard microbial pasteurization (which only requires 72°C for 15 seconds); rather, it is the primary biochemical driver of the cheese's final physical structure, textural smoothness, and volumetric yield¹.

Kinetics of β -Lactoglobulin and α -Lactalbumin Unfolding

Bovine milk serum contains highly soluble, globular whey proteins, the most abundant of which are β -lactoglobulin (β -Lg) and α -lactalbumin (α -La), representing roughly 50-60% and 20% of total whey proteins, respectively⁵¹. In standard, unheated milk, these globular proteins are thermodynamically stable, highly folded structures. Their hydrophobic amino acid residues and highly reactive free sulfhydryl (thiol) groups are buried deep within their tertiary β -barrel structures, completely shielded from the aqueous solvent⁵². Because they remain highly soluble in the aqueous phase, native whey proteins do not naturally participate in rennet-induced coagulation and are entirely lost in the whey drainage of traditional cheeses (like Cheddar or Gouda), representing a significant loss of potential nutritional and economic yield⁴.

The fundamental biochemical purpose of heating the cream to 85-95°C is to forcibly and irreversibly denature these whey proteins. The thermal denaturation of β -Lg is a complex, multistage kinetic reaction⁵³. As the temperature of the cream exceeds 65°C, the applied thermal kinetic energy overcomes the intramolecular hydrogen bonds and van der Waals forces stabilizing the protein. This causes the β -Lg molecule to dissociate from its native dimeric state into monomers, and undergo a profound conformational shift into a partially unfolded, intermediate "molten globule" state⁵³. This partial unfolding exposes the previously buried hydrophobic domains to the aqueous environment and, critically, exposes the highly reactive free sulfhydryl group located at the Cysteine-121 (Cys121) residue⁵³. The reaction kinetics governing this denaturation are best described mathematically as an apparent 1.5-order reaction for β -Lg⁵¹. Thermodynamic modeling utilizing Arrhenius plots demonstrates an abrupt discontinuity or "bend" in the reaction rate constant between 80°C and 90°C. This indicates a fundamental shift in the reaction kinetics: below this threshold, the reaction is limited by the rate of physical unfolding (requiring extremely high activation energies between 280 and 524 kJ/mol), whereas above 85°C, the unfolding happens almost instantaneously, and the overall reaction rate becomes entirely limited by the speed of protein-protein aggregation⁵¹.

Co-precipitation and Covalent Interaction with κ -Casein

Once the highly reactive Cys121 thiol group is exposed to the hot solvent at 85-95°C, it becomes extremely susceptible to thiol-disulfide interchange reactions⁵³. The denatured β -Lg monomers aggressively seek out and react with other available sulfur-containing proteins to minimize their free energy. The primary target for this interchange is κ -casein, the stabilizing protein that naturally resides on the outer, "hairy" surface of the suspended casein micelles²⁷. The denatured β -Lg, and subsequently α -La, chemically and covalently bond to the κ -casein via strong intermolecular disulfide bridges, effectively coating the exterior of the casein micelles in a thick, dense layer of denatured whey protein⁵². Additionally, these denatured whey proteins complex with the proteins newly adsorbed onto the recombinant Milk Fat Globule Membrane, creating extensive, rigid cross-linkages between the aqueous protein phase and the dispersed lipid phase⁵. This heat-induced co-precipitation is the defining structural hallmark of Mascarpone. By chemically welding the highly hydrophilic whey proteins onto the casein micelles *prior* to acid coagulation, the dairy processor ensures that the whey proteins are permanently, physically trapped within the cheese matrix during the subsequent draining and separation phases⁴. The retention of these heavily hydrated whey proteins dramatically increases the final volumetric yield of the cheese. Furthermore, this bound hydration layer significantly softens the resulting matrix and actively restricts the violent contraction (syneresis) of the casein network, preserving the delicate, creamy, high-moisture texture that defines the product⁴.

Scraped Surface Heat Exchangers (SSHE) and Emulsion Preservation

While sustained high heat is biochemically essential for whey denaturation and yield enhancement, meticulous thermal management is required to preserve the physical stability of the 20-25% lipid emulsion. At temperatures of 85-95°C, the triglycerides comprising the core of the fat globules are entirely molten, existing purely as liquid oil¹. The macroscopic stability of these liquid oil droplets is entirely dependent upon the structural integrity of the surrounding protein/phospholipid membrane.

If the cream is subjected to excessive hydrodynamic shear, turbulent flow, or localized surface boiling while at these extreme temperatures, the delicate membrane will rupture¹. The destruction of the membrane allows the molten lipid cores of adjacent globules to undergo rapid, irreversible coalescence. This localized melting and pooling of free oil entirely destroys the emulsion architecture, resulting in severe fat loss into the whey, drastically depressed yields, and a highly defective, greasy, and uneven final texture¹.

Furthermore, denatured β -Lg is highly prone to adhering to hot metallic surfaces, causing severe fouling and scaling ("burn-on") in standard plate or tubular heat exchangers, which destroys heat transfer efficiency⁵⁴. To circumvent these massive physical and engineering challenges, industrial Mascarpone production relies on advanced Scraped Surface Heat Exchangers (SSHE)⁶¹. Systems such as the Gerstenberg Schröder Kombinator, the Rotapro, and the Hercules utilize an inner cylinder surrounded by a heating jacket. A central rotating shaft equipped with floating scraper blades continuously physically sweeps the inner wall of the cylinder⁶¹.

This continuous mechanical scraping achieves three critical outcomes: it violently prevents the highly viscous, denatured whey proteins from adhering to and fouling the heat transfer surface; it provides instantaneous, uniform heat transfer to the highly viscous fluid; and it gently mixes the product without inducing the high-velocity hydrodynamic shear forces that would otherwise fracture the molten fat globules and destroy the emulsion⁶¹. The SSHE effectively manages the complex rheology of the high-fat matrix while safely achieving the exact 85-90°C target required for optimal β -Lg unfolding.

Heat Exchanger Type	Mechanism of Heat Transfer	Shear Profile	Fouling Resistance	Suitability for Mascarpone 25% Cream
Plate Heat Exchanger	Corrugated metal plates	High turbulence/shear	Extremely Poor	Low; causes fat coalescence and severe protein burn-on.

Tubular Heat Exchanger	Concentric tubes	Moderate shear	Poor	Low; struggles with high viscosity and protein fouling.
Scraped Surface (SSHE)	Jacketed cylinder with internal rotating scraper blades	Gentle, continuous sweeping	Excellent	Optimal; prevents fouling, manages extreme viscosity, protects molten fat globules.

Table 2: Comparison of thermal processing equipment and their physical suitability for handling the highly viscous, thermally sensitive 20-25% cream matrix during the 85-95°C holding period.
[cite: 54, 61, 62, 63, 64]

The Physical Chemistry of Non-Enzymatic Coagulation

Unlike traditional hard and soft cheeses (e.g., Cheddar, Gouda, or Brie), the coagulation of Mascarpone relies entirely on thermo-acidic precipitation, completely omitting the use of rennet (chymosin)¹. The mechanism of this direct chemical precipitation involves the fundamental manipulation of the electrical double layer surrounding the casein micelles in the presence of extreme heat.

Zeta Potential and Isoelectric Neutralization

In their native state at a physiological pH of 6.7, casein micelles are highly stable, suspended colloidal particles. This stability is provided almost exclusively by the dense layer of κ -casein protruding from the micelle surface, which imparts both a profound steric hindrance (a physical "hairy" barrier preventing close contact) and a strong net negative electrostatic charge, known as the zeta potential¹³. The zeta potential of native casein micelles typically measures approximately -20 mV¹³. The strong electrostatic repulsion between adjacent, negatively charged micelles prevents them from colliding and aggregating, maintaining the fluid, liquid state of the cream despite the high concentration of proteins¹³.

In traditional enzymatic cheesemaking, rennet is introduced to specifically and exclusively cleave the protruding, hydrophilic macropeptide tail of the κ -casein²⁷. This enzymatic severing physically removes the steric barrier and strips away the negative charge, allowing the structurally naked micelles to instantly aggregate via calcium bridging and hydrophobic

interactions at near-neutral pH¹³.

Mascarpone bypasses this highly specific enzymatic pathway entirely through the direct volumetric titration of organic acids. As the acidulant is introduced into the hot cream, it

dissociates and releases free protons (H^+), causing a rapid decline in the pH of the plasma¹². As the pH drops, the excess protons rapidly protonate the highly acidic carboxyl and phosphoserine residues located on the surface of the casein micelles⁶⁷. This progressive protonation systematically neutralizes the net negative charge of the micelle, driving the zeta potential from -20 mV steadily toward zero¹³.

Temperature-Dependent Isoelectric Shifts and Coagulation pH

In cold or ambient milk, casein micelles reach their absolute isoelectric point (pI)—the specific thermodynamic point of zero net electrical charge and maximum insolubility—at a pH of exactly 4.6⁶⁷. At this specific pH, electrostatic repulsion is completely annihilated, and the proteins spontaneously precipitate to form a gel⁶⁷.

However, the physical chemistry of the system is radically and permanently altered by the extreme thermal environment utilized in Mascarpone production. Because the cream is held at 85°C to 95°C during acidification, the thermodynamic activation energy required for the micelles to overcome their remaining repulsive barriers is drastically lowered⁴². The elevated kinetic energy of the hot fluid vastly increases the frequency and violence of micellar collisions¹³.

Furthermore, as the pH is lowered, the Colloidal Calcium Phosphate (CCP) that acts as the internal structural glue of the micelle begins to rapidly solubilize into the aqueous phase, causing the micelles to swell and loosen, reaching a highly unstable transition state between pH 5.5 and 5.0¹³. This CCP solubilization generates a massive chemical buffering peak in the fluid

around pH 5.0⁷². Critically, the thick layer of denatured whey proteins (β -Lg and α -La) that was covalently bound to the outside of the casein micelles during the heating phase physically shifts the effective isoelectric point of the entire co-precipitate complex⁵⁷.

Consequently, the high-heat, whey-coated micelles do not require the pH to fall all the way to 4.6 to precipitate. Instead, massive, spontaneous coagulation occurs at a significantly higher pH, typically between 5.7 and 6.6⁴. As the pH crosses this critical, elevated threshold, the kinetic energy of the hot fluid easily overcomes the artificially diminished electrostatic repulsion.

Hydrophobic interactions, van der Waals forces, and hydrogen bonding instantly dominate, causing the micelles to rapidly collide, fuse, and cross-link into a macroscopic, continuous tridimensional gel network that physically entraps the entire 25% lipid phase within its interstitial spaces¹³.

Precision Titration: Selection and Impact of Specific Organic Acids

The chemical agent utilized to drive this isoelectric precipitation is of paramount technical and organoleptic importance. The selection of the specific organic acid, its exact concentration, and the physical kinetics of its administration directly govern the final structural uniformity and the temporal sensory profile of the Mascarpone. Historically and commercially, the acids utilized

include citric acid, tartaric acid, acetic acid (vinegar), malic acid, and lactic acid¹.

Biochemical Profiles and Dissociation Constants

The physical efficacy of an acidulant is dictated by its molecular structure and its specific acid dissociation constants (pK_a), which define its thermodynamic willingness to release protons into the aqueous matrix, and critically, its subsequent buffering capacity near the target pH.

- **Citric Acid:** The vast majority of modern industrial Mascarpone is coagulated utilizing citric acid ($C_6H_8O_7$)¹. Citric acid is a weak, triprotic organic acid, meaning it possesses three separate carboxyl groups capable of sequential deprotonation. The three corresponding dissociation constants are $pK_{a1} = 3.13$, $pK_{a2} = 4.76$, and $pK_{a3} = 6.40$ ⁷⁹. Because the third dissociation constant (pK_{a3}) sits remarkably close to the target, high-heat coagulation pH of 5.8-6.6, citric acid provides an exceptional, highly specific chemical buffering capacity precisely at the critical moment of micellar precipitation⁷⁹. This massive buffering effect physically prevents the pH from crashing too rapidly or unpredictably, allowing the dairy technologist to exert exquisite, slow control over the rate of isoelectric aggregation⁸¹.
- **Tartaric Acid:** Traditionally utilized in historical artisanal Lombardy recipes, tartaric acid ($C_4H_6O_6$) is a diprotic acid featuring dissociation constants of $pK_{a1} = 2.98$ and $pK_{a2} = 4.34$ ². Because it completely lacks a pK_a in the 5.5 to 6.5 range, it offers far less buffering capacity at the exact point of Mascarpone coagulation, resulting in a much sharper, more aggressive, and harder-to-control pH drop⁸².
- **Acetic Acid (Vinegar):** Acetic acid (CH_3COOH) is a simple monoprotic acid with a single pK_a of 4.76. While functionally capable of acting as a coagulant, its high volatility and specific organoleptic properties make it highly problematic in modern, delicately flavored dairy applications¹.

Sensory Psychophysics and Time-Intensity Profiling

The choice of acidulant extends far beyond fluid physical chemistry; it dictates the temporal sensory profile of the cheese. Mascarpone is globally renowned for its mild, slightly sweet flavor, heavily dominated by ketones (such as 2-heptanone and 2-pentanone)¹⁰. This delicate profile must remain perfectly balanced without the overwhelming, lingering tanginess typical of fermented products¹.

Sensory threshold analysis, time-intensity mapping, and adherence to the Weber-Fechner law of psychophysics reveal profound differences in how the human palate perceives these coagulants in a high-fat matrix.

- **Citric Acid ("The Sprinter"):** Elicits a fast, sharp, and highly aggressive "burst" of sourness that reaches its maximum intensity almost immediately, but critically, dissipates very rapidly⁸⁵. This rapid clearance from the palate is highly desirable in Mascarpone, as it

provides a bright initial note that quickly yields to the rich, creamy sweetness of the dairy fat without leaving a lingering, acidic aftertaste⁸⁵. Furthermore, sensory studies demonstrate that the presence of citric acid specifically increases the absolute perception threshold for sucrose, altering the sweet-sour dynamic to feel more balanced⁸⁶.

- **Acetic Acid:** Imparts a pungent, highly persistent sourness and a distinct volatile aroma that clashes severely with the mild, sweet profile of the cream. In extensive industrial trials, acetic acid formulations were rapidly discontinued due to overwhelmingly negative sensory feedback indicating an "overly sour" and unbalanced taste¹.
- **Tartaric Acid ("The Dry Punch"):** Possesses a highly aggressive, astringent, and distinctly "dry" sensory attack that can induce a severe puckering sensation⁸². While historically accurate for the style, its intense astringency can mask the delicate dairy notes, leading to its near-universal replacement by citric acid in modern commercial formulations⁷⁶.
- **Malic Acid ("The Marathon Runner"):** Exhibits a slow onset but a long, persistent finish that lingers on the palate⁸⁵. This lingering sourness is generally avoided in Mascarpone, as it competes with the desired rich, fatty finish of the cream⁷⁸.

Acidulant	Classification	Dissociation Constants (pKa)	Buffering Efficacy Near Coagulation pH (5.8-6.6)	Sensory Time-Intensity Profile
Citric Acid	Triprotic	3.13, 4.76, 6.40	Extremely High	Fast onset, rapid dissipation; clean finish preserving fat sweetness.
Tartaric Acid	Diprotic	2.98, 4.34	Low	Intense, aggressive attack; induces dryness and strong astringency.
Acetic Acid	Monoprotic	4.76	Low	Pungent, persistent aroma; imparts lingering, negative

				sourness.
Malic Acid	Diprotic	3.40, 5.05	Low	Slow onset, lingering plateau; persistent sour finish.

Table 3: Chemical parameters, dissociation constants, and empirical psychophysical sensory profiles of organic acidulants utilized in dairy thermo-acid coagulation.

[cite: 1, 79, 82, 85]

Mixing Dynamics and the Prevention of Localized Isoelectric Shock

The physical administration of the chosen acidulant into the hot cream represents one of the most critical mechanical failure points in the entire manufacturing process. When a concentrated organic acid is poured into the 85°C cream, the immediate microscopic fluid environment surrounding the acid droplets experiences a violent, localized pH crash, instantly dropping far below the 5.7 threshold and violently precipitating the proteins in that exact microscopic vicinity¹. If the cream is static or poorly agitated, this localized precipitation forms massive, uneven, highly dense protein clumps. The fat globules become trapped in disorganized protein aggregates rather than a smooth, continuous matrix¹. Macroscopically, this rheological defect manifests as a severely uneven, grainy, or "sandy" texture, entirely ruining the glossy, smooth finish expected of a premium Mascarpone¹.

To definitively prevent this structural defect, contemporary production protocols demand continuous, high-shear stirring while the acidifying agent is continuously and slowly titrated into the vortex¹. This intense hydrodynamic mixing immediately disperses the acidulant throughout the entire volume of the fluid, preventing localized isoelectric shock and ensuring that the entire batch crosses the critical pH threshold simultaneously, resulting in a perfectly uniform, smooth, and glossy particulate gel network¹.

Downstream Processing, Whey Drainage, and Yield Optimization

Immediately following the addition of the acid and the achievement of the target pH, the cream physically transitions into the weak, viscoelastic particulate gel¹³. However, this freshly coagulated matrix still contains the entirety of the original water volume from the milk plasma. To achieve the dense, rich texture of the final product, the coagulum must undergo a process of syneresis and physical separation to expel the excess moisture (whey) and concentrate the fat and protein fractions to the required 50-55% absolute fat level²⁹.

Traditional Gravity Drainage vs. High-Speed Centrifugation

Historically, syneresis was achieved by pouring the hot coagulum into fine linen or muslin cloth

bags and allowing gravity and passive capillary action to drain the whey over a prolonged period of 20 to 24 hours in a cold environment (4°C to 10°C)¹. While effective for artisanal, small-batch production, this method introduces severe microbiological vulnerabilities and creates inconsistencies in the final moisture content, leading to the uneven textures noted in classical production¹.

To circumvent these risks, drastically reduce processing time, and ensure exact compositional standardization, modern continuous dairy operations utilize advanced centrifugal separators or Tangential Flow Filtration (TFF)¹⁶. The hot coagulum can be pumped directly into specialized, self-cleaning centrifugal separators designed specifically for high-viscosity, high-fat fluids (Double Cream Separators)²⁹. The extreme centripetal force instantly fractions the heavy aqueous whey from the lighter, highly concentrated lipid-protein curd, recovering the solid mass with extraordinary speed and allowing for immediate hot-filling into sterile packaging, thereby bypassing the 24-hour vulnerability window entirely²⁹.

Ultrafiltration (UF) and Membrane Fouling Mechanics

Alternatively, cross-flow membrane ultrafiltration (UF) is increasingly utilized to selectively force water, lactose, and soluble minerals (the permeate) through a microscopic ceramic or polymeric membrane under pressure, while perfectly retaining the concentrated fat and denatured protein matrix (the retentate)¹⁶. This method completely eliminates the loss of residual fat and soluble proteins in the whey, driving yield efficiency to near 100% and providing absolute, real-time control over the final total solids composition¹⁶.

However, the ultrafiltration of highly acidic, high-protein whey and curd introduces the severe engineering challenge of membrane fouling. During operation, a rapid, initial decline in permeate flux occurs due to concentration polarization—the accumulation of a dense layer of retained solutes immediately adjacent to the membrane surface⁵⁹. This is rapidly followed by severe, long-term irreversible fouling caused by the physical deposition and adsorption of denatured whey proteins, humic acids, and colloidal silica into the pores of the membrane, mathematically modeled by cake filtration and intermediate pore-blocking dynamics⁵⁹.

Because the isoelectric point of the fouled membrane shifts to the alkaline range, cleaning the UF system requires complex and aggressive Clean-In-Place (CIP) protocols. This typically involves alkaline agents like Sodium Hydroxide (NaOH) combined with strong oxidative agents like Sodium Hypochlorite (NaClO) to solubilize the organic foulants and restore the permeate flux from heavily degraded levels back to operational baselines⁶⁰.

Conclusion

The manufacture of Mascarpone represents a profound, highly sophisticated exercise in dairy physical chemistry, operating on mechanisms fundamentally distinct from standard enzymatic cheesemaking protocols. The absolute necessity of standardizing the initial cream to 20-25% fat is driven by microstructural geometry; the massive influx of lipid volume physically interrupts extensive casein-casein crosslinking, ensuring the final network exhibits the low yield stress necessary for a soft, highly spreadable texture rather than a rubbery gel.

The application of intense, sustained thermal processing (85-95°C) within Scraped Surface Heat

Exchangers leverages the specific thermodynamic unfolding kinetics of β -lactoglobulin. By exceeding the 80°C threshold, the reaction shifts to an aggregation-limited state, initiating rapid thiol-disulfide interchange reactions that covalently weld the highly hydrophilic whey proteins

directly to the κ -casein on the micelle surface, drastically augmenting final yield and moisture retention while preserving the molten fat emulsion.

Finally, the precise, high-shear volumetric titration of specific organic acids—overwhelmingly

citric acid due to its superior buffering capacity at pK_{a3} and its rapid, non-lingering sensory profile—drives the isoelectric neutralization of the casein micelles. Because the extreme heat lowers the thermodynamic barrier for aggregation and solubilizes colloidal calcium phosphate, precipitation occurs rapidly at a remarkably high pH of 5.7 to 6.6. By avoiding localized isoelectric shock through continuous hydrodynamic mixing, and efficiently recovering the resulting coagulum via high-speed centrifugation or ultrafiltration, dairy technologists perfectly execute the creation of a pristine, heavily hydrated, non-Newtonian fat-protein matrix.

Works cited

1. Mascarpone Cheese Production - Узловский молочный комбинат, <https://uzmolkom.com/en/blog/mascarpone-cheese-production/>
2. Mascarpone vs Cream Cheese: Fat Content, Texture Differences, <https://world.taobao.com/lang/en-us/shopping-guide/2030082023363706880.htm>
3. Mascarpone: the Italian cream with a velvety texture | WikiCibo, <https://wikicibo.com/en/prodotti/latticini-e-formaggi/mascarpone>
4. Unveiling the Molecular Basis of Mascarpone Cheese Aroma: VOCs, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7179404/>
5. Mediterranean milk and milk products | Request PDF - ResearchGate, https://www.researchgate.net/publication/8647679_Mediterranean_milk_and_milk_products
6. Doppelrahmkäse - Wikipedia, <https://de.wikipedia.org/wiki/Doppelrahmk%C3%A4se>
7. Was ist Mascarpone? - eat.de, <https://eat.de/magazin/was-ist-mascarpone/>
8. Lebensmittel: Kennzeichnung des Fettgehalts bei Käse, https://www.lgl.bayern.de/lebensmittel/warengruppen/wc_03_kaese/et_fett_kaese.htm
9. Microbial dynamics during shelf-life of industrial Ricotta cheese and, https://www.researchgate.net/publication/288889564_Microbial_dynamics_during_shelf-life_of_industrial_Ricotta_cheese_and_identification_of_a_Bacillus_strain_as_a_cause_of_a_pink_discolouration
10. Unveiling the Molecular Basis of Mascarpone Cheese Aroma: VOCs, https://www.researchgate.net/publication/339843494_Unveiling_the_Molecular_Basis_of_Mascarpone_Cheese_Aroma_VOCs_analysis_by_SPME-GCMS_and_PTR-ToF-MS
11. s4365780_phd_finalthesis.pdf - UQ eSpace,

- https://espace.library.uq.edu.au/view/UQ:5c18970/s4365780_phd_finalthesis.pdf
12. PHYSICAL AND CHEMICAL CHARACTERIZATION OF ... - novomar,
https://novomar.cetmar.org/wp-content/uploads/2015/10/PDF_24.pdf
 13. Micellar Transition State in Casein Between pH 5.5 and 5.0,
https://www.researchgate.net/publication/229502095_Micellar_Transition_State_in_Casein_Between_pH_5.5_and_5.0
 14. 81 Gravity Cream Separation Physics.pdf
 15. (PDF) An alternative method for serum protein depletion/enrichment,
https://www.researchgate.net/publication/271589958_An_alternative_method_for_serum_protein_depletionenrichment_by_precipitation_at_mildly_acidic_pH_values_and_low_ionic_strength
 16. Cheese Production by Ultrafiltration (UF) Using Tangential Flow,
<https://perinox.com/en/cheese-production-ultrafiltration-tangential-flow-membrane-filtration/>
 17. Cheese Production- Manufacturing scheme (basic recipe),
https://dca.au.dk/fileadmin/DJF/Kontakt/Besog_DJF/Oevelsesvejledning_og_baggrundsmateriale/Cheese_Production_procedure.pdf
 18. Protein Precipitation by Metal Hydroxides as a Convenient ... - MDPI,
<https://www.mdpi.com/1420-3049/30/1/2>
 19. Effect of coagulants, stage of cream addition and inulin incorporation,
<https://epubs.icar.org.in/index.php/IJDS/article/view/76570>
 20. Association of denatured whey proteins with casein micelles in,
<https://pubmed.ncbi.nlm.nih.gov/12617395/>
 21. Käsegruppen - so wird Käse eingeteilt,
<https://www.schoenegger.com/kaesewissen/kaesegruppen>
 22. Einteilung von Käse - Käsekompass NRW,
<https://www.kaesekompass-nrw.de/kaeseinfos/infos-im-ueberblick/einteilung-von-kaese>
 23. Frozen storage is detrimental to the functionality of cream cheese,
https://www.researchgate.net/publication/401012042_Frozen_storage_is_detrimental_to_the_functionality_of_cream_cheese_and_mascarpone_even_in_high-pressure-treated_samples
 24. Status and Technological Aspects of Fresh Cheeses,
<https://basu.org.in/wp-content/uploads/2020/04/Status-and-Technological-Aspects-of-Fresh-Cheeses.pdf>
 25. Determination of the yield stress by the Herschel–Bulkley model and,
https://www.researchgate.net/figure/Determination-of-the-yield-stress-by-the-Herschel-Bulkley-model-and-the-Casson-model-for_fig10_373673885
 26. On the Response of a Herschel–Bulkley Fluid Due to a Moving Plate,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9502123/>
 27. Fundamentals of Cheese Science | Request PDF - ResearchGate,
https://www.researchgate.net/publication/311994895_Fundamentals_of_Cheese_Science
 28. Protein precipitation: a comprehensive guide - Abcam,
<https://www.abcam.com/en-us/knowledge-center/proteins-and-protein-analysis/pro>

[tein-precipitation](#)

29. Mascarpone, what is it? - GEA,
<https://www.gea.com/en/campaigns/fresh-cheese-uncovered/mascarpone/>
30. Perspective of Membrane Technology in Dairy Industry: A Review,
<https://www.scribd.com/document/539825360/Milk-Journal>
31. Principles of Filtration for Milk, Cheese & Dairy Products,
<https://www.cdic.net/wp-content/uploads/2026/03/9.-Eric-Terada-Principles-of-Filtration-for-Milk-Cheese-and-Dairy-Products.pdf>
32. effects of high-pressure homogenization on the physicochemical,
<https://journal.pan.olsztyn.pl/pdf-98679-30492?filename=30492.pdf>
33. Integrated Membrane Operations: In the Food Production,
<https://dokumen.pub/integrated-membrane-operations-in-the-food-production-9783110285666-9783110284676.html>
34. (PDF) Microstructure of Cheese Products - ResearchGate,
https://www.researchgate.net/publication/326600936_Microstructure_of_Cheese_Products
35. (PDF) Microstructure of Dairy Products - ResearchGate,
https://www.researchgate.net/publication/326060152_Microstructure_of_Dairy_Products
36. Microstructure - Department of Chemical Engineering,
<https://chemical.eng.unimelb.edu.au/dairy/research/microstructure>
37. Microstructure and Texture of Khoa - USU Digital Commons,
https://digitalcommons.usu.edu/context/foodmicrostructure/article/1275/viewcontent/Microstructure_and_Texture_of_Khoa.pdf
38. The effect of milk processing on the microstructure of the milk fat,
<https://pubmed.ncbi.nlm.nih.gov/20492286/>
39. Homogenizers | Dairy Processing Handbook - Tetra Pak,
<http://dairyprocessinghandbook.tetrapak.com/chapter/homogenizers>
40. A New Control Strategy for High-Pressure Homogenization to ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9412542/>
41. Comparative analysis of technological efficiency of various methods,
https://www.researchgate.net/publication/395183626_Comparative_analysis_of_technological_efficiency_of_various_methods_of_formulating_cream_mixtures
42. Rheological Changes During Slow Acid Induced Gelation of Milk by,
https://www.researchgate.net/publication/230001811_Rheological_Changes_During_Slow_Acid_Induced_Gelation_of_Milk_by_D-glucono--lactone
43. Rheological and Viscoelastic Analysis of Hybrid Formulations for,
<https://www.mdpi.com/1999-4923/15/10/2392>
44. Hershel-Bulkley Parameters - Hydrologic Engineering Center,
<https://www.hec.usace.army.mil/confluence/rasdocs/rasmuddebris/non-newtonian-user-s-manual/user-inputs-and-model-parameters/hershel-bulkley-parameters>
45. Viscoelasticity and yield behavior of loess mudflows - Frontiers,
<https://www.frontiersin.org/journals/earth-science/articles/10.3389/feart.2026.1767084/full>
46. Herschel-Bulkley Model - Documentation,

- <https://docs.aft.com/impulse/Herschel-Bulkley-Model.htm>
47. Sensory Texture and Mastication Physics of Multi-Phase Meat,
<https://www.mdpi.com/2076-3417/12/21/11076>
 48. Protein-based Fat Replacers – A Review of Recent Advances,
https://www.academia.edu/90878599/Protein_based_Fat_Replacers_A_Review_of_Recent_Advances
 49. Effects of Ingredients on the Functionality of Fat-free Process,
https://www.researchgate.net/publication/228481314_Effects_of_Ingredients_on_the_Functionality_of_Fat-free_Process_Cheese_Spreads
 50. The effect of κ - and λ -carrageenan concentrations on the viscoelastic,
https://www.researchgate.net/publication/350987416_The_effect_of_k-_and_l-carrageenan_concentrations_on_the_viscoelastic_and_sensory_properties_of_cream_desserts_during_storage
 51. Reaction Kinetics of the Denaturation of Whey Proteins in Milk,
https://www.researchgate.net/publication/229743737_Reaction_Kinetics_of_the_Denaturation_of_Whey_Proteins_in_Milk
 52. Heat-induced interactions of β -lactoglobulin and α -lactalbumin with,
https://www.researchgate.net/publication/223762985_Heat-induced_interactions_of_b-lactoglobulin_and_a-lactalbumin_with_the_casein_micelle_in_pH-adjusted skim_milk
 53. Milk Processing Affects Structure, Bioavailability and ... - MDPI,
<https://www.mdpi.com/2304-8158/9/7/874>
 54. Reaction kinetic pathway of reversible and irreversible thermal,
<https://lait.dairy-journal.org/articles/lait/pdf/2007/04/I07s4505.pdf>
 55. (PDF) Heat-induced interactions of β -lactoglobulin A and κ -casein B,
https://www.researchgate.net/publication/10874109_Heat-induced_interactions_of_b-lactoglobulin_A_and_k-casein_B_in_a_model_system
 56. Influence of calcium on β -lactoglobulin denaturation kinetics - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/22118070/>
 57. Use of thermal-treated or high-pressure treated liquid micellar,
<https://academica-e.unavarra.es/bitstreams/0e4efdbd-0d1c-4234-8ae6-85977b3a53bb/download>
 58. The Effect of Processing on Some Microstructural Characteristics of,
<https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=1357&context=foodmicrostructure>
 59. Flux Decline in Ultrafiltration Processes - <https://ris.utwente.nl>,
<https://ris.utwente.nl/ws/files/6486241/flux.pdf>
 60. Comprehensive Guide to Ultrafiltration Membrane Fouling - Globaluf,
<https://www.globaluf.com/ultrafiltration-uf-membrane-fouling-and-cleaning-methods/>
 61. ROTAPRO SCRAPED SURFACE HEAT EXCHANGER,
https://jhstaal.dk/wp-content/uploads/2019/06/JH-Steelindustri-AS_Brochure_Rotapro_UK.pdf
 62. Scraped Surface Heat Exchangers | Efficient Heating & Cooling,
<https://lgnngroup.co/scraped-surface-heat-exchangers/>

63. (PDF) Heat transfer analysis of scraped surface heat exchanger, https://www.researchgate.net/publication/319500954_Heat_transfer_analysis_of_scraped_surface_heat_exchanger
64. Kombinator 250 | Scraped Surface Heat Exchanger for Margarine, <https://www.spxflow.com/gerstenberg-schroder/products/kombinator-250/>
65. Hercules - Scraped surface heat exchangers - Cleanroom Processes, <https://cleanroom-processes.de/en/product/hercules-schabewarmetauscher/>
66. OPTIMIZATION OF CRUDE PAPAYA (Carica papaya) PROTEASE, <http://202.45.146.37:8080/jspui/bitstream/123456789/131/2/Nabindra%20Thesis.pdf>
67. The mechanism and properties of acid-coagulated milk gels, https://www.researchgate.net/publication/26472637_The_mechanism_and_properties_of_acid-coagulated_milk_gels
68. (PDF) Application of Membrane Separation Technology to Cheese, https://www.researchgate.net/publication/316900215_Application_of_Membrane_Separation_Technology_to_Cheese_Production
69. The Effect of Precipitation pH on Protein Recovery Yield and ... - MDPI, <https://www.mdpi.com/1420-3049/27/9/2957>
70. Cheese problems solved - E-Book's, <https://thunderbooks.files.wordpress.com/2009/05/cheese-problems-solved-woodhead-2007.pdf>
71. Gelation of the complex between kappa-casein and beta-lactoglobulin, <https://pubmed.ncbi.nlm.nih.gov/6427434/>
72. Acid-base buffering curves of milks: (a) titration of milk from initial, https://www.researchgate.net/figure/Acid-base-buffering-curves-of-milks-a-titration-of-milk-from-initial-pH-to-pH-30-with_fig4_51750085
73. Buffering capacity of dairy products | Request PDF - ResearchGate, https://www.researchgate.net/publication/223457883_Buffering_capacity_of_dairy_products
74. Acid and rennet-induced coagulation behavior of casein micelles, https://www.researchgate.net/publication/332273141_Acid_and_rennet-induced_coagulation_behavior_of_casein_micelles_with_modified_structure
75. The Four Broad Categories of Cheese | The Beverage People, <https://www.thebeveragepeople.com/how-to/cheese/articles/four-categories-cheese.html>
76. Mascarpone turns out acidic : r/cheesemaking - Reddit, https://www.reddit.com/r/cheesemaking/comments/1uugtjk/mascarpone_turns_out_acidic/
77. CHEESEMAKING 101 - Make Cheese Inc., <https://www.makecheese.ca/pages/cheesemaking-101>
78. Sensory perception of different acidulants in flavoured sports drinks, <https://repository.up.ac.za/bitstreams/d749efa0-8c34-45c6-97ad-f26dbddd7be8/download>
79. Mind the buffering capacity of citric acid, <https://www.fgsc.net/fgn42/nozawa.html>
80. Citric Acid: A Multifunctional Pharmaceutical Excipient - PMC - NIH,

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC9148065/>
81. Citrate Buffer Preparation Guide | PDF | Buffer Solution | Ph - Scribd,
<https://www.scribd.com/document/597015806/Citrate-Buffer-recipe>
 82. Solubility, acidifying power and sensory properties of fumaric acid in,
<https://oeno-one.eu/article/view/5455>
 83. The Acid Index: Every Culinary Acid Ranked by Science - Littoralicious,
<https://littoralicious.com/articles/the-acid-index>
 84. TASTER RESPONSES TO ACIDITY IN WHITE WINES,
<https://repositorio.ulisboa.pt/bitstream/10400.5/13344/1/Francesco%20-%20TASTER%20RESPONSES%20TO%20ACIDITY%20Final.pdf>
 85. Decoding the Flavor Profiles of Citric, Malic, Lactic, and Tartaric Acid,
<https://www.foodadditivesasia.com/en/market-insights/applications-and-buyers/decoding-flavor-profiles-food-acidulants>
 86. An optimized organic acid human sensory sourness analysis method,
<https://pubmed.ncbi.nlm.nih.gov/33811349/>
 87. Sensory sweetness and sourness interactive response of sucrose,
<https://d-nb.info/126930321X/34>
 88. "Generic" Acids and their effects : r/vegancheesemaking - Reddit,
https://www.reddit.com/r/vegancheesemaking/comments/kl4dkj/generic_acids_and_their_effects/
 89. Chemical Determinants of Sourness and Related Attributes - Blacklight,
<https://etda.libraries.psu.edu/catalog/31029aed5493>
 90. (PDF) Mascarpone cheese from sheep's milk - a new option for the,
https://www.researchgate.net/publication/345771782_Mascarpone_cheese_from_sheep's_milk_-_a_new_option_for_the_consumer
 91. Cake formation and the decreased performance of whey ultrafiltration,
<https://www.redalyc.org/journal/3032/303258119001/html/>
 92. New Insights into the Fouling of a Membrane during the ... - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10054249/>
 93. Membrane surface properties and their effects on real waste oil-in,
<https://watersa.net/article/view/6733>
 94. Fouling and Chemical Cleaning Strategies for Submerged ... - MDPI,
<https://www.mdpi.com/2077-0375/14/12/251>