

Comprehensive Analysis of Vegetable Lacto-Fermentation: Microbial Succession, Osmotic Dynamics, and Biochemical Mechanisms

Introduction to Lactic Acid Fermentation in Vegetable Matrices

The preservation of vegetable matter via lactic acid fermentation represents one of the oldest, most biochemically complex, and ecologically elegant food processing methodologies in human history. At its core, lacto-fermentation is a microbially driven biotransformation wherein indigenous epiphytic lactic acid bacteria (LAB) metabolize plant carbohydrates into organic acids—predominantly lactic and acetic acids¹. This process rapidly lowers the pH of the food matrix, effectively eliminating pathogenic and putrefactive microorganisms while preserving the structural integrity and enhancing the nutritional profile of the plant tissue⁴.

The success of spontaneous vegetable fermentations, such as those yielding sauerkraut (fermented cabbage) and kimchi (traditional Korean fermented vegetables), relies entirely on the precise manipulation of environmental parameters to dictate microbial succession¹. Chief among these parameters is the concentration of sodium chloride (NaCl). The application of salt initiates a cascade of physical, chemical, and biological events that select for highly specialized, halotolerant, and aciduric LAB, while imposing insurmountable osmotic and acidic stress on spoilage organisms⁵. The metabolic exudation of lactic and acetic acids by the LAB consortium fundamentally restructures the thermodynamic and biochemical equilibrium of the brine, transforming a highly perishable agricultural commodity into a stable, functional food rich in bioactive metabolites, bacteriocins, and structural biopolymers¹⁰.

The Physicochemical Foundation of Sodium Chloride (NaCl) Feedstocks

Before examining the biological interactions within the fermentation vessel, it is necessary to understand the physicochemical nature of the primary selective agent: sodium chloride. The origin, purity, and trace mineral composition of the salt heavily dictate the osmotic dynamics and structural outcomes of the fermented vegetable matrix¹³.

Salt Origins, Fractional Crystallization, and Impurities

Industrial and artisanal salt production generally falls into two categories: the solar evaporation of seawater (yielding sea salt) and the solution mining or dry mining of subterranean halite deposits (yielding rock or vacuum pan salt)¹³. The extraction of high-purity sodium chloride via solar evaporation involves a sequential fractional crystallization process monitored historically

by the Baumé scale⁶. As seawater concentrates in shallow ponds, calcium carbonate precipitates first, followed by calcium sulfate (gypsum) in the sulfate stage¹³. Only when the brine reaches a highly supersaturated state (between 25.0 and 29.5 degrees Baumé) does sodium chloride crystallize as halite¹³.

The residual liquor, known as bittern, contains high concentrations of magnesium, potassium, and sulfate ions⁷. Because unrefined sea salts often retain fractions of these trace alkaline earth metals, their use in fermentation introduces calcium and magnesium ions into the brine. As will be discussed regarding pectin cross-linking, these divalent cations play a critical role in preserving the structural crispness of the vegetable tissue¹⁶. Conversely, highly refined vacuum pan salts—which undergo rigorous chemical purification using the lime-soda process to precipitate competing multivalent ions—lack these trace minerals¹³.

Furthermore, modern environmental analyses indicate that minimally processed, unrefined sea salt extracted from coastal solar pans contains significantly higher concentrations of microplastic particles compared to vacuum salt or mined rock salt¹⁸. Subterranean brine deposits predate synthetic polymers, and the closed-loop thermal crystallization (often using Mechanical Vapor Recompression) effectively eliminates microplastic contamination¹³. Consequently, the selection of salt directly impacts both the trace mineral availability and the microscopic purity of the fermentation brine.

Anti-Caking Agents and Fermentation Viability

Because sodium chloride is highly hygroscopic, commercial table salts are frequently treated with anti-caking agents to prevent agglomeration caused by solid crystalline bridges forming between particles during moisture evaporation¹³. The most universally applied agent is Yellow Prussiate of Soda (YPS), or sodium ferrocyanide decahydrate¹³. YPS acts as a microscopic crystal habit modifier, adsorbing onto the faces of nascent salt crystals to preserve flowability¹³. While regulatory authorities approve YPS at trace levels (10 to 250 ppm) due to its high stability and lack of toxicity, artisanal fermentation practitioners often advocate for unrefined or additive-free salts (such as Kosher or Himalayan salts) to avoid potential, albeit poorly documented, interferences with sensitive microbial ecosystems¹³.

Plasmolysis and the Establishment of the Brine Matrix

The addition of salt to raw vegetables is the fundamental trigger for lacto-fermentation. A specific salt concentration of 2.0% to 3.0% by mass or volume is universally recognized as the optimal parameter for initiating a controlled vegetable fermentation⁴.

Methods of Salting and Mathematical Calibration

There are two primary techniques for introducing salt to the vegetable matrix: dry salting and wet brining. Dry salting involves massaging solid sodium chloride directly into shredded vegetables (such as cabbage for sauerkraut). The hypertonic environment forces cellular plasmolysis, drawing water, dissolved soluble carbohydrates, amino acids, and minerals out of the plant cells to form a self-generated, nutrient-rich exudate⁵. In wet brining, whole or thickly chopped vegetables (such as cucumbers or carrots) are submerged in an externally prepared saline solution⁵.

Calculating the precise 2.0% to 3.0% concentration requires accounting for the intrinsic water content of the vegetables, which are typically 93% to 98% water¹⁵. For maximum precision, the percentage must be calculated based on the combined weight of the produce and any added water⁵. For example, to achieve a 2.5% salinity, an artisan would weigh the produce and water (e.g., 100 grams total), multiply by 0.025, and add exactly 2.5 grams of salt¹⁵. This rigorous mass-based mathematical approach ensures consistency, regardless of the crystalline density of the salt chosen (e.g., dense Himalayan salt versus low-density, hollow Alberger process flake salt)¹³.

Vegetable Type	Salting Methodology	Optimum Salinity (%)	Rationale and Matrix Adjustments
Shredded Cabbage (Sauerkraut)	Dry Salting	2.0% – 2.5%	Relies on high surface area for rapid plasmolysis; 2% yields optimal heterofermentative initiation ⁵ .
Napa Cabbage (Kimchi)	Dry or Wet Brining	2.0% – 3.0%	Complex matrix; rapid pH drop required to stabilize diverse auxiliary ingredients (e.g., alliums, radish) ¹ .
Cucumbers / Root Vegetables	Liquid Brine Submergence	3.0% – 5.0%	High initial water content dilutes the external brine rapidly; requires higher starting salinity to prevent pectinolytic softening ⁶ .
Hot Peppers (Mash or Whole)	Dry or Wet Brining	3.0% – 5.0%	High propensity for mold proliferation requires a more aggressive hypertonic barrier ⁶ .

Thermodynamic Water Activity and Anaerobiosis

The introduction of 2-3% salt significantly depresses the thermodynamic activity of water within the matrix. While standard thermodynamic models (like Debye-Hückel) apply to dilute solutions, the ionic strength of hypersaline environments necessitates complex geochemical modeling, such as the Pitzer equations, to calculate the mean ionic activity coefficients and specific ion-ion interactions¹³. By depressing water activity, the brine immediately limits the availability of unbound water required for the metabolic processes of competitive putrefactive microorganisms⁸.

Furthermore, once the vegetables are submerged beneath this exudate or brine, an obligate anaerobic environment is established⁵. This oxygen deprivation acts as a secondary selective pressure, rapidly inhibiting strictly aerobic spoilage bacteria (such as *Pseudomonas* species) and highly oxidative molds⁴. LAB, conversely, are aerotolerant anaerobes or facultative anaerobes; they lack a functional electron transport chain for oxidative phosphorylation but possess unique enzymatic adaptations allowing them to dominate in oxygen-depleted matrices³³.

The Plasmolysis of Pathogens and Enterobacteriaceae

At salt concentrations below 1.5%, the osmotic pressure is insufficient to suppress competitive microbiota. In hypotonic conditions, enteric bacteria (e.g., *Escherichia coli*), pectinolytic spoilage organisms, and undesirable yeasts outcompete the LAB, leading to tissue maceration and off-flavors⁴. The 2.0–3.0% optimum is a precise biological compromise: it is high enough to induce lethal osmotic shock in Gram-negative pathogens, yet low enough to permit the rapid proliferation of halotolerant LAB⁴.

Gram-negative bacteria, possessing a thinner peptidoglycan layer enclosed between an inner and outer membrane, are particularly susceptible to this shock³⁵. The hypertonic brine forces an immediate, massive efflux of water from their cytoplasm, causing a sudden drop in turgor pressure³⁵. The cell membrane detaches from the cell wall, cellular volume collapses, and metabolic processes cease in a phenomenon known as plasmolysis³⁵.

During the initial 24 to 48 hours of spontaneous fermentation, a transient population of *Enterobacteriaceae* often attempts to proliferate before the pH drops⁷. This brief activity is responsible for a characteristic "nitrite peak" observed in vegetable fermentations, as these organisms reduce naturally occurring plant nitrates into nitrites³⁹. However, the combination of osmotic stress and the onset of acid production by early-stage LAB rapidly triggers the collapse of the *Enterobacteriaceae* populations, subsequently allowing the nitrite peak to be degraded and ensuring the safety of the final product³⁹.

Mechanisms of Osmotic Tolerance in Lactic Acid Bacteria

The ability of LAB to thrive in a hypertonic brine while pathogenic bacteria undergo lethal plasmolysis is governed by sophisticated molecular defense mechanisms, primarily the active accumulation of osmoprotectants.

Turgor Pressure Maintenance and the Gram-Positive Advantage

Bacterial cell viability depends on maintaining positive turgor pressure—an outward hydrostatic pressure exerted by the cytoplasm against the cell membrane and cell wall. LAB, being Gram-positive, possess a thick, highly cross-linked peptidoglycan cell wall that provides a robust mechanical scaffold against environmental stress³⁵. While this structural advantage offers baseline resilience against membrane rupture, the primary biochemical defense against hyperosmotic shock is the accumulation of compatible solutes.

Compatible Solutes and Glycine Betaine Transport

To counteract the hypertonic external environment without interfering with internal enzymatic functions, LAB actively transport or synthesize "compatible solutes" (osmoprotectants)³¹. These are highly soluble, low-molecular-weight organic compounds that accumulate to molar concentrations within the cytoplasm, restoring osmotic equilibrium and preventing the lethal influx of sodium ions (Na^+)³¹.

Glycine betaine (N,N,N-trimethylglycine) is the most critical osmoprotectant for LAB in vegetable fermentations²⁹. LAB such as *Lactiplantibacillus plantarum* and *Lactococcus lactis* generally cannot synthesize glycine betaine *de novo*; instead, they rely on high-affinity membrane transport systems to scavenge it from the surrounding plant matrix³⁷.

In *L. plantarum*, the uptake of quaternary ammonium compounds like glycine betaine, carnitine, and choline is mediated by osmoregulated secondary transport proteins and ATP-binding cassette (ABC) transporters, such as the QacT system³⁰. When the external osmolality increases due to NaCl addition, these transport systems are rapidly activated⁴⁵. This mechanism shares evolutionary parallels with the highly studied OpuA, OpuB, and OpuC ABC-type transporters found in *Bacillus subtilis*, which similarly import choline for subsequent oxidation into glycine betaine via the GbsB and GbsA enzymes³⁰.

The master regulatory protein CcpA (Catabolite Control Protein A) plays a vital role in integrating these responses in LAB. Transcriptomic studies reveal that under salt stress, CcpA regulates the uptake of glycine betaine, alters fatty acid composition to maintain membrane lipid homeostasis, and modulates ion transport⁴⁴. The accumulation of glycine betaine not only retains intracellular water but also stabilizes the native state of cellular proteins and lipids against denaturation by salt³¹. Consequently, osmoprotectant-loaded LAB can maintain their

internal Na^+ concentration at a manageable 200–300 nmol/mg (dry weight), while

unprotected bacteria suffer massive Na^+ influx (exceeding 2,000 nmol/mg), leading to rapid cell death³⁷.

Microbial Succession: The Dynamics of Sauerkraut and Kimchi

Vegetable fermentation is not a static microbiological event; it is a dynamic, multi-stage ecological succession. As the environmental parameters—pH, oxygen availability, and nutrient profile—shift over time, one dominant bacterial population yields to another. The successional

pathways of sauerkraut and kimchi illustrate how different matrix compositions lead to divergent microbiological outcomes¹. The contemporary resurgence of artisanal fermentation workshops and community laboratories (e.g., Edible Alchemy in Berlin) specifically capitalizes on these diverse pathways to educate consumers on microbiome health, demonstrating how wild versus cultured starter techniques can radically alter the final organoleptic profile²¹.

Sauerkraut: A Two-Stage, Homofermentative Dominated Succession

Sauerkraut represents a classic, prolonged microbial succession dominated fundamentally by two key species, resulting in a homofermentative, highly acidic final product¹.

1. **Early Fermentation (Days 1–5):** The succession is initiated by *Leuconostoc mesenteroides*, an obligate heterofermentative bacterium¹. *L. mesenteroides* possesses a distinct physiological advantage: it exhibits a faster growth rate at lower temperatures (15°C - 20°C) and higher salt concentrations than other indigenous LAB³². Crucially, as a heterofermenter, it metabolizes hexose sugars via the phosphoketolase pathway, producing equimolar amounts of D-lactic acid, acetic acid, ethanol, and carbon dioxide (CO_2)¹. The vigorous production of CO_2 by *L. mesenteroides* purges residual oxygen from the brine, solidifying the strict anaerobic conditions required for subsequent phases¹⁰. Additionally, it secretes mannitol and specific flavor precursors that contribute to the sensory profile¹. During this phase, the pH drops rapidly from roughly 6.5 to approximately 4.5¹.
2. **Mid-to-Late Fermentation (Days 7–30+):** As the pH falls below 4.5, *L. mesenteroides* becomes inhibited by its own acidic byproducts¹. The ecological niche is then rapidly occupied by more acid-tolerant, homofermentative lactobacilli, primarily *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*), alongside *Lactiplantibacillus pentosus* and *Levilactobacillus brevis*¹. *L. plantarum* utilizes the highly efficient Embden-Meyerhof-Parnas (EMP) pathway to convert remaining sugars almost exclusively into lactic acid¹. This phase drives the pH down to a highly stable terminal level of 3.2 to 3.8¹. The extended maturation period allows *L. plantarum* to generate complex secondary metabolites, resulting in the sharp, clean, and highly acidic flavor profile characteristic of mature sauerkraut¹.

Kimchi: Rapid, Heterofermentative Complexity

Kimchi fermentation diverges significantly from sauerkraut due to its highly complex ingredient matrix, which includes high-sugar vegetables (daikon radish), alliums (garlic, green onions), ginger, chili powder (gochugaru), and often animal-derived amino acids from fish sauce or salted shrimp¹.

Unlike sauerkraut, which trends heavily toward a homofermentative climax, kimchi is characterized by a persistent and dominant heterofermentative profile¹. The abundance of soluble sugars from the diverse ingredients fuels an explosive early growth of *Leuconostoc* species (e.g., *L. mesenteroides*, *L. citreum*, *L. holzapfelii*) and *Weissella* species (e.g., *W. koreensis*, *W. soli*)¹.

This explosive heterofermentative activity causes the pH of kimchi to plummet much faster than

sauerkraut, dropping to roughly 4.5 in just 3 to 5 days¹. The dominance of *Leuconostoc* and *Weissella* produces a complex metabolite mix featuring high levels of CO_2 (giving kimchi its characteristic effervescence), mannitol (contributing a cooling sweetness), acetic acid, and gamma-aminobutyric acid (GABA)¹. As the pH stabilizes around 3.9 to 4.3 (the optimal edible range), the microbial community shifts toward *Latilactobacillus sakei* and, to a lesser extent, *L. plantarum*¹. *L. sakei* is particularly well-adapted to the cold temperatures often used in traditional kimchi fermentation and possesses unique metabolic pathways to cope with oxygen changes, heme uptake, and the complex spice matrix¹. Because the fermentation kinetics are highly accelerated and effectively complete by day 20, kimchi retains a highly complex, multi-metabolite profile that differs markedly from the pure lactic acid dominance of aged sauerkraut¹.

Microbiological & Chemical Parameter	Sauerkraut (Cabbage-based)	Kimchi (Complex Matrix)
Metabolic Pathway Dominance	Shifts to Homofermentative at maturity	Persistent Heterofermentative
Primary Initiating Genera	<i>Leuconostoc mesenteroides</i>	<i>Leuconostoc</i> spp., <i>Weissella</i> spp.
Climax Dominance	<i>Lactiplantibacillus plantarum</i>	<i>Latilactobacillus sakei</i> , <i>Leuconostoc</i> spp.
Kinetics of Acidification	Slow (weeks to achieve pH < 4.0)	Rapid (days to achieve pH ~ 4.2)
Terminal pH Range	3.2 – 3.8	3.4 – 4.5
Key Metabolite Profile	Lactic acid (overwhelmingly dominant)	Lactic acid, Acetic acid, Mannitol, GABA, CO_2

The Biochemistry of Pathogen Inhibition

The safety and preservation of lacto-fermented vegetables rely on multiple overlapping biochemical mechanisms, primarily centered on organic acid toxicity and the targeted production of bacteriocins.

The Proton Effect: Membrane Permeability of Weak Organic Acids

The primary mechanism by which lacto-fermentation inhibits putrefying and pathogenic bacteria is the accumulation of weak organic acids—namely lactic acid and acetic acid. It is a

fundamental misconception that mere hydrogen ion (H^+) concentration (measured as ambient pH) is solely responsible for bacterial death. While strong mineral acids like hydrochloric acid (HCl) fully dissociate in aqueous solutions, weak organic acids only partially dissociate, governed by their specific acid dissociation constant (pK_a)⁵⁸.

The pK_a of lactic acid is 3.86, and the pK_a of acetic acid is 4.76²⁰. According to the Henderson-Hasselbalch equation, when the environmental pH falls near or below the pK_a of a specific acid, a significant proportion of the acid molecules exist in their undissociated (protonated) state⁵⁸.

The uncharged, undissociated forms of lactic and acetic acid are lipophilic, allowing them to freely diffuse across the lipid bilayer of pathogenic bacterial cell membranes⁶¹. Once inside the pathogen's cytoplasm, where the internal pH is maintained near neutrality (pH ~7.0), the acids immediately encounter an environment well above their pK_a . Consequently, they rapidly dissociate, releasing protons (H^+) and acid anions (lactate/acetate) directly into the cell interior⁵⁸.

This internal dissociation triggers a lethal biological crisis for the pathogen via the "proton effect":

1. **Collapse of the Proton Motive Force (PMF):** The massive influx of internal protons destroys the transmembrane electrochemical gradient (the ΔpH component of the PMF), paralyzing the pathogen's ability to synthesize ATP or transport nutrients across the membrane²⁰.
2. **ATP Depletion:** The pathogen desperately attempts to pump the excess protons back out of the cell using its own membrane ATPases. This futile cycle rapidly exhausts cellular ATP reserves, leading to metabolic exhaustion and cell death⁵⁸.
3. **Anion Toxicity:** The concurrent accumulation of intracellular lactate and acetate anions further disrupts metabolic enzymes and destroys internal osmotic balance⁵⁸.

Because acetic acid possesses a higher pK_a (4.76) than lactic acid (3.86), it maintains a substantially higher proportion of undissociated molecules at typical fermentation pH ranges (e.g., pH 4.5)⁶⁰. Therefore, the acetic acid produced by heterofermentative bacteria like *L. mesenteroides* acts as an exceptionally potent antimicrobial agent in the early stages of fermentation, providing a synergistic preservation effect alongside the higher volumes of lactic acid⁶².

Bacteriocins and Quorum Sensing

In addition to organic acids, LAB synthesize and secrete ribosomally synthesized antimicrobial peptides known as bacteriocins. These proteinaceous toxins provide LAB with a significant

competitive advantage in the crowded vegetable matrix by targeting and neutralizing closely related Gram-positive competitors and, in specific configurations, Gram-negative pathogens⁴³. Bacteriocins are broadly classified into distinct subgroups, with Class I (lantibiotics, e.g., Nisin, which feature post-translational modifications like lanthionine) and Class II (small, heat-stable, non-lanthionine peptides) being the most prominent in LAB⁶⁴. In vegetable fermentations, Class IIa (pediocin-like, featuring a conserved YGNGV motif) and Class IIb (two-peptide) bacteriocins play critical ecological roles⁶⁵. For example, *L. plantarum* produces plantaricin, a highly potent Class IIb bacteriocin that targets the cell envelope of competing microbes⁴³.

The mechanism of action for Class IIa bacteriocins is highly specific. They recognize and bind to the membrane-embedded portion of the mannose phosphotransferase system (Man-PTS) IIC and IID subunits located on the plasma membrane of target cells⁶⁵. Upon binding, the bacteriocins undergo a conformational change, inserting themselves into the lipid bilayer to form

pores. This pore formation causes a rapid, uncontrolled efflux of intracellular ions (K^+ , Mg^{2+}) and amino acids, obliterating the target cell's proton motive force and resulting in immediate cell death⁶³. While primarily targeting Gram-positive spoilage organisms (such as the dangerous foodborne pathogen *Listeria monocytogenes*), certain plantaricins and Class IIa variants have demonstrated broad-spectrum activity capable of permeabilizing the outer membrane of Gram-negative pathogens such as *E. coli* and *Salmonella* species⁴³.

Crucially, the biosynthesis of bacteriocins is not a constant, constitutive process; it is tightly regulated by Quorum Sensing (QS)⁶⁴. LAB continually secrete low molecular weight signaling molecules called autoinducer peptides (AIPs). When the population density of the LAB reaches a critical threshold, the extracellular concentration of AIPs triggers a two-component regulatory system (involving a histidine kinase receptor and a response regulator)⁶⁴. This phosphorylation cascade initiates the transcription of bacteriocin structural genes. By delaying bacteriocin production until they have achieved sufficient biomass, LAB efficiently utilize their metabolic energy to mount a coordinated, overwhelming chemical attack against competitors at the precise moment it will yield ecological dominance⁶⁴.

Endogenous Acid Tolerance Mechanisms in LAB

The aggressive production of organic acids fundamentally alters the vegetable matrix. While this acid-shock kills pathogens, it demands an evolutionary solution: how do LAB survive in the highly acidic, low-pH environment (frequently falling below pH 4.0) that they themselves create? LAB rely on highly specialized, energy-dependent biochemical pathways to regulate their own intracellular pH, ensuring homeostasis despite the hostile extracellular gradient.

The F_1F_0 -ATPase Proton Pump

The primary, universal defense mechanism against intracellular acidification in LAB is the

F_1F_0 -ATPase enzyme complex⁶⁸. Unlike pathogenic bacteria that catastrophically exhaust their ATP trying to pump out protons in acidic environments, LAB have evolved highly efficient,

acid-inducible F_1F_0 -ATPase systems specifically tuned for this purpose⁶⁸. Located in the cytoplasmic membrane, this multi-subunit complex actively extrudes protons (H^+) from the cytoplasm to the extracellular environment, a mechanism powered directly by the hydrolysis of ATP⁶⁸. Under severe acid stress, transcriptomic analyses of *L. plantarum* reveal a significant, rapid upregulation of the *atp* operon (which encodes the structural subunits of the F_1F_0 -ATPase)⁶⁸. By continuously pumping out the excess protons delivered by the influx of their own organic acids, LAB maintain an intracellular pH close to neutrality (pH 6.0–7.0). This targeted proton extrusion preserves the functionality of their critical metabolic enzymes and nucleic acids, allowing them to remain metabolically active while the external environment remains aggressively acidic⁶⁸.

The Glutamate Decarboxylase (GAD) System and GABA Production

A vital, specialized secondary acid-resistance mechanism utilized heavily by *Latilactobacillus sakei*, *Levilactobacillus brevis*, and *L. plantarum* is the Glutamate Decarboxylase (GAD) system⁵⁷. This pathway provides a highly efficient method for consuming intracellular protons while simultaneously generating a valuable, bioactive metabolic byproduct.

The GAD system relies fundamentally on the enzyme glutamate decarboxylase (encoded by the *gadA* or *gadB* genes) and a specific membrane-bound antiporter protein (*gadC*)⁵⁷. When the intracellular environment becomes dangerously acidic, the LAB import extracellular L-glutamate—which is naturally abundant in plant tissues, and particularly concentrated in auxiliary kimchi ingredients like fish sauce or soy sauce—via the GadC antiporter⁷¹.

Inside the cell, the GAD enzyme catalyzes the irreversible α -decarboxylation of the L-glutamate molecule to form gamma-aminobutyric acid (GABA). The biochemical elegance of this specific reaction lies in its stoichiometry: the decarboxylation of a single glutamate molecule consumes one free intracellular proton (H^+) and releases one molecule of carbon dioxide (CO_2)⁵⁷. The newly synthesized GABA is then exported back out of the cell via the GadC antiporter in a continuous exchange for a fresh molecule of glutamate⁷¹.

This continuous biochemical cycle acts as a powerful intracellular proton sink, actively raising the internal pH without expending precious cellular ATP⁵⁷. The optimal parameters for this system vary slightly by species; for example, *L. sakei* exhibits maximum GAD activity at 55°C and pH 5.0, while *L. plantarum* optimizes at 40°C and pH 4.5⁵⁷. The massive accumulation of GABA in the extracellular brine is particularly notable in kimchi, functionally transforming the fermented vegetable into a nutraceutical food rich in this non-protein amino acid, which acts as the primary inhibitory neurotransmitter in the mammalian central nervous system and assists in regulating blood pressure and anxiety¹⁰.

Manganese Accumulation and Iron-Free Oxidative Stress Defense

Although lacto-fermentation ultimately establishes an anaerobic environment, the initial stages of fermentation expose LAB to oxygen. Furthermore, several LAB species retain the potential to

undergo aerobic respiration via specific enzymes³³. This oxygen exposure leads to the inevitable formation of lethal Reactive Oxygen Species (ROS), such as superoxide anions (O_2^-), hydrogen peroxide (H_2O_2), and highly reactive hydroxyl radicals ($OH^\bullet$) generated by enzymes like NADH oxidase (Nox), pyruvate oxidase (Pox), and lactate oxidase (Lox)³³. Most aerobic organisms neutralize ROS using iron-dependent heme enzymes, most notably classic catalase. However, LAB are highly unique in that they typically lack heme and cannot synthesize classic iron-dependent catalases³⁴. To survive oxidative stress without iron, *L. plantarum* and related species rely on the massive, active intracellular accumulation of manganese ions (Mn^{2+})³⁴. Using high-affinity transport systems, LAB import manganese to levels thousands of times higher than those found in standard bacteria³⁴.

Once inside the cytoplasm, Mn^{2+} serves as an essential cofactor for specialized antioxidant enzymes, such as manganese-dependent superoxide dismutase (Mn-SOD) and a non-heme pseudocatalase (Mn-Kat)⁷³. Furthermore, highly concentrated intracellular Mn^{2+} can complex with inorganic phosphate, peptides, and organic acids to directly scavenge and decompose superoxide radicals non-enzymatically³⁴. This unique, iron-free redox defense system allows LAB to dominate the aerotolerant niche during early fermentation without risking the catastrophic oxidative damage that iron-driven Fenton reactions would otherwise cause in the presence of H_2O_2 ⁷⁴.

Biochemical Transformations: Texture, Flavor, and Nutritional Enhancement

The intersection of salt concentration, enzymatic activity, and microbial metabolism ultimately defines the sensory, rheological, and nutritional qualities of the final fermented product.

Enzymatic Modulation of Plant Cell Walls and Texture Preservation

The structural integrity, bite, and crispness of fermented vegetables are heavily dependent on the management of plant cell wall components, specifically the highly complex polysaccharide pectin²⁵. Pectin methylesterase (PME) and polygalacturonase (PG) are endogenous plant enzymes released during the mechanical tissue maceration inherent in preparation²⁵. If left

unchecked, PG rapidly hydrolyzes the α -(1,4)-glycosidic bonds in the polygalacturonic acid backbone of pectin, leading to catastrophic tissue softening and mushiness—a primary cause of commercial product failure¹⁶.

The precise application of salt (2-3%) critically modulates these enzymes. The high ionic strength of the sodium chloride brine alters the electrostatic interactions within the plant cell wall, partially inhibiting polygalacturonase activity¹⁶. Concurrently, the salt facilitates the extraction of calcium and magnesium ions from the plant tissue (or provides them directly if unrefined solar salts are used)¹³. These divalent cations interact with the demethylated pectin

chains (which are the product of continued PME activity) to form robust calcium-pectate cross-links¹⁶. This biochemical bridging reinforces the middle lamella of the cell walls, ensuring the vegetables retain a rigid, crisp texture throughout months of highly acidic storage¹⁶.

Glucosinolate Degradation and the Generation of Bioactive

Isothiocyanates

Vegetables belonging to the *Brassica* genus (including cabbage, daikon radishes, and mustard greens) are characterized by high intracellular concentrations of secondary sulfur-containing metabolites called glucosinolates¹¹. In their intact form within the plant vacuole, glucosinolates are biologically inert. However, during the mechanical disruption of salting and packing, endogenous plant enzymes known as myrosinases are released from specialized cells, coming into contact with the glucosinolates and hydrolyzing them into a suite of bioactive breakdown products²⁷.

The acidic, microbially active environment of fermentation heavily influences this degradation pathway. Rather than forming toxic nitriles, the low pH and specific microbial milieu direct the breakdown of glucosinolates (such as glucobrassicin and sinigrin) into isothiocyanates, sulforaphane, and ascorbigen¹¹. These sulfur-containing volatile compounds are directly responsible for the pungent, characteristic aromas of sauerkraut and kimchi¹¹. Furthermore, isothiocyanates and ascorbigen are highly stable in acidic environments and represent potent bioactive compounds known to induce Phase II detoxification enzymes (such as glutathione S-transferase, GST) in mammalian systems, conferring significant chemopreventive and antioxidant properties to the fermented matrix¹¹.

Control of Biogenic Amines

A critical safety function of successful, rapid lacto-fermentation is the strict suppression of biogenic amines (BAs). BAs, such as histamine, tyramine, and putrescine, are low-molecular-weight organic bases formed by the microbial decarboxylation of free amino acids (e.g., histidine converted to histamine, tyrosine converted to tyramine)⁸². Ingestion of foods with high BA concentrations can lead to severe physiological reactions, including vasodilation, migraines, tachycardia, and localized allergic responses—symptoms clinically identical to scombroid food poisoning typically associated with temperature-abused fish⁸².

The formation of BAs in fermented vegetables is primarily driven by contaminating spoilage organisms or a dangerously slow initial fermentation rate⁸⁴. The strict adherence to a 2-3% salt concentration and the subsequent rapid succession of *L. mesenteroides* directly mitigates this safety risk. By establishing an immediate, aggressive pH drop via organic acid production, the specific decarboxylase enzymes responsible for BA formation in competitive bacteria are either permanently inhibited, or the aminobiogenetic bacteria themselves are eradicated via plasmolysis before significant amine accumulation can occur⁸⁵.

Exopolysaccharides and Dextran Synthesis

In fermentations where the matrix contains sufficient amounts of sucrose, or those specifically formulated with added sugars, *Leuconostoc mesenteroides* and certain *Weissella* species can utilize the extracellular enzyme dextransucrase to synthesize massive extracellular polymers

known as dextrans⁸⁷. Dextranucrase cleaves the disaccharide sucrose, utilizing the high energy from the glycosidic bond to polymerize the glucose monomer into a long, branched

homopolysaccharide chain (predominantly an α -1,6-linked glucan), while releasing the residual fructose into the brine to be metabolized⁴⁹.

The extensive production of dextran significantly increases the rheological viscosity of the brine, sometimes leading to a transient "ropy" or viscous fermentation¹². While excessive ropiness is occasionally viewed as a textural defect in some commercial products, moderate dextran production acts as a highly functional prebiotic, enhancing the mouthfeel of the brine and providing immunomodulatory and cholesterol-lowering benefits to the host upon consumption¹².

Stereochemistry of Lactic Acid Production

The specific strains of LAB dominating the terminal phase of the fermentation dictate the stereochemical orientation of the final lactic acid product. Lactic acid exists as two distinct optical isomers: L-lactate and D-lactate⁹⁰.

Humans possess the enzyme L-lactate dehydrogenase in high quantities, making L-lactate readily metabolizable and safe in high volumes. Conversely, humans metabolize D-lactate much more slowly; high systemic accumulations can theoretically lead to D-lactic acidosis in highly susceptible individuals with short-bowel syndrome⁹². In vegetable fermentations, *Lactiplantibacillus plantarum* possesses both L- and D-lactate dehydrogenases, typically producing a racemic mixture (DL-lactate)⁹⁰. *Leuconostoc* species, however, primarily synthesize D-lactate³². The ultimate ratio of D- to L-lactate in the mature product is therefore a direct biochemical biomarker of the successional timeline; a prolonged heterofermentative phase (as seen in cold-fermented kimchi dominated by *Leuconostoc*) tends to yield higher proportions of D-lactate, while an accelerated homofermentative climax balances the stereochemical profile³².

Synthesized Conclusions

The lacto-fermentation of vegetables represents an exceptionally calibrated, multistage interplay of physical chemistry, microbial ecology, and molecular biology. The precise integration of 2.0% to 3.0% sodium chloride serves as the primary environmental master-switch, dictating the entire thermodynamic and biological trajectory of the system. By inducing rapid cellular plasmolysis and establishing an anaerobic, hypertonic environment, the salt selectively eradicates putrefactive *Enterobacteriaceae* and highly oxidative spoilage fungi, allowing a highly specialized cohort of lactic acid bacteria to thrive and initiate succession.

Organisms such as *Leuconostoc mesenteroides*, *Lactiplantibacillus plantarum*, and *Latilactobacillus sakei* dominate this specific ecological niche by deploying advanced, evolutionary survival mechanisms. These range from the active transport of osmoprotectants like glycine betaine via the QacT system to the iron-independent scavenging of reactive oxygen species via massive intracellular manganese accumulation. As these bacteria orchestrate a complex ecological succession—shifting from the rapid heterofermentative gas production of kimchi to the prolonged homofermentative acidification of sauerkraut—they fundamentally reshape the chemical landscape of the plant matrix. Through the coordinated operation of

ATP-driven F_1F_0 proton pumps and the GABA-producing glutamate decarboxylase (GAD) pathway, they successfully survive the very organic acids they excrete.

These organic acids, operating synergistically with quorum-sensing regulated, pore-forming bacteriocins, collapse the proton motive force of pathogenic competitors, ensuring absolute microbiological safety and the strict suppression of toxic biogenic amines. Simultaneously, enzymatic modulation of pectin preserves the cellular crispness of the plant tissue, while secondary microbial metabolism transforms inert plant glucosinolates into highly bioactive isothiocyanates and synthesizes functional exopolysaccharides like dextran. Ultimately, vegetable lacto-fermentation stands as a flawless natural paradigm of biopreservation, where extreme, carefully calibrated environmental stress is harnessed to synthesize a safe, biochemically complex, and nutritionally enhanced food product.

Works cited

1. Sauerkraut vs Kimchi — A Chemist Compares | SourChad,
<https://sourchad.com/compare/sauerkraut-vs-kimchi>
2. Characterization of microbiota of naturally fermented sauerkraut by,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10082884/>
3. Improved sauerkraut production with probiotic strain Lactobacillus,
<https://pubmed.ncbi.nlm.nih.gov/21535775/>
4. Studying the Influence of Salt Concentrations on Betalain and,
<https://www.mdpi.com/1420-3049/29/20/4803>
5. Salt and Brine For Fermenting Vegetables (Ultimate Guide),
<https://revolutionfermentation.com/en/blogs/fermented-vegetables/salt-brine-fermenting-vegetables/>
6. How Much Salt Should I Use For My Fermentation?,
<https://nwedible.com/how-much-salt-fermentation/>
7. The cutting type of vegetables influences the spontaneous,
<https://peercommunityjournal.org/articles/10.24072/pcjournal.553/>
8. (PDF) Sodium Reduction in Traditional Fermented Vegetables,
https://www.researchgate.net/publication/402695272_Sodium_Reduction_in_Traditional_Fermented_Vegetables_Challenges_Strategies_and_Perspectives
9. Change in pH and acidity during kimchi fermentation at 8 (A), 15 (B),,
https://www.researchgate.net/figure/Change-in-pH-and-acidity-during-kimchi-fermentation-at-8-A-15-B-and-25-o-C-C-for_fig1_348067024
10. A Review of the Health Benefits of Kimchi Functional Compounds,
<https://www.mbl.or.kr/journal/view.html?doi=10.48022/mbl.2310.10011>
11. Changes in glucosinolates and their breakdown products during the,
https://www.researchgate.net/publication/352893036_Changes_in_glucosinolates_and_their_breakdown_products_during_the_fermentation_of_cabbage_and_prolonged_storage_of_sauerkraut_Focus_on_sauerkraut_juice
12. Biotechnological potential and genomic analysis of the Leuconostoc,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12628567/>
13. 68 Sodium Chloride Extraction Methods.pdf

14. A Primer on Lactic-Fermented Vegetables,
<https://homebrewersassociation.org/beer-food/a-primer-on-lactic-fermented-vegetables/>
15. The Perfect Lacto Fermentation Salt Ratio for Fermenting Vegetables,
<https://cultured.guru/blog/the-perfect-lacto-fermentation-salt-ratio-for-fermenting-vegetables>
16. Effects of Salts and Preheating Temperature of Brine on the Texture,
https://www.researchgate.net/publication/227561423_Effects_of_Salts_and_Preheating_Temperature_of_Brine_on_the_Texture_of_Pickled_Cucumbers
17. How Microbiome Composition Correlates with Biochemical ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9430578/>
18. Workshops - Mimi Ferments, <https://mimiferments.com/en/collections/workshop>
19. vegetables workshop 26.04.2026 - Berlin - Mimi Ferments,
<https://mimiferments.com/en/products/gemuse-workshop-am-20-september-2026>
20. Minimal inhibitory concentrations of undissociated lactic, acetic, citric,
<https://www.nizo.com/wp-content/uploads/Wemmenhove-et-al.-2016-Minimal-inhibitory-concentrations-of-undissociated-lactic-acetic-citric-and-propionic-acid-for-Listeria-monocytogenes-under-conditions-relevant-to-cheese.pdf>
21. Veggie Fermentation - Berlin - The Makery,
<https://themaker.de/veggie-fermentation/Berlin>
22. Learn - EDIBLE ALCHEMY, <https://ediblealchemy.co/learn/>
23. Edible Alchemy - Alexis Goertz - Eventbrite,
<https://www.eventbrite.de/o/edible-alchemy-alexis-goertz-7775070161>
24. Edible Alchemy - Umweltbildung Treptow-Köpenick,
<https://umweltbildung-treptnick.berlin/akteurin/edible-alchemy/>
25. Blogs - Bubble Organic,
<https://www.bubbleorganic.net/index.php?route=information/blogger>
26. Edible Alchemy - EDIBLE ALCHEMY, <https://ediblealchemy.co/>
27. Services - BioProfile Testing Laboratories, LLC,
<https://www.bioprofilelabs.com/FateGlucosinoatesSulfidesKimchi.htm>
28. Workshops - EDIBLE ALCHEMY, <https://ediblealchemy.co/workshops/>
29. Potential osmoprotectants for the lactic acid bacteria *Pediococcus*,
<https://www.scilit.com/publications/fe95b9451a1287cc4fa88b9dc86be724>
30. Genetic Control of Osmoadaptive Glycine Betaine Synthesis in,
<https://journals.asm.org/doi/10.1128/jb.06642-11>
31. Bacterial osmoprotectants—a way to survive in saline conditions,
<https://academic.oup.com/femsre/article/doi/10.1093/femsre/fuaf020/8133896>
32. Changes in Microbial Community Composition during Production of,
https://www.jstage.jst.go.jp/article/fstr/20/3/20_693/_html/-char/ja
33. Antioxidant Potential of Probiotics and Postbiotics Derived from,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC13072972/>
34. Regulation and distinct physiological roles of manganese in bacteria,
<https://academic.oup.com/femsre/article/45/6/fuab028/6284802>
35. Bacterial Cell Wall Quality Control during Environmental Stress | mBio,
<https://journals.asm.org/doi/10.1128/mbio.02456-20>

36. Bacteriocins from lactic acid bacteria: purification, properties and, <https://www.scielo.br/j/babt/a/htZvm5MQCwK7BzjLsM7TqVg/?lang=en>
37. Glycine Betaine, Carnitine, and Choline Enhance Salinity ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC91856/>
38. Microbial communities of a variety of 75 homemade fermented, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10757351/>
39. Sodium Reduction in Traditional Fermented Vegetables - MDPI, <https://www.mdpi.com/2311-5637/12/3/162>
40. Sodium Reduction in Traditional Fermented Foods - ACS Publications, <https://pubs.acs.org/doi/10.1021/acs.jafc.1c01687>
41. Production of Pickles by Mixed Culture Fermentation - ResearchGate, https://www.researchgate.net/publication/385416288_Production_of_Pickles_by_Mixed_Culture_Fermentation
42. Leuconostoc mesenteroides AA001: A High-Efficiency Nitrite ... - MDPI, <https://www.mdpi.com/2076-2607/14/2/411>
43. (PDF) Bacteriocin, Plantaricin and Pediocin Biosynthesis in Lactic, https://www.researchgate.net/publication/373538436_Bacteriocin_Plantaricin_and_Pediocin_Biosynthesis_in_Lactic_Acid_Bacteria_Antimicrobial_Mechanism_and_Applications_as_Food_Preservatives_Review
44. The protective role of glycine betaine in Lactobacillus plantarum ST, https://www.researchgate.net/publication/262195965_The_protective_role_of_glycine_betaine_in_Lactobacillus_plantarum_ST-III_against_salt_stress
45. Mechanism of Osmotic Activation of the Quaternary Ammonium, <https://pmc.ncbi.nlm.nih.gov/articles/PMC107610/>
46. Stress responses of Bacillus subtilis to high osmolarity environments, https://www.uni-marburg.de/de/fb17/fachgebiete/mikrobiologie/ag-bremer/publikationen/pdf-zu-reviews/rev_04.pdf/@download/file/rev_04.pdf
47. Kimchi to kombucha: The art of fermentation at Edible Alchemy, <https://www.the-berliner.com/berlin/edible-alchemy-diy-miso-kombucha-and-kimchi/>
48. Improving the bacterial community, flavor, and safety properties of, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11068551/>
49. Development of the Efficient Electroporation Protocol for ... - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12733099/>
50. Microbial and metabolic characterization of organic artisanal ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9606823/>
51. Effects of inoculation with Lactiplantibacillus plantarum PC170 and, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13138060/>
52. Some Important Metabolites Produced by Lactic Acid Bacteria ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8465840/>
53. Microbial succession and metabolite changes during fermentation of, https://www.researchgate.net/publication/236184356_Microbial_succession_and_metabolite_changes_during_fermentation_of_dongchimi_traditional_Korean_watery_kimchi
54. Microbial succession and metabolite changes during long-term,

- <https://pubmed.ncbi.nlm.nih.gov/23550842/>
55. Microbial Succession and Metabolite Changes during Long-Term, https://www.researchgate.net/publication/236101843_Microbial_Succession_and_Metabolite_Changes_during_Long-Term_Storage_of_Kimchi
 56. Study of γ -Amino Butyric Acid (GABA) Production by Lactobacillus, https://www.researchgate.net/publication/386180136_Study_of_g-Amino_Butyric_Acid_GABA_Production_by_Lactobacillus_sakei_B2-16
 57. Contributions of Gamma-Aminobutyric Acid (GABA) Produced ... - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11311711/>
 58. Elucidation of toxicity of organic acids inhibiting growth of, https://www.researchgate.net/publication/287575040_Elucidation_of_toxicity_of_o rganic_acids_inhibiting_growth_of_Escherichia_coli_W
 59. The Acid Response Network of Staphylococcus aureus - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7311314/>
 60. Surface physiological changes induced by lactic acid on pathogens, https://www.researchgate.net/publication/273544517_Surface_physiological_chan ges_induced_by_lactic_acid_on_pathogens_in_consideration_of_pKa_and_pH
 61. Extractive Fermentation of Lactic Acid in Lactic Acid Bacteria, <https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2017.02285/full>
 62. Understanding the Tolerance of Different Strains of Human ... - MDPI, <https://www.mdpi.com/2076-3417/13/1/305>
 63. Bacteriocins of Lactic Acid Bacteria as Potent Antimicrobial Peptides, <https://www.intechopen.com/chapters/74860>
 64. Lactic Acid Bacteria Bacteriocins: Classification, Biosynthesis, <https://pubs.acs.org/jafcau/article/74/7/5859/5086299/Lactic-Acid-Bacteria-Bacteri ocins-Classification>
 65. FliA regulates the antibacterial activity of plantaricin BM-1 against, <https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2025.1606567/full>
 66. advances in classification, synthesis, mode of action, and food, <https://academic.oup.com/jambio/article/135/11/lxae274/7875240>
 67. Bacteriocin production enhancing mechanism of Lactiplantibacillus, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9998909/>
 68. The acid tolerance association with expression of H⁺-ATPase in, https://www.researchgate.net/publication/230053345_The_acid_tolerance_associ ation_with_expression_of_H-ATPase_in_Lactobacillus_casei
 69. Exploring Probiotic Potential: A Comparative Genomics and In Silico, <https://www.biorxiv.org/content/10.1101/2024.05.15.594408v1.full-text>
 70. Survival of Probiotic Lactobacilli in Acidic Environments Is Enhanced, https://www.researchgate.net/publication/7809197_Survival_of_Probiotic_Lactoba cilli_in_Acidic_Environments_Is_Enhanced_in_the_Presence_of_Metabolizable_S ugars
 71. Production of Gamma (γ)-Aminobutyric Acid by Lactic Acid Bacteria, <https://www.mdpi.com/2674-1334/5/3/47>

72. Advances in Biotechnological GABA Production: Exploring Microbial,
<https://www.mdpi.com/1422-0067/27/1/306>
73. Manganese-Based Antioxidant Agents: Understanding Structure,
https://www.researchgate.net/publication/399151072_Manganese-Based_Antioxidant_Agents_Understanding_Structure-Activity_Relationships_in_Biological_Applications
74. Isolation, identification, and evaluation of the antioxidant properties,
<https://journals.plos.org/plosone/article/file?type=printable&id=10.1371/journal.pone.0327225>
75. (PDF) Antioxidant Activity and Metabolomic Characterization of,
https://www.researchgate.net/publication/407130285_Antioxidant_Activity_and_Metabolomic_Characterization_of_Lactiplantibacillus_plantarum_MCS1903_Isolated_from_Naturally_Fermented_Tofu_Whey
76. Antioxidant Activity and Metabolomic Characterization of ... - MDPI,
<https://www.mdpi.com/2076-2607/14/6/1348>
77. Image analysis of kimchi cabbage penetrated with brine and,
https://www.researchgate.net/publication/387200109_Image_analysis_of_kimchi_cabbage_penetrated_with_brine_and_seasoning_using_a_serial_block_face_scanning_electron_microscope_and_energy_dispersive_X-ray_spectroscopy
78. Yogurt Production and Health Benefits | PDF - Scribd,
<https://www.scribd.com/document/630550119/C-3-pdf>
79. Microbiological Spoilage of Fruits and Vegetables - USDA ARS,
<https://www.ars.usda.gov/ARSUserFiles/60701000/Pickle%20Pubs/p363.pdf>
80. Changes in glucosinolates and their breakdown products ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/34243119/>
81. Changes in volatile flavor compounds of Kimchi cabbage (Brassica,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11016023/>
82. A review of biogenic amines in fermented foods - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10830535/>
83. (PDF) A review of biogenic amines in fermented foods - ResearchGate,
https://www.researchgate.net/publication/377583963_A_review_of_biogenic_amin es_in_fermented_foods_Occurrence_and_health_effects
84. Biogenic amines in foods - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6049710/>
85. Technological Factors Affecting Biogenic Amine Content in Foods,
<https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2016.01218/full>
86. The inhibitory effects of plant additives on biogenic amine formation,
<https://www.tandfonline.com/doi/full/10.1080/10408398.2023.2258964>
87. Dextransucrase production by *Leuconostoc mesenteroides*,
https://www.researchgate.net/publication/285765035_Dextransucrase_production_by_Leuconostoc_mesenteroides
88. Production of Dextran from Domestic *Weissella cibaria* Isolated,
https://www.researchgate.net/publication/376548409_Production_of_Dextran_from_Domestic_Weissella_cibaria_Isolated

89. Purification, Characterization and N-terminal Protein Sequencing of,
<https://www.biotech-asia.org/vol18no2/purification-characterization-and-n-terminal-protein-sequencing-of-the-enzyme-dextranucrase-produced-by-leuconostoc-mesenteroides/>
90. Exploring the Potential of Lactic Acid Bacteria Fermentation as a,
<https://www.mdpi.com/2076-3417/15/5/2686>
91. Metabolism Characteristics of Lactic Acid Bacteria and the ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8149962/>
92. Comprehensive Evaluation of Safety, Functional Properties, and,
https://www.jmb.or.kr/journal/download_pdf.php?doi=10.4014/jmb.2605.05037
93. Effect of lactic acid bacteria on D- and L-lactic acid contents of kimchi,
https://www.researchgate.net/publication/299261172_Effect_of_lactic_acid_bacteria_on_D-_and_L-lactic_acid_contents_of_kimchi