

Advanced Viticulture and Enology of Cold-Hardy Fruit Musts: PIWI Grapes, Cider Pomology, and Wild Fermentation Dynamics in the Berlin-Brandenburg Terroir

The cultivation of fermentable fruit and the subsequent vinification of the harvested must represent a highly intricate intersection of agronomy, microbiology, and biochemical engineering. In marginal, cool-climate regions such as the Berlin-Brandenburg area of Germany, historical viticultural and pomological practices have been profoundly constrained by short growing seasons, late spring frosts, and elevated humidity, which traditionally fostered devastating fungal and bacterial pathogens¹. However, the convergence of shifting climatic baselines, the advanced breeding of cold-hardy, fungus-resistant grape varieties (PIWI), the preservation of heirloom cider apple genetics, and a resurgence in low-intervention, wild-fermentation enology has established a dynamic new frontier for northern fermented beverages.

The exhaustive analysis presented herein explores modern viticulture and orchard management tailored to cool-climate, sandy-soil terroirs. It meticulously details the agronomic management of PIWI grape varieties and heritage cider apples, the complex microbial succession underlying spontaneous alcoholic and malolactic fermentations, the physicochemical mechanisms of must clarification, and the oxidative thermodynamics governing the long-term maturation of wine in wooden casks.

Pedoclimatic Realities and Microclimate Engineering

The Berlin-Brandenburg region is characterized by challenging edaphic conditions—predominantly dry, rapidly draining sandy soils with notoriously low water-holding capacities and poor nutrient retention—coupled with a temperate, continental climate¹. While the region boasts a viticultural history dating back to the 13th century, often associated with monastic settlements and modern urban micro-sites such as the Viktoriapark in Kreuzberg, the Humboldthain park, and the Britzer Weinkultur, it demands highly specialized agricultural interventions to achieve commercial viability⁶.

The climatic realities of this region expose traditional *Vitis vinifera* and standard *Malus domestica* cultivars to severe abiotic and biotic stress. High precipitation and humidity during the vegetative cycle create ideal vectors for the proliferation of downy mildew (*Plasmopara viticola*), powdery mildew (*Erysiphe necator*), botrytis bunch rot (*Botrytis cinerea*), and apple scab (*Venturia inaequalis*)³. Traditional cultivation in such environments necessitates the intensive application of synthetic agrochemicals or heavy organic copper and sulfur spraying, which contradicts modern sustainability paradigms, risks phytotoxicity (such as fruit russetting

from copper formulations), and exacerbates soil compaction via repeated mechanical interventions¹⁰.

Windbreaks and Evapotranspiration Dynamics

A critical microclimate engineering tool for sandy soils is the integration of windbreaks or shelterbelts. The strategic placement of linear tree strips profoundly modifies atmospheric evaporative demand (AED)¹⁶. Windbreaks function by presenting a porous obstacle to approaching airflow, significantly reducing wind velocities by up to 70–80% on the leeward side¹⁷. This drastic reduction in turbulence directly depresses the vapor pressure deficit within the canopy, resulting in a parallel reduction of reference evapotranspiration by 15% to 30% during the summer growing season¹⁷. In soils with limited water retention, mitigating this evaporative loss preserves critical moisture, ensuring that the soil water content does not drop below the permanent wilting point of -1500 kPa ¹⁷.

Furthermore, windbreaks buffer extreme diurnal temperature fluctuations. By reducing convective heat loss, soils remain warmer during the day and retain more heat at night¹⁶. This microclimatic buffering provides a decisive defense against late spring frost events—such as the European Eiseiligen (Ice Saints) in mid-May—which frequently threaten the sensitive floral primordia of early-blooming cultivars in northern latitudes².

Viticulture of Cold-Hardy PIWI Grape Varieties

To navigate pathogenic constraints, modern viticulture in Berlin and Brandenburg is pivoting toward the cultivation of fungus-resistant grape varieties, known by the German acronym PIWI (*Pilz widerstandsfähige Rebsorten*). PIWIs are interspecific hybrids created by crossing high-quality European *Vitis vinifera* cultivars with American or Asian wild grape species (e.g., *Vitis labrusca*, *Vitis riparia*, or *Vitis amurensis*) that possess inherent evolutionary resistance to fungal pathogens and extreme cold¹⁰.

Genetic Introgressions and Appellation Legislation

Historically, interspecific hybrids suffered from a poor enological reputation due to the presence of undesirable organoleptic traits, most notably the "foxy" aroma profile associated with American genetics¹⁹. European appellation laws strictly mandated that quality wines bearing a Protected Designation of Origin (PDO) could only be produced from 100% pure *Vitis vinifera* vines²¹.

However, decades of backcrossing by pioneer breeders such as Valentin Blattner, and institutes including the Julius Kühn Institute (Geilweilerhof) and the Freiburg Wine Institute, have marginalized the non-*vinifera* genetic footprint³. Modern PIWIs retain critical resistance loci (such as the *Rpv* genes for downy mildew and *Ren* genes for powdery mildew) and utilize physical barriers like thickened cuticles to prevent Botrytis, while completely restoring the complex aromatic and structural profile of *Vitis vinifera*³. Recognizing the necessity of these sustainable cultivars, the European Union formally revised its agricultural regulations in 2021, granting member states the right to include resistant hybrid varieties in their appellation wines¹⁹. In Brandenburg, initiatives like the Vitis BB project are actively evaluating these modern PIWI vines specifically for their combined drought and frost tolerance in sandy terroirs⁴.

Cultivar Selection for the Northern Terroir

In cool-climate viticulture, the selection of the correct PIWI cultivar dictates the trajectory of the must. The most successful varieties engineered for regions like Brandenburg include both white and red cultivars optimized for early ripening, polygenic pathogen resistance, and cold hardiness:

PIWI Cultivar	Skin Color	Ripening Kinetics	Disease Resistance Profile	Key Aromatic & Enological Characteristics
Solaris	White	Very Early	Very High (Downy/Powdery mildew)	Aromatic, Muscat-like, tropical fruit notes. Accumulates massive sugar levels rapidly (often >24° Brix). Ideal for cool climates but requires careful harvest timing to prevent flabbiness ²² .
Regent	Red	Medium	High	Deep color, high tannin structure. Aromas of cherry and plum. Highly suitable for oak maturation and produces complex polymeric pigments ¹³ .
Cabernet	White	Medium-Late	High	Complex,

Blanc				spicy, extract-rich wines with aromas of green pepper, gooseberry, and citrus. Closely mimics the profile of Sauvignon Blanc ¹³ .
Calardis Soleil	White	Medium	Very High (includes Black Rot)	Loose cluster structure highly resistant to Botrytis. Ripens later than Solaris, making it ideal for avoiding premature acid degradation in warming seasons ²⁵ .
Souvignier Gris	White	Medium	High	Powerful, full-bodied wines with honeydew and apricot aromas. Often reminiscent of Pinot Gris with a creamy phenolic melt ¹³ .
Muscaris	White	Medium	High	Slender to full-bodied wines with vibrant acidity and intense,

				spicy Muscat bouquets. Excellent for aromatic dry styles or sweet wines ¹³ .
--	--	--	--	---

Architectural Canopy Management: Minimal Pruning vs. Guyot

The architectural manipulation of the vine canopy directly influences microclimate airflow, light interception, and carbohydrate partitioning. Traditional European viticulture relies heavily on cane-pruning systems such as the Single or Double Guyot²⁸. In a Guyot system, the vine is severely pruned during winter dormancy, leaving a single or double fruiting cane with a specific bud count²⁸. This restricts yields, forcing the vine to concentrate secondary metabolites into a smaller number of clusters, which is historically correlated with higher quality²⁸. However, emerging agronomic research in PIWI cultivation demonstrates the profound efficacy of "Minimal Pruning" or mechanical light pruning systems¹⁵. In minimal pruning systems, the vines are mechanically hedged rather than manually cane-pruned, leaving a vastly higher bud load³⁰.

The physiological response of the vine to minimal pruning fundamentally alters the phenological timeline. The vine produces significantly higher yields characterized by a massive proliferation of clusters with substantially smaller individual berries³⁰. Because the canopy volume and crop load are immense, the ripening process is intentionally delayed³². In warming climates, this delay is highly advantageous. It shifts the final ripening phase into the cooler autumn weeks, mitigating the rapid respiration of organic acids and preventing the excessive, rapid sugar spikes that lead to unbalanced, high-alcohol wines³². Research indicates that minimal pruning does not detrimentally lower the oenological quality of the must; rather, total sugar production per vine increases, while parameters like pH and Total Polyphenol Index (TPI) remain relatively stable, and manual labor costs decrease³⁰.

Cold-Hardy Pomology: Heirloom Orchards for Craft Cider

To comprehensively understand the conversion of fruit must into fermented beverages in cold, marginal soils, the viticultural analysis must be paralleled with the advanced pomology of heirloom cider apples (*Malus domestica*). Craft cider production from specialized, high-tannin apple varieties requires pedoclimatic adaptations and biochemical processing analogous to high-end viticulture, particularly in the arid sands of Brandenburg¹.

Rootstock Dynamics and Interstem Grafting

Just as grapevines rely on phylloxera-resistant rootstocks, the foundation of a productive apple orchard lies in the successful union of a scion cultivar with a selected rootstock¹. The rootstock dictates the tree's vigor, precocity, and susceptibility to both abiotic and biotic stresses¹. While global commercial production relies heavily on the highly dwarfing M.9 rootstock due to

its exceptional precocity, M.9 is extremely susceptible to fire blight (*Erwinia amylovora*) and struggles in the dry, rapidly draining sandy soils of Brandenburg due to its shallow root architecture¹. Alternatively, the Geneva series (e.g., G.41, G.214, G.890) provides unparalleled immunity to fire blight and replant disease¹. However, biomechanical constraints limit certain genotypes; histological evaluations reveal that G.41 forms a structurally weak, brittle graft union with specific scions, resulting in catastrophic breakage during high wind events¹.

To overcome the limitations of dwarfing rootstocks in arid landscapes, arborists employ interstem (or interstock) grafting. This technique involves a double-grafting process where a 15-30 cm section of a dwarfing rootstock (such as M.9) is grafted between a vigorous, deep-rooting, drought-tolerant rootstock (such as MM.111 or Antonovka) and the chosen fruiting scion¹. The vigorous root system penetrates deeply into the sandy soil profile, accessing lower water tables¹. Simultaneously, the M.9 interstem biochemically restricts vegetative vigor by altering the basipetal transport of auxins and the acropetal flow of cytokinins¹. This structural bottleneck shifts the tree's energy partitioning away from vegetative wood toward early floral initiation, combining the drought survival traits of a standard tree with the precise size control of a dwarf¹.

Architectural Canopy Management: The Oeschberg Method

For extensive *Streuobstwiese* (meadow orchard) systems utilizing vigorous rootstocks, traditional pruning paradigms often resulted in structural failure or the premature dieback of lower branches due to severe shading by the upper canopy¹. To counter this, the Oeschberg method (*Oeschbergschnitt*) is utilized to shape large, free-standing fruit trees¹.

The Oeschberg architecture establishes a strict hierarchical dominance: a single, dominant central leader (*Stammverlängerung*) trained in a narrow spindle, supported by three to four structurally dominant scaffold branches (*Leitäste*) radiating at steep 45-degree angles¹. By aggressively shortening and redirecting the scaffold branches outward, the arborist builds robust structural wood capable of bearing immense fruit loads¹. This specific pyramidal geometry ensures that sunlight easily penetrates the interior of the tree, maintaining a low Leaf Area Index (LAI) and maximizing Photosynthetically Active Radiation (PAR) interception, which is biochemically necessary for consistent annual floral initiation and even ripening across the entire volumetric space of the canopy¹.

Cellular Homeostasis: The Pathogenesis of Bitter Pit

Achieving premium fruit must requires exact management of plant nutrition. Apples are highly susceptible to physiological disorders, most notably bitter pit, which manifests as dark, sunken, necrotic lesions on the fruit peel¹.

Bitter pit is fundamentally a disorder of localized cellular calcium (Ca^{2+}) homeostasis¹.

Calcium is highly immobile and transported almost exclusively through the xylem¹. Because the expansive leaf canopy transpires at a vastly higher rate than the developing fruit, calcium is

preferentially partitioned to the vegetative shoots¹. At the cellular level, Ca^{2+} plays critical structural roles in stabilizing phospholipid bilayers and cross-linking pectic polysaccharides in

the middle lamella¹. When apoplastic free calcium is deficient, cell membranes lose structural integrity, leading to plasmolysis and localized necrosis¹.

Paradoxically, fruit suffering from bitter pit may exhibit high total calcium levels due to the excessive activity of pectin methyl esterase (PME). PME demethylates cell wall pectins,

exposing carboxyl groups that irreversibly bind available Ca^{2+} , locking it into the cell wall matrix and depleting the free apoplastic pool required for membrane stabilization¹. High

concentrations of potassium (K^{+}) and magnesium (Mg^{2+}) severely antagonize calcium uptake, competing directly at the root exchange sites and displacing calcium in the cell wall, resulting in tissue collapse¹. Mitigation requires strict vigor control, foliar calcium chloride sprays directly onto the fruit epidermis, and the avoidance of high-potassium fertigation¹.

LARS Classification and Cider Biochemistry

The production of premium hard cider requires the meticulous blending of specific apple typologies to achieve a harmonious organoleptic profile. The Long Ashton Research Station (LARS) established the definitive classification system based on titratable malic acid and total polyphenols (tannins) in the pressed must¹.

LARS Classification	Malic Acid (% w/v)	Tannin (% w/v)	Sensory Role in Must Blending
Sweet	< 0.45	< 0.20	Provides the fermentable sugar base and alcohol potential without astringency ¹ .
Sharp	> 0.45	< 0.20	Adds acidity, crispness, and lowers pH to microbiologically protect the must ¹ .
Bittersweet	< 0.45	> 0.20	The backbone of complex cider; imparts mouthfeel, body, and crucial phenolic structure ¹ .
Bittersharp	> 0.45	> 0.20	Extremely rare; provides perfect

			inherent balance of acid and tannin for single-varietal ciders ¹ .
--	--	--	---

Tannins are crucial for their potent antioxidant properties, which protect the fermenting must from deleterious oxidation and microbial spoilage¹.

The Biochemistry of Pre-Fermentation Processing

Whether processing grapes or apples, the biochemical manipulation of the crushed fruit determines the structural potential of the final wine.

Organic Acid Evolution in PIWI Musts

In grape viticulture, a critical challenge exacerbated by warming climates is the rapid respiration of organic acids during the ripening phase³³. The two dominant organic acids are tartaric acid and malic acid, which together constitute up to 90% of the berry's total acidity³³.

While tartaric acid concentrations remain relatively stable throughout ripening, malic acid is actively respired by the berry as a cellular energy source following véraison³³. In warm years, this respiration accelerates, leading to musts with elevated pH levels, decreased microbial stability, and flabby taste profiles³³. However, PIWI varieties exhibit distinct genetic controls over malic acid degradation rates. Cultivars like Solaris and Regent may retain sufficient malic acid in cooler northern limits, but require precise harvest timing to prevent a dramatic drop in total acidity³³. The tartaric/malic ratio at harvest is a fundamental metric, dictating the must's chemical buffering capacity and resistance to oxidative degradation³⁸.

Maceration and Keeving in Cider Production

In cider enology, pre-fermentation processing relies heavily on maceration (*cuvage*) and keeving (*défécation*). During maceration, milled apple pomace rests in open vats, where oxygen exposure triggers polyphenol oxidase (PPO) activity¹. This initiates the enzymatic browning of tannins, causing them to irreversibly adsorb to the solid cellular pulp¹. Paradoxically, extended oxygen contact reduces soluble tannin in the final juice, resulting in a softer mouthfeel, while facilitating the vital leaching of endogenous pectins¹.

Keeving is a method of deliberate nutrient deprivation designed to arrest fermentation prematurely, yielding a naturally sweet, clear cider. When juice is held at very low temperatures, pectin methyl esterase (PME) catalyzes the demethylation of linear homogalacturonan chains in the dissolved pectin¹. The newly exposed, negatively charged carboxyl groups cross-link with calcium ions to form a dense gel of calcium pectate¹. This negatively charged pectic acid matrix acts as a magnet, attracting positively charged nutrients, primarily yeast-assimilable nitrogen (YAN) and thiamine¹. As sluggish initial fermentation produces trace carbon dioxide, the bubbles lift this complex matrix of pectin, yeast cells, and bound amino acids to the surface, forming a dense brown cap (*chapeau brun*)¹. The winemaker racks the severely nutrient-depleted clear must from beneath the cap. Bereft of vital nitrogen, the subsequent fermentation proceeds at a highly retarded rate, eventually arresting completely while residual

fructose remains¹.

Ecological Succession in Wild Fermentation

The conversion of fruit must into wine via spontaneous, or "wild," fermentation harnesses the uninoculated indigenous microbiomes of the vineyard and the cellar, managing complex ecological successions rather than relying on sterile, monocultural inoculations of commercial yeast¹⁷.

The Non-Saccharomyces Pre-Fermentative Phase

The fresh must represents a highly selective, hyper-osmotic environment characterized by high sugar concentrations, low pH, and high dissolved oxygen¹⁸. In the initial 48 to 72 hours, the microbial landscape is entirely dominated by non-Saccharomyces yeasts, primarily species belonging to the genera *Hanseniaspora* (e.g., *H. uvarum*), *Candida* (*Starmerella bacillaris*), *Metschnikowia*, *Lachancea*, and *Torulaspora*¹⁷. Historically disparaged as spoilage organisms, these indigenous yeasts are now recognized as the primary drivers of aromatic complexity and terroir expression².

Their contribution is fundamentally enzymatic. Non-Saccharomyces yeasts secrete a vast array of extracellular hydrolytic enzymes that *Saccharomyces cerevisiae* lacks, most critically β -glucosidase⁴³. Many potent aromatic compounds in grapes, such as monoterpenes (linalool, geraniol) and C_{13} -norisoprenoids, exist bound to sugar molecules as odorless, non-volatile glycosides⁴⁴. The β -glucosidase enzymes cleave these β -glucosidic bonds, liberating the volatile aglycones into the wine matrix and massively amplifying the floral and fruity aromatic profile⁴⁴. Additionally, esterases and β -lyases secreted during this phase contribute to the formation of acetate esters and the liberation of volatile thiols⁴³.

Bio-Acidification via *Lachancea thermotolerans*

Within the non-Saccharomyces consortium, *Lachancea thermotolerans* holds exceptional biotechnological significance for mitigating climate-induced acid loss⁴⁸. *L. thermotolerans* possesses a unique metabolic pathway characterized by high lactate dehydrogenase (LDH) activity⁴². While standard yeasts ferment hexoses purely into ethanol and carbon dioxide via pyruvate, *L. thermotolerans* redirects a portion of the carbon flux away from ethanol production and into the synthesis of *L*-lactic acid⁵¹.

This "bio-acidification" process can generate substantial lactic acid, drastically decreasing the wine's pH (frequently by 0.3 to 0.5 units) and simultaneously lowering the final alcoholic strength by 0.2% to 0.7% (v/v)⁴². By naturally augmenting the total acidity without the artificial addition of tartaric acid powder, *L. thermotolerans* restores freshness, improves color stability, and fundamentally protects the wine by establishing a lower pH baseline for subsequent sulfur dioxide efficacy⁴².

Saccharomyces cerevisiae Ascension and Cell-to-Cell Contact Death

As the fermentation progresses, the accumulation of ethanol, the depletion of dissolved

oxygen, and nutrient scarcity create a highly toxic matrix³⁹. Non-Saccharomyces yeasts generally exhibit low ethanol tolerance (succumbing between 4% and 9% v/v)⁴¹. At this juncture, *Saccharomyces cerevisiae*, present only in trace amounts on the grape skin but highly endemic to winery environments, exponentially proliferates and seizes total ecological dominance to drive the fermentation to dryness².

The mechanism by which *S. cerevisiae* eradicates competing yeast species transcends simple ethanol toxicity; it is an aggressive, active biochemical warfare mechanism. *S. cerevisiae* secretes specific antimicrobial peptides (AMPs) into the fermenting must, which are lethal to non-Saccharomyces species such as *Hanseniaspora* and *Lachancea*⁵³. Mass spectrometry reveals that these AMPs are fragmented derivatives of the highly conserved glycolytic enzyme glyceraldehyde-3-phosphate dehydrogenase (GAPDH)⁵³.

During the exponential and early stationary growth phases, *S. cerevisiae* accumulates these GAPDH-derived peptides (termed saccharomycin) on the outer surface of its cell wall⁵³.

Through direct physical cell-to-cell contact, *S. cerevisiae* utilizes these surface-bound peptides to rupture and kill competing yeast cells, ensuring it monopolizes the remaining hexose

sugars⁵³. Additionally, indigenous *S. cerevisiae* strains intrinsically excrete sulfur dioxide (SO_2) as an intermediate metabolite in their sulfate assimilation pathway (ranging from 10 to 90 mg/L), further paralyzing sensitive competing microflora⁵⁸.

Nutrient Deprivation and Hydrogen Sulfide (H_2S) Metabolism

A critical risk factor in wild fermentations, particularly in the nutrient-poor sandy soils of Brandenburg, is Yeast Assimilable Nitrogen (YAN) starvation⁵⁹. YAN consists of primary amino acids and ammonium ions required for yeast protein synthesis⁵⁹.

When *S. cerevisiae* exhausts the available nitrogen pool, a severe metabolic crisis occurs within its sulfur reduction pathway. The yeast continuously reduces environmental sulfates and

sulfites into sulfide ions (S^{2-}) with the intention of binding them to nitrogenous carbon

skeletons (specifically O -acetylserine and O -acetylhomoserine) to synthesize the essential amino acids methionine and cysteine⁵⁸. If nitrogen is depleted, the yeast cannot synthesize these carbon-nitrogen backbones. The highly reactive sulfide ions accumulate intercellularly with no metabolic sink. To avoid immediate toxicity, the yeast cell rapidly vents these ions into

the wine as hydrogen sulfide (H_2S) gas⁵⁹. The presence of H_2S imparts severe reductive aromas of rotten eggs, requiring rapid intervention to prevent permanent taint⁵⁹.

Spontaneous Malolactic Fermentation (MLF)

Following alcoholic fermentation, wine frequently undergoes a secondary biological deacidification known as malolactic fermentation (MLF). In wild, low-intervention winemaking, this is facilitated by indigenous lactic acid bacteria (LAB), predominantly *Oenococcus oeni*⁶⁴. *O. oeni* thrives in the hostile post-fermentation matrix (high ethanol, low pH, nutrient depletion) where other bacteria perish. It utilizes the malolactic enzyme to catalyze the direct

decarboxylation of L-malic acid into L-lactic acid and carbon dioxide⁶⁶. Because lactic acid contains only one carboxylic acid group compared to malic acid's two, MLF naturally raises the wine's pH, significantly softens the acidic perception, and contributes complex secondary metabolites like diacetyl (imparting buttery aromas)⁶⁸.

The Danger of Biogenic Amines

A perilous consequence of spontaneous MLF by wild bacterial strains is the potential generation of biogenic amines (BAs). BAs are low-molecular-weight nitrogenous compounds produced via the enzymatic decarboxylation of free amino acids⁶⁶.

Certain wild strains of *O. oeni*, *Lactobacillus*, and *Pediococcus* possess specific, pyridoxal-5-phosphate-independent decarboxylase genes⁷⁰. If the wine pH is elevated and SO_2 levels are low, these bacteria actively decarboxylate amino acids to generate energy and forcibly raise their intracellular pH to survive⁷⁰.

The primary biogenic amines formed in wine are:

- **Histamine:** Derived from histidine. Highly vasoactive, causing severe allergic reactions, flushing, headaches, and respiratory distress in sensitive consumers⁷⁰.
- **Putrescine & Cadaverine:** Derived from ornithine/arginine and lysine, respectively. They utterly degrade the wine's organoleptic profile, masking varietal fruit with aromas of rotting meat, rancidity, and putrefaction⁷¹.

There are no legally mandated global upper limits for biogenic amines in wine, though safety advocates heavily monitor them⁶⁹. Managing BA production requires maintaining low-pH stability, minimizing wild *Pediococcus* populations, and racking off the lees early to remove the amino acid precursors liberated by yeast autolysis⁷⁰.

Colloidal Stability: Racking, Clarification, and Fining

The transition from a microbiologically active fermentation to a stable wine destined for aging requires precise physical and biochemical interventions.

Lees Management and Racking

Following alcoholic fermentation, dead yeast cells, grape pulp, and precipitated tartrates sink to form a dense sludge known as the "gross lees" (particles > 100 micrometers)³⁴. The wine must be meticulously racked (siphoned) off these gross lees within days to prevent cellular

putrefaction, the release of heavy mercaptans, and reductive H_2S defects³⁴.

Conversely, the smaller, suspended cellular fragments known as the "fine lees" are frequently retained and periodically agitated (*bâtonnage*)⁷⁸. During maturation, fine lees undergo autolysis—the enzymatic self-destruction of yeast cell walls. This releases mannoproteins, polysaccharides, and peptides into the wine matrix, increasing mid-palate volume, integrating harsh tannins, and providing a powerful reductive buffer against oxidation¹⁸.

Protein Stability and Bentonite Fining

White PIWI wines require rigorous clarification to prevent protein haze. Grape pathogenesis-related (PR) proteins, primarily thaumatin-like proteins (TLPs) and chitinases, are robust molecules with molecular weights ranging from 10 to 40 kDa and an isoelectric point

(pI) typically below 6.0⁸³. Resistant to low pH and proteolytic enzymes, these proteins are soluble at room temperature but denature upon exposure to heat, unfolding and aggregating to form a milky flocculation or "protein haze"⁸³.

The standard for eradicating protein instability is bentonite fining. Bentonite is a complex, multilayered hydrated aluminum silicate clay, predominantly composed of montmorillonite⁸⁶. Its efficacy relies entirely on electrostatic adsorption. In the acidic environment of wine (pH 3.0 - 3.6), PR proteins possess a net positive charge⁸⁶. When sodium or calcium bentonite is hydrated, the clay layers swell (with sodium bentonite exhibiting significantly higher swelling indices than calcium bentonite), exposing massive surface areas laden with exchangeable cations and presenting an overwhelming negative electrostatic charge⁸⁶. The negatively charged bentonite platelets violently attract the positively charged thaumatin and chitinase proteins, collapsing them into insoluble complexes that precipitate to the bottom⁸⁶. However, bentonite indiscriminately strips the wine of highly desirable aromatic compounds. Mass spectrometry confirms the removal of key volatile esters (e.g., ethyl octanoate) and monoterpenes. This is not solely due to direct clay adsorption, but is predominantly driven by hydrophobic interactions where the aromatic compounds remain physically bound to the proteins that are being precipitated⁸⁸.

Phenolic Management via Proteinaceous Fining Agents

Red wines (such as Regent) rely on proteinaceous fining agents to remove excessively harsh, astringent polyphenols and condensed tannins⁸⁹. Agents such as egg white (albumin), gelatin, casein, and isinglass (a highly pure collagen derived from fish bladders) operate by complexation⁸⁹.

When these positively charged proteins are introduced into the wine, they seek out negatively charged, highly reactive tannins. The resulting protein-tannin complexes grow in molecular weight until they become insoluble and precipitate⁸⁹. Isinglass is particularly prized for white wines and light reds, as it has a lower affinity for large condensed tannins but efficiently removes smaller polyphenolic monomers, yielding brilliant clarity without stripping the wine's intrinsic body⁹⁰. Furthermore, potassium caseinate demonstrates exceptional utility in selectively adsorbing ochratoxin A (OTA), a dangerous fungal metabolite, from contaminated musts⁹⁵.

Fining Agent	Primary Target	Mechanism of Action	Typical Application
Bentonite	PR-Proteins (Thaumatin/Chitinase)	Electrostatic adsorption (negative clay binds positive proteins) ⁸⁶ .	White wines (Haze prevention) ⁸⁶ .
Egg White	Large Condensed	Protein-tannin	Red wines (Tannin

	Tannins	complexation and precipitation ⁸⁹ .	softening) ⁸⁹ .
Isinglass	Small Phenolic Monomers	Gentle protein-polyphenol binding ⁹³ .	White / Light Red wines (Clarification) ⁹⁴ .
Potassium Caseinate	Oxidized Phenolics / Ochratoxin A	Adsorption and protein coagulation ⁸⁹ .	Defect remediation / Toxin removal ⁹⁵ .

Long-Term Storage in Wooden Casks

Following clarification, the long-term maturation or *élevage* of the wine in oak barrels chemically stabilizes pigment structure and alters tactile mouthfeel through a process of slow, continuous oxidation.

Micro-Oxygenation (MOX) and Phenolic Polymerization

The fundamental utility of the wooden cask lies in its physical porosity. The continuous, infinitesimal permeation of atmospheric oxygen through the oak staves and the bung is technically defined as the Oxygen Transmission Rate (OTR)⁹⁷. A standard 225-liter Bordeaux barrique introduces approximately 10 to 30 mg/L of oxygen into the wine per year⁹⁷.

This slow ingress of oxygen drives a cascade of metal-catalyzed redox reactions⁹⁷. The oxygen oxidizes small amounts of ethanol into acetaldehyde. In red wine maturation, acetaldehyde acts as an essential molecular bridge⁹⁷. It facilitates the covalent cross-linking and polymerization of free anthocyanins (unstable red pigment molecules) with condensed tannins⁹⁷. This tannin-anthocyanin condensation creates massive polymeric pigments that permanently lock

in the wine's deep color, rendering it highly resistant to bleaching by SO_2 ⁹⁷. The elongation of these polymers physically softens perceived astringency, transforming a harsh red wine into a structurally integrated product⁹⁷.

To simulate this without immense barrel expenditures, active Micro-Oxygenation (MOX) devices inject calibrated, continuous micro-doses of oxygen (e.g., 0.5 – 3 mg/L/month post-MLF) through porous ceramic diffusers into stainless steel tanks holding oak staves, mirroring the OTR of a natural barrel⁹⁷.

Cellar Thermodynamics and Evaporative Kinetics

The physical dynamics of a barrel are completely subservient to the thermodynamic realities of the cellar. As the barrel breathes, capillary evaporation occurs, where the wine migrates through the wood matrix and evaporates from the outer surface¹⁰¹. This creates a negative pressure vacuum and a physical void space at the top of the barrel known as the "ullage"¹⁰¹. If this ullage is not regularly eliminated by "topping up," the exposed surface area absorbs massive amounts of oxygen, leading to catastrophic oxidation and volatile acidity spikes¹⁰¹. The rate and composition of the evaporative loss are dictated by ambient humidity and

temperature¹⁰⁰. Water and ethanol possess different molecular weights and evaporation kinetics. In high humidity cellars (>70% RH), water evaporation is suppressed by the saturated atmosphere, causing ethanol to evaporate at a faster relative rate, ultimately *decreasing* the wine's total alcoholic strength¹⁰⁶. In low humidity cellars (<70% RH), the dry air extracts water molecules faster than the heavier ethanol vapors, leading to an overall concentration effect that *increases* the alcoholic strength¹⁰⁶.

Microbiological Spoilage in Barrels: The *Brettanomyces* Threat

While the barrel provides the optimal environment for oxidative maturation, its porous microstructure makes it an impregnable fortress for the most feared spoilage yeast in winemaking: *Dekkera bruxellensis* (anamorph: *Brettanomyces bruxellensis*)¹¹⁰.

Unlike *Saccharomyces*, *Brettanomyces* thrives in the nutrient-depleted, high-ethanol environment of a finished wine in a barrel¹¹⁰. *Brettanomyces* possesses a highly specific, two-step enzymatic pathway designed to metabolize the naturally occurring hydroxycinnamic acids in grapes (specifically *p*-coumaric and ferulic acids)¹¹⁰. First, phenolic acid decarboxylase (PAD) strips the carboxyl group, converting them into highly reactive vinylphenols¹¹⁰. Subsequently, vinylphenol reductase (VPR) reduces these intermediates into 4-ethylphenol (4-EP) and 4-ethylguaiaicol (4-EG)¹¹⁰.

Above sensory thresholds (around 400–600 $\mu\text{g/L}$ for 4-EP), these volatile phenols mask all varietal aromatics with overwhelming odors described as "horse sweat," "barnyard," and "medicinal band-aids"¹¹⁰.

Because *Brettanomyces* lodges up to 5.5 mm deep within the oak staves, the primary defense mechanism relies on the careful management of molecular sulfur dioxide (SO_2)¹⁰¹. The antimicrobial efficacy of SO_2 is strictly and logarithmically dependent upon the pH of the wine¹¹³. SO_2 exists in an equilibrium state; only the uncharged, molecular SO_2 can penetrate the yeast cell membrane to exert toxicity¹²⁰.

Wine pH	% Molecular SO2	Free SO2 Required for 0.8 mg/L Molecular Target (mg/L)	Free SO2 Required for 0.5 mg/L Molecular Target (mg/L)
3.00	6.1%	13	8
3.20	3.9%	21	13
3.40	2.5%	32	20

3.60	1.6%	50	31
3.80	1.0%	79	49
4.00	0.7%	125	78

Table: The critical relationship between wine pH and the concentration of antimicrobial molecular SO_2 ¹²⁰.

For *Brettanomyces* control, a molecular SO_2 concentration of 0.5 to 0.8 mg/L is generally required⁵⁸. A wine resting at a stable pH of 3.20 requires only 21 mg/L of free SO_2 for total protection¹²⁰. Conversely, a wine with a high pH of 3.80 requires an immense 79 mg/L of free SO_2 to achieve the exact same level of protection¹²⁰. Such high levels bleach the wine's color, harden tannins, and present regulatory issues⁵⁸. This underscores the absolute necessity of maintaining high acidity—either through early harvest, acid-retaining PIWI cultivars, or bio-acidification via *Lachancea thermotolerans*—to secure the wine during its oxidative maturation⁴².

Works cited

1. 35 Heirloom Apple Orchard Management.pdf
2. Not your ordinary yeast: non-Saccharomyces yeasts in wine, <https://academic.oup.com/femsyr/article/14/2/215/531574>
3. EU co-financed WiVitis project aims to promote PIWI cultivation - JKI, <https://www.julius-kuehn.de/en/news/nachricht/n/eu-kofinanziertes-projekt-wivitis-moechte-piwi-anbau-foerdern>
4. Vitis BB - Institut für Lebensmittel- und Umweltforschung e.V. (ILU), <https://www.ilu-ev.de/en/portfolio-items/vitis-bb-wine-is-also-grown-in-brandenburg-the-vitis-bb-project-aims-to-establish-a-climate-tolerant-and-modern-range-of-grape-varieties-in-brandenburg/>
5. Wine Country Brandenburg - MLEUV - Land Brandenburg, <https://mleuv.brandenburg.de/sixcms/media.php/9/Weinland-Brandenburg-en.pdf>
6. Britzer Wein, Berliner Weinkultur (ehem. Britzer Weingut) - Wein aus, <https://www.berliner-weinkultur.de/>
7. Britzer Weinkultur - Berlin.de, <https://www.berlin.de/tourismus-neukoelln/entdecken/artikel.1156813.php>
8. In the heart of Berlin, 300 grapevines grow at this enchanting spot in, <https://secretberlin.co/en/grapevines-in-berlin-kreuzberg-victoria-park/>

9. Humboldthain Park - visitBerlin.de,
<https://www.visitberlin.de/en/humboldthain-park>
10. Fungus-resistant grape varieties - PIWI - Austrian Wine,
<https://www.austrianwine.com/our-wine/grape-varieties/fungus-resistant-grape-varieties>
11. Biotransformation and Biological Regulation Mechanism of Aroma,
<https://www.sciopen.com/article/10.7506/spkx1002-6630-20220412-131>
12. PIWI factsheet Solaris 23-25_EN,
<https://agriculture.public.lu/dam-assets/innovation-forschung-projekte/piwi3/piwi-factsheet-solaris-23-25-en.pdf>
13. PIWI - Pilzresistente Rebsorten - Wines of Germany,
<https://www.winesofgermany.com/our-wine/wine-and-food/wine-and-food/387/piwi-pilzwidestandsf%C3%A4hige-reben>
14. AJEV PAPERS IN PRESS,
<https://www.ajevonline.org/content/early/2015/04/15/ajev.2015.14108.full.pdf>
15. PIWI sustainable - PIWI International,
<https://piwi-international.org/en/about-piwi/piwi-nachhaltig/>
16. Adoption and Impact of Fungus-Resistant Grape Varieties within,
<https://www.mdpi.com/2071-1050/16/14/6068>
17. Multi-omic characterization of microbial dynamics during ... - Frontiers,
<https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2026.1857803/pdf>
18. The Role and Use of Non-Saccharomyces Yeasts in Wine Production,
<https://www.gustodivino.it/allegati/The%20Role%20and%20Use%20of%20Non-Saccharomyces%20Yeasts%20in%20Wine%20Production.pdf>
19. Dipping Into PIWI Wines - Wineinsights,
https://www.wineinsights.org/2584395_dipping-into-piwi-wines
20. ProWein 2024: PiWis are the Next-Gen Grapes,
https://www.prowein.com/en/Media_News/Academy/Trend_topics/ProWein_2024_PiWis_are_the_Next-Gen_Grapes
21. EU grants member states the right to use resistant hybrid varieties in,
<https://www.decanter.com/wine-news/eu-grants-member-states-the-right-to-use-resistant-hybrid-varieties-in-appellation-wines-470864/>
22. Solaris grape | Cool Climate Wine Summit,
<https://coolclimatesummit.com/knowledge/piwi-varietals/solaris-grape>
23. EU grants member states the right to use resistant hybrid varieties in,
<https://piwi-international.org/en/2022/01/eu-grants-member-states-the-right-to-use-resistant-hybrid-varieties-in-appellation-wines/>
24. Solaris | EnoCodex, <https://eno.news/grapes/solaris>
25. cultivars - Förderverein Geilweilerhof, <https://geilweilerhof.eu/en/cultivars>
26. PIWI - Pilzresistente Rebsorten - Wines of Germany,
<https://www.winesofgermany.com/our-wine/wine-and-food/wine-and-food/387/piwi-fungus-resistant-vines>
27. Fungus-resistant white wine varieties - Vine nursery Kiefer & Sester,
<https://www.rebschule-sester.de/en/variety-description/fungus-resistant-white-v>

[arieties/](#)

28. Vine School: Part 1 – Common Vine-Training Systems,
<https://www.winescholarguild.com/blog/viticulture-insights/vine-school-part-1-common-vine-training-systems>
29. Pruning Vines: A Practical Guide to Professional Success,
<https://www.lebaroudeurduvin.com/en-es/blogs/infos/la-taille-de-la-vigne-guide-pratique-pour-une-reussite-professionnelle>
30. (PDF) Impact of pruning methods on the yield and the oenological,
https://www.researchgate.net/publication/287906355_Impact_of_pruning_methods_on_the_yield_and_the_oenological_quality_of_bobal_and_Tempranillo_grapevines
31. INCIDENCE OF MINIMAL PRUNING ON WINE QUALITY - ISHS,
https://ishs.org/ishs-article/978_36/
32. Evaluation of the impact of vine pruning periods on grape production,
<https://oeno-one.eu/article/view/8239>
33. Assessing Estonia's viticultural potential based on the compositional,
<https://journal.fi/afs/article/download/160985/115730/424369>
34. Racking Strategies: Why and when to employ - WineMakerMag.com,
<https://winemakermag.com/technique/racking-strategies-why-and-when-to-employ>
35. Characterization of varietal effects on the acidity and pH of grape,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2024.1439114/full>
36. Adaptive winemaking technologies using PIWI varieties in the wine,
<https://ives-openscience.eu/oiv2025-858/>
37. Investigating Evolution and Balance of Grape Sugars and Organic,
https://www.researchgate.net/publication/353740548_Investigating_Evolution_and_Balance_of_Grape_Sugars_and_Organic_Acids_in_Some_New_Pathogen-Resistant_White_Grapevine_Varieties
38. Modification of the organic acid profile of grapes due to climate,
<https://ojs.openagrar.de/index.php/VITIS/article/view/14834/14631>
39. (PDF) Non-Saccharomyces Yeasts: Biotechnological Role for Wine,
https://www.researchgate.net/publication/311456642_Non-Saccharomyces_Yeasts_Biotechnological_Role_for_Wine_Production
40. Multi-strain indigenous Yeast and Bacterial starters for 'Wild-ferment',
<https://cordis.europa.eu/project/id/315065/reporting>
41. Perspectives and Uses of Non-Saccharomyces Yeasts in Fermented,
<https://www.intechopen.com/chapters/64367>
42. pH Control and Aroma Improvement Using the Non-Saccharomyces,
<https://www.intechopen.com/chapters/78993>
43. (PDF) Non-Saccharomyces Yeasts from Organic Vineyards as,
https://www.researchgate.net/publication/374412608_Non-Saccharomyces_Yeasts_from_Organic_Vineyards_as_Spontaneous_Fermentation_Agents
44. Indigenous Non-Saccharomyces Yeasts With β -Glucosidase Activity,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9133630/>

45. Malolactic Fermentation—Theoretical Advances and Practical,
<https://www.mdpi.com/2311-5637/8/10/521>
46. Time-resolved volatile profiling and endpoint microbiome ... - SMUJO,
<https://smujo.id/jnpb/article/download/25013/9650/1230547>
47. Microbial Fermentation: A Sustainable Strategy for Producing High,
<https://www.preprints.org/manuscript/202512.2086>
48. Progress in Research and Application of Lachancea thermotolerans,
<https://www.sciopen.com/article/10.7506/spkx1002-6630-20211223-266>
49. Lachancea thermotolerans Applications in Wine Technology,
<https://pdfs.semanticscholar.org/a444/3f897f8dc0f9e2cd8c76c397440329a5f254.pdf>
50. External appearance of Lachancea thermotolerans - infowine.com,
<https://www.infowine.com/en/external-appearance-of-lachancea-thermotolerans/>
51. BIOACIDIFICATION WITH LACHANCEA THERMOTOLERANS - Laffort,
<https://laffort.com/en/bioacidification/>
52. Assessing the Impact of Commercial Lachancea thermotolerans,
<https://www.mdpi.com/2311-5637/10/6/303>
53. Saccharomyces cerevisiae accumulates GAPDH-derived peptides,
<https://pubmed.ncbi.nlm.nih.gov/28449125/>
54. Identification of novel GAPDH-derived antimicrobial peptides,
<https://pubmed.ncbi.nlm.nih.gov/24292082/>
55. (PDF) Identification of novel GAPDH-derived antimicrobial peptides,
https://www.researchgate.net/publication/259109765_Identification_of_novel_GAPDH-derived_antimicrobial_peptides_secreted_by_Saccharomyces_cerevisiae_and_involved_in_wine_microbial_interactions
56. Saccharomyces cerevisiae accumulates GAPDH-derived peptides,
<https://academic.oup.com/femsec/article/93/5/fix055/3753550>
57. Saccharomyces cerevisiae accumulates GAPDH-derived peptides,
https://www.researchgate.net/publication/316552401_Saccharomyces_cerevisiae_accumulates_GAPDH-derived_peptides_on_its_cell_surface_that_induce_death_of_non-Saccharomyces_yeasts_by_cell-to-cell_contact
58. SO₂ PRODUCTION BY WINE YEAST DURING ALCOHOLIC,
<https://www.lallemandwine.com/wp-content/uploads/2014/09/WE6-China.pdf>
59. Regulation of hydrogen sulfide liberation in wine-producing ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC167303/>
60. Taking the stink out of fermentation,
<https://www.awri.com.au/files/attachment/s1327/>
61. A Parallel Analysis of H₂S and SO₂ Formation by Brewing Yeast in,
https://www.researchgate.net/publication/281600914_A_Parallel_Analysis_of_H2S_and_SO2_Formation_by_Brewing_Yeast_in_Response_to_Sulfur-Containing_Amino_Acids_and_Ammonium_Ions
62. Hydrogen sulfide formation during fermentation: Effect of nitrogen,
https://www.researchgate.net/publication/284057210_Hydrogen_sulfide_formation_during_fermentation_Effect_of_nitrogen_composition_in_model_grape_must

63. Effects of rehydration nutrients on H₂S metabolism and formation of,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3226641/>
64. Impact of non-Saccharomyces in malolactic fermentation of white,
<https://ives-openscience.eu/3796/>
65. Biogeography of Oenococcus oeni Reveals Distinctive but,
<https://journals.asm.org/doi/10.1128/aem.02322-16>
66. Biogenic amines in wine: gene transcription as a tool for selection of,
<https://run.unl.pt/entities/publication/Oba4a2bc-48d2-465b-bc52-f0df9b0688eb>
67. New Malolactic Bacteria Strains Isolated from Wine Microbiota - MDPI,
<https://www.mdpi.com/2311-5637/8/1/31>
68. Physiological, Genetic, and Fermentative Traits of Oenococcus oeni,
<https://www.mdpi.com/2311-5637/11/8/440>
69. Malolactic & Biogenic Amines: Organoleptic alterations of wine,
<https://www.tentamus.com/news/malolactic-biogenic-amines-organoleptic-alterations-of-wine/>
70. Biogenic amines in wines: role of lactic acid bacteria,
https://www.profp.com.br/Qu%C3%ADmica%20do%20Vinho/lactic_acid_bacteria.pdf
71. LALLEMAND Wine Expert,
<https://www.lallemantwine.com/wp-content/uploads/2021/03/Wine-Expert-Biogenic-Amines-and-MLB-ENG-AUS.pdf>
72. Biogenic amine production by Oenococcus oeni - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/11927990/>
73. Biogenic amine production by Oenococcus oeni isolates ... - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/19435249/>
74. Biogenic Amine Production by Oenococcus oeni - ResearchGate,
https://www.researchgate.net/publication/11437431_Biogenic_Amine_Production_by_Oenococcus_oeni
75. BIOGENIC AMINES IN WINE - Wineitupanotch,
<https://www.wineitupanotch.com/blog/wineitupanotch/home/biogenic-amines-wine-allergic-reactions>
76. A review of biogenic amines in fermented foods - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10830535/>
77. The Influence of Winemaking Processes on the Formation of ... - MDPI,
<https://www.mdpi.com/2311-5637/11/12/687>
78. Gross Lees vs. Fine Lees — When to Rack, When to Stay | WineWiki,
<https://www.winewithseth.com/winewiki/gross-lees-vs-fine-lees/>
79. When to rack from secondary | Winemaking Talk,
<https://www.winemakingtalk.com/threads/when-to-rack-from-secondary.76921/>
80. Winemaking Techniques and Stylistic Choices,
<https://internationalwineauthority.com/winemaking-techniques-and-styles/>
81. About - Techniques in Home Winemaking,
<https://techniquesinhomewinemaking.com/blog/comment-page-11/>
82. Key concepts in winemaking;,
<https://www.gencowinemakers.com/docs/Key%20Concepts%20in%20Winemaking>

[ng-Zoom-9-13-2023.pdf](#)

83. Haze Formation and the Challenges for Peptidases in Wine Protein,
<https://pubs.acs.org/jafcau/article/69/48/14402/426483/Haze-Formation-and-the-Challenges-for-Peptidases>
84. Chemically Modified Clay Adsorbents Used in the Retention of,
<https://repository.utm.md/bitstream/handle/5014/27358/Nanomaterials-2024-V14-N7-AN588.pdf?sequence=1&isAllowed=y>
85. The Role of Proteins On Colloidal Formation in Italian Red Wine,
https://thesis.unipd.it/retrieve/dd6c1cb6-5238-4460-b897-9eca9d8ec0f9/Chikasha_Ratidzai%20Patience.pdf
86. White Wine Protein Instability: Mechanism, Quality Control and,
<https://www.mdpi.com/2306-5710/6/1/19>
87. Comparison of swelling index test (a) calcium bentonite, and (b),
https://www.researchgate.net/figure/Comparison-of-swelling-index-test-a-calcium-bentonite-and-b-sodium-bentonite_fig3_349924609
88. Study of combined effect of proteins and bentonite fining on the wine,
<https://pubmed.ncbi.nlm.nih.gov/25665100/>
89. Finings meaning in winemaking - Familia Morgan Wine,
<https://familiamorgan.com/learn/glossary/finings/>
90. faq on wine fining - ENARTIS NEWS,
<https://www.enartis.com/en/newsletters/enartis-news-faq-on-wine-fining/>
91. 218. Effect of fining with various proteins on tannin fractions,
<https://wine.ucdavis.edu/sites/g/files/dgvnsk2676/files/research-summaries/218%20effect%20of%20fining%20with%20various%20proteins%20on%20tannin%20fractions.pdf>
92. Different effectiveness of protein fining agents when tested for,
<https://oenone.eu/article/view/7264>
93. Fining Agents - The Australian Wine Research Institute,
https://www.awri.com.au/industry_support/winemaking_resources/frequently_asked_questions/fining_agents/
94. Fining agents - French wine tours,
<https://www.grapesandcorks.com/blog/fining-agents>
95. Removal of Ochratoxin A in Red Wines by Means of Adsorption,
<https://pubs.acs.org/doi/10.1021/jf010137o>
96. Fining Agents | WineWiki - Wine with Seth,
<https://www.winewithseth.com/winewiki/fining-agents/>
97. Micro-Oxygenation & Tannin Polymerization (MOX) | WineWiki,
<https://www.winewithseth.com/winewiki/micro-oxygenation-tannin-polymerization-mox/>
98. Effect of micro-oxygenation on color of wines made with toasted,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11726594/>
99. Artificial Intelligence Methods for Constructing Wine Barrels with a,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7397268/>
100. How much oxygen enters through a barrel? - Agrovin,
<https://agrovin.com/en/how-much-oxygen-enters-through-a-barrel/>

101. Impact of Barrel Kinetics and Dynamics on Wine - WineMakerMag.com,
<https://winemakermag.com/technique/barrel-kinetics-impact-on-wine>
102. Micro-oxygenation principles - Enartis,
https://www.enartis.com/en-us/technical_materials/micro-oxygenation-principles/
103. Micro-oxygenation: how this technique influences the style of,
<https://www.lasommeliere.com/en/blog/micro-oxygenation-how-this-technique-influences-the-style-of-modern-wines--n848>
104. Red Wine Aging by Different Micro-Oxygenation Systems and Oak,
<https://www.mdpi.com/2227-9717/8/10/1250>
105. Use of new tools for red wine aging: active and passive ... - Infowine,
<https://www.infowine.com/en/use-of-new-tools-for-red-wine-aging-active-and-passive-microoxygenation-with-oak-wood/>
106. Barrel temp & humidity | Winemaking Talk,
<https://www.winemakingtalk.com/threads/barrel-temp-humidity.36573/>
107. Understanding Wine Barrel Physics | PDF | Chemistry - Scribd,
<https://www.scribd.com/document/522105836/How-Barrels-Work>
108. The Benefits Of Implementing A Barrel Tracking System - BarrelWise,
<https://www.barrelwise.ca/blog/the-benefits-of-implementing-a-barrel-tracking-system>
109. In-situ monitoring of wine volume, barrel mass, ullage pressure and,
https://www.researchgate.net/publication/342653772_In-situ_monitoring_of_wine_volume_barrel_mass_ullage_pressure_and_dissolved_oxygen_for_a_better_understanding_of_wine-barrel-cellar_interactions
110. Use of Autochthonous Yeasts and Bacteria in Order to Control,
<https://www.mdpi.com/2311-5637/3/4/65>
111. Wine Bacteria to Control Volatile Phenols and Brettanomyces,
<https://www.lallemandwine.com/wp-content/uploads/2017/05/WUP-2-2014-South-Africa.pdf>
112. Significance of Brettanomyces and Dekkera during Winemaking,
<https://pdfs.semanticscholar.org/46f8/cc60ebf7df6ee44fd3272d1fb5a9b861ee0d.pdf>
113. Molecular Tools to Exploit the Biotechnological Potential of ... - MDPI,
<https://www.mdpi.com/2076-3417/11/16/7302>
114. Survival patterns of Dekkera bruxellensis in wines and inhibitory,
<https://repositorio.ulisboa.pt/server/api/core/bitstreams/c520fe7e-fd4b-4c86-a337-bed771b2846f/content>
115. Brettanomyces bruxellensis yeasts: impact on wine and winemaking,
https://iris.unito.it/bitstream/2318/1655567/4/Agnolucci%20et%20al.%2C%202017_wjmb.pdf
116. Relation between coumarate decarboxylase and vinylphenol,
<https://pubmed.ncbi.nlm.nih.gov/25955288/>
117. brettanomyces bruxellensis and off-odours - DISAFA,
https://www.disafa.unito.it/do/documenti.pl/ShowZip?_id=nlkd;field=file;file=CAMPOLONGO%2Epdf;key=oJlIT6tTgjcaeKAtP5JdYgQOzB5O1Q2IUpVYHPmrNL6HwtDLP7m1CcwXDAOAEHd5k;t=0435

118. The formation of volatile phenols from their hydroxycinnamic acids,
https://www.researchgate.net/figure/The-formation-of-volatile-phenols-from-their-hydroxycinnamic-acids-precursors-Slika-1_fig1_309490750
119. Molecular Diagnosis of *Brettanomyces bruxellensis*' Sulfur Dioxide,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6004410/>
120. DISTRIBUTION OF FREE SO₂ | Enartis,
<https://www.enartis.com/wp-content/uploads/2020/04/Distribution-of-Free-SO2.pdf>
121. (PDF) Metabolomic Analysis of SO₂ Tolerance and pH Effects on,
https://www.researchgate.net/publication/295633497_Metabolomic_Analysis_of_SO2_Tolerance_and_pH_Effects_on_Brettanomyces_bruxellensis
122. Innovative Fermentation Approach Employing *Lachancea* ... - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12385897/>