

# **The Biochemical and Rheological Engineering of a Regionalized Tarte Tatin: A Berlin-Brandenburg Case Study**

The execution of a classic Tarte Tatin represents one of the most demanding thermodynamic and biochemical orchestrations in the culinary arts. Fundamentally, the pastry is a biphasic system: an inverted, load-bearing matrix of dehydrated, caramelized fruit suspended within a pectin-stabilized hydrocolloid gel, hermetically sealed by a highly laminated, steam-leavened pastry crust. While standard, globalized culinary formulations rely on highly refined commercial ingredients with predictable behaviors, constructing this architectural pastry utilizing strictly localized ingredients from the Berlin and Brandenburg regions introduces immense physicochemical complexity. The reliance on regionally milled flour, cultured dairy, wild yeast sourdough fermentation, raw beet sugar, and root-cellar-stored heritage apples forces a radical departure from conventional baking parameters.

This comprehensive report provides an exhaustive analysis of the biophysical, rheological, and biochemical mechanisms required to engineer a localized Berlin Tarte Tatin. It systematically delineates the agricultural sourcing of the constituent elements, the proteolytic degradation of the gluten network via lactic acid bacteria, the complex crystal geometry and phase behavior of cultured butter during lamination, the Arrhenius kinetics of beet sugar caramelization, and the thermodynamic variables governing the final cast-iron baking and demolding processes. The resultant protocol serves as a blueprint for manipulating localized agricultural outputs into highly refined pastry architectures.

## **Agricultural Terroir and Ingredient Specifications**

The structural integrity, thermodynamic resilience, and organoleptic profile of this specific Tarte Tatin are dictated entirely by the biochemical composition of the regional ingredients selected from the agricultural zones surrounding Berlin. Each component introduces specific kinetic and mechanical variables that must be tightly controlled.

### **The Apple Matrix: Post-Harvest Physiology of Brandenburg Heritage Cultivars**

The selection of the apple cultivar is the most critical variable in determining the structural survival of the fruit matrix under extreme thermal stress. Modern dessert apples, cultivated primarily for immediate raw consumption, fail catastrophically during the prolonged boiling phase of a Tarte Tatin, rapidly disintegrating into a structureless puree due to the rapid thermal degradation of their cellular middle lamella. To combat this, heritage cultivars historically adapted to the Brandenburg climate must be employed; specifically, the 'Belle de Boskoop' and the 'Freiherr von Berlepsch'<sup>1</sup>.

The 'Belle de Boskoop', discovered as a chance seedling in the Netherlands in 1856 and widely adapted across German orchards, is a classic winter rennet apple characterized by its immense size (often reaching up to 200 grams per fruit), rough russeted skin, and profoundly dense, tart

flesh<sup>2</sup>. The 'Freiherr von Berlepsch', cultivated in the Rhineland in 1880 by Diedrich Uhlhorn Jr. from a cross of Ananas Reinette and Ribston Pippin, possesses exceptional acidity, an unusually high vitamin C content (up to 23.5 mg per 100 g), and distinct, spicy, pineapple-like aromatic notes<sup>3</sup>. Both cultivars are highly acidic, which is an absolute biochemical necessity for both the organoleptic balance of the intensely sweet caramel and the chemical catalysis of pectin gelation and sucrose inversion<sup>2</sup>.

Crucially, these apples must undergo prolonged root-cellar storage prior to utilization. Immediately following the autumn harvest, these winter apples exhibit excessive firmness and overwhelming astringency due to high concentrations of insoluble protopectin and complex phenolic compounds<sup>7</sup>. During prolonged storage in a cold, high-humidity root cellar (typically maintained at 3°C to 4°C), the harvested fruit remains metabolically active and continues to respire<sup>2</sup>. Over a period of several months, endogenous enzymes, primarily polygalacturonase and pectin methylesterase, gradually hydrolyze the rigid cell wall protopectins into water-soluble pectins<sup>8</sup>. This enzymatic activity slightly softens the flesh while simultaneously preserving enough structural rigidity in the cellulose matrix to withstand high-temperature baking<sup>8</sup>.

Concurrently, post-harvest physiology dictates a slow metabolic shift. While complex starches are enzymatically converted into soluble sugars (fructose, glucose, and sucrose), the total titratable acidity—dominated by malic and citric acids—decreases only slightly, preserving the sharp tartness required for the final dish<sup>7</sup>. This highly controlled cellular senescence, unique to cellar-stored winter varieties, yields an apple matrix with the precise physicochemical properties necessary for extreme high-heat caramelization.

| Cultivar               | Origin | Primary Organic Acid   | Pectin Quality      | Structural Baking Integrity                     | Organoleptic Profile            |
|------------------------|--------|------------------------|---------------------|-------------------------------------------------|---------------------------------|
| Belle de Boskoop       | 1856   | Malic Acid (High)      | High Methoxyl (HMP) | Exceptional; retains distinct cellular boundary | Tart, sharp, intensely robust   |
| Freiherr von Berlepsch | 1880   | Malic Acid (Very High) | High Methoxyl (HMP) | Moderate to High                                | Pineapple-like, spicy, aromatic |

### The Carbohydrate Matrix: Localized Beet Sugar Thermodynamics

While globalized pastry applications rely almost exclusively on highly refined sugarcane, a strictly Berlin-localized recipe necessitates the use of sugar derived from *Beta vulgaris* (sugar

beets) cultivated in the surrounding agricultural belts. Chemically, fully refined beet sugar and cane sugar are nearly identical, both consisting of roughly 99.95% pure sucrose<sup>10</sup>. However, the remaining 0.05% of trace impurities drastically alters their thermodynamic behavior and flavor development during severe thermal caramelization<sup>10</sup>.

Because sugar beets are root vegetables grown entirely underground, the extracted juice contains unique non-sugar components, including specific amino acids, complex carbohydrates, inorganic minerals, and organic compounds such as betaine<sup>10</sup>. During the thermal extraction and crystallization processes, minute fractions of these impurities remain inextricably bound to the sucrose crystals<sup>10</sup>. Advanced sensory and chromatographic analyses demonstrate that under high thermal stress, these specific beet-derived impurities undergo unique, localized Maillard reactions. These reactions generate volatile compounds characterized by earthy, oxidized, and slightly barnyard-like aromas, along with a distinct burnt-sugar aftertaste that contrasts sharply with the fruity aromas typically generated by cane sugar<sup>10</sup>.

These trace impurities also act as localized catalytic sites, meaning unrefined beet sugar caramelizes less uniformly and requires far more stringent thermal management than cane sugar to prevent premature scorching and catastrophic bittering<sup>12</sup>. Consequently, the application of local beet sugar in a Tarte Tatin grounds the flavor profile heavily in the regional terroir, providing a dark, robust caramel capable of standing up to the extreme acidity of the cellar-stored apples.

## **The Lipid Phase: Sauerrahmbutter and Emulsion Chemistry**

Lamination requires a highly plastic, low-moisture continuous fat phase capable of forming discrete, unbroken sheets between alternating layers of dough. For this regionalized iteration, *Sauerrahmbutter* (cultured butter) produced by regional dairies such as the Gläserne Molkerei in Brandenburg is utilized<sup>14</sup>. Unlike standard American sweet cream butter, which relies on a neutral pH and a standardized 80% fat content, European-style cultured butter undergoes a rigorous biological fermentation prior to the mechanical phase inversion of churning<sup>15</sup>.

The raw or pasteurized cream is inoculated with a carefully calibrated mesophilic starter culture consortium, typically comprising *Lactococcus lactis* subsp. *cremoris*, *Lactococcus lactis* biovar. *diacetylactis*, and *Leuconostoc mesenteroides*<sup>16</sup>. As these obligate fermenters proliferate at an optimized incubation temperature of approximately 20°C, they metabolize available lactose into lactic acid via the homofermentative glycolytic pathway, rapidly depressing the pH of the plasma from a neutral 6.7 to an acidic optimum of 4.4 to 5.0<sup>16</sup>. This acidification profoundly impacts the physical chemistry of the emulsion by neutralizing the highly protective negative zeta potential of the Milk Fat Globule Membrane (MFGM)<sup>16</sup>. The electrostatic repulsion that previously maintained the dispersion collapses, allowing the fat globules to flocculate and rendering them highly susceptible to the mechanical shear forces applied during churning<sup>16</sup>. Furthermore, the signature organoleptic profile of cultured butter is driven by the microbial synthesis of diacetyl (2,3-butanedione)<sup>16</sup>. Citrate-positive bacterial strains transport endogenous citric acid from the milk plasma into the intracellular matrix, enzymatically cleaving it into acetate and oxaloacetate, and eventually decarboxylating it into highly aromatic

diacetyl<sup>16</sup>. The Gläserne Molkerei Sauerrahmbutter possesses an 82% minimum fat content, ensuring a lower aqueous fraction (approximately 16% moisture)<sup>15</sup>. This specific lipid-to-water ratio is critical; it minimizes unwanted, excessive gluten development during the mechanical lamination process while simultaneously providing the exact volume of aqueous plasma required for optimal steam generation during baking<sup>15</sup>.

**The Structural Scaffolding: Regional Wheat Milling Profiles**

The architectural foundation of the puff pastry requires precise protein optimization to ensure maximum extensibility without sacrificing gas-retention capabilities. Flour sourced from regional mills, such as the Adler Mühle in Bahlingen, provides highly specific milling profiles tailored to these exact requirements<sup>19</sup>. The German flour classification system dictates the mineral ash content per 100 grams of dry matter. For laminated sourdough pastry, a carefully calibrated blend of *Weizenmehl Type 550* and *Type 812* is strictly required<sup>19</sup>. Type 550 serves as the primary structural scaffolding. With an extraction rate of roughly 64-71%, it is milled primarily from the starchy endosperm, possessing exceptional viscoelastic properties<sup>21</sup>. It provides the strong, resilient gluten network necessary to physically trap expanding steam and separate the hundreds of discrete butter layers during baking<sup>22</sup>. However, to introduce regional terroir and provide complex fermentation substrates for the wild sourdough culture, a smaller percentage of Type 812 is introduced into the *détrempe*<sup>19</sup>. Type 812 contains a higher mineral ash content, incorporating microscopic fractions of the wheat bran and germ<sup>21</sup>. This darker flour enhances the enzymatic activity during bulk fermentation by providing micronutrients to the lactic acid bacteria, while also contributing a darker, more robust flavor profile that synergizes with the earthy bitterness of the beet sugar caramel<sup>24</sup>.

| German Flour Classification | Extraction Profile                 | Primary Application in Lamination       | Rheological Contribution                                                 |
|-----------------------------|------------------------------------|-----------------------------------------|--------------------------------------------------------------------------|
| Weizenmehl Type 550         | Low ash, endosperm dominant        | Base matrix (approx. 80% of total mass) | High elasticity, optimal gas retention, strong protein network           |
| Weizenmehl Type 812         | Higher ash, partial bran inclusion | Flavor/Enzyme supplement (approx. 20%)  | Increased extensibility, enhanced Maillard browning, microbial substrate |

**Microbiological Kinetics and Rheology of the**

# Sourdough Détrempe

Traditional *pâte feuilletée* (puff pastry) relies entirely on mechanical steam generation for leavening, largely devoid of any biological fermentation<sup>25</sup>. However, integrating a localized sourdough starter (levain) into the *détrempe* (base dough) transforms the pastry into a highly complex, dynamic biochemical system. This biological addition serves a dual purpose: it

provides secondary leavening via carbon dioxide ( $CO_2$ ) production, but more importantly, it biochemically modifies the macroscopic rheology of the gluten network through extreme, acid-driven proteolysis<sup>25</sup>.

## Acidification Kinetics and the Lactic-to-Acetic Ratio

The sourdough starter represents a symbiotic microbial consortium of wild yeasts (e.g., *Saccharomyces cerevisiae*) and Lactic Acid Bacteria (LAB), predominantly *Lactobacillus sanfranciscensis*, *Lactiplantibacillus plantarum*, and *Lactobacillus brevis*<sup>29</sup>. The metabolic byproducts of these microorganisms radically alter the physical chemistry of the dough over

time. While the yeasts primarily metabolize simple sugars into ethanol and  $CO_2$  for volumetric expansion, the LAB dominate the biochemical conditioning of the dough via the continuous production of organic acids<sup>33</sup>.

Homofermentative LAB species utilize the Embden-Meyerhof-Parnas pathway to produce strictly lactic acid, imparting a smooth, relatively mild acidity<sup>33</sup>. Heterofermentative LAB, conversely, utilize the phosphoketolase pathway to produce a mixture of lactic acid, acetic acid, ethanol, and  $CO_2$ <sup>35</sup>. The precise ratio of lactic to acetic acid within the dough governs both the final flavor profile and the physical strength of the gluten matrix. Acetic acid imparts a sharp, aggressive tang, while lactic acid provides a gentler profile that is far less destructive to the rheology of the dough<sup>33</sup>.

For a highly extensible laminated pastry, the baker must actively manipulate the thermodynamics of the fermentation to strongly favor homofermentative lactic acid production. Biological fermentation roughly doubles or triples in rate per 10°C rise within the

working range (governed by the  $Q_{10}$  temperature coefficient)<sup>33</sup>. By maintaining a highly hydrated starter and proofing the *détrempe* at warmer ambient temperatures (approximately 26°C to 28°C), the microbial metabolism is forcefully steered toward lactic acid dominance<sup>33</sup>. This is critical, as excessive acetic acid creates a highly rigid, short gluten network that will tear catastrophically during the extreme mechanical shear stress applied during lamination<sup>33</sup>.

## Proteolysis and Depolymerization of the Glutenin Macropolymer (GMP)

The most profound biophysical consequence of incorporating sourdough into puff pastry is the severe proteolytic degradation of the structural gluten proteins. Wheat gluten is a massive, hydrated protein complex comprising two primary fractions: gliadins (monomeric, single-chain proteins responsible for viscosity and extensibility) and glutenins (polymeric proteins linked by

immense intermolecular disulfide bonds, responsible for elasticity and dough strength)<sup>37</sup>. The largest of these polymeric structures aggregate to form the Glutenin Macropolymer (GMP), which ultimately dictates the absolute structural integrity of the dough<sup>28</sup>.

As the LAB proliferate exponentially during the bulk fermentation of the *détrempe*, they rapidly excrete lactic and acetic acids, forcing the pH of the system down from a neutral 6.0 to a highly acidic range of 3.8 to 4.2<sup>31</sup>. This specific pH depression acts as a universal biochemical trigger. At pH values below 4.4, the endogenous aspartic proteinases natively present within the wheat flour are hyper-activated<sup>27</sup>. Simultaneously, the acidic environment increases the net positive charge of the protein chains, inducing massive electrostatic repulsion that physically unfolds the gluten molecules, exposing their previously hidden, highly conserved peptide bonds to enzymatic attack<sup>27</sup>.

Once activated, these cereal proteases, working in synergistic tandem with the extracellular peptidases and cell-wall-associated serine proteinases secreted directly by the LAB (such as *L. sanfranciscensis* and *L. plantarum*), begin systematically cleaving the peptide bonds of the gluten proteins<sup>31</sup>. Advanced electrophoretic analyses and Fourier transform infrared spectroscopy (FTIR) demonstrate that during this phase, the secondary structure of the GMP undergoes radical modification; the alpha-helix structures diminish, while beta-turns and random coils increase, signifying increasing molecular flexibility<sup>42</sup>. The high-molecular-weight (HMW) glutenin subunits are aggressively hydrolyzed, resulting in the rapid, macroscopic depolymerization of the Glutenin Macropolymer<sup>28</sup>. Furthermore, the LAB exhibit a specific proteolytic affinity for gliadin fractions, actively degrading the proline- and glutamine-rich peptides, which additionally alters the dough's flow behavior<sup>31</sup>.

## **Rheological Implications for Extensibility and Elastic Recoil**

This microscopic biochemical degradation translates directly into macroscopic, highly desirable rheological changes for the lamination process. The cleavage of the disulfide bonds and the systematic depolymerization of the GMP fundamentally shift the dough's viscoelastic behavior.

Analytical measurements confirm that the storage modulus ( $G'$ ), representing elasticity and structural recovery, drops precipitously, while the loss modulus ( $G''$ ), representing viscous flow, increases relative to unfermented controls<sup>27</sup>. The dough transitions from a highly resilient, rubbery mass into an exceptionally soft, extensible matrix<sup>27</sup>.

For the pastry chef, this sourdough-induced relaxation is an engineered advantage. Traditional unfermented puff pastry suffers from severe elastic recoil; the intact gluten network resists the rolling pin, causing the dough to snap back and permanently rupture the delicate, microscopic internal butter layers<sup>25</sup>. The targeted proteolysis achieved via sourdough fermentation chemically neutralizes this elasticity, allowing the dough to be sheeted out to a fraction of a millimeter in thickness without fighting the physical force of the baker<sup>44</sup>.

However, this represents a highly precarious thermodynamic balance. If the bulk fermentation proceeds for too long, the proteolytic degradation becomes absolute; the gluten network liquefies entirely, losing its gas-trapping capability and yielding a dense, flat, and structurally collapsed pastry<sup>27</sup>. Strict thermal management—specifically, cold retarding the dough at 3°C to

4°C—is therefore required to suddenly halt the enzymatic activity immediately prior to the lamination phase, locking the gluten network in a state of optimal extensibility<sup>25</sup>.

## The Physical Chemistry and Geometry of Lamination

Lamination is the precise architectural process of suspending hundreds of alternating, microscopic layers of aqueous dough (the *détrempe*) and solid lipid (the *beurrage*)<sup>25</sup>. The success of this biophysical assembly hinges entirely on managing the thermal plasticity of the cultured butter and executing mathematically precise geometric folds.

### Triglyceride Polymorphism and Solid Fat Content Optimization

Bovine milk fat is an exceptionally complex and heterogeneous mixture of thousands of different triacylglycerol (TAG) species<sup>2</sup>. Because of this immense molecular diversity, milk fat does not possess a single, discrete melting point; rather, it melts over a broad thermodynamic continuum<sup>16</sup>. To function as a continuous, waterproof structural barrier during lamination, the *Sauerrahmbutter* must be thermally engineered to possess a specific Solid Fat Content (SFC)—ideally stabilized between 30% and 40%<sup>16</sup>.

If the SFC is too high (i.e., the butter is excessively chilled), the fat becomes a brittle, crystalline block that will shatter into discrete shards inside the dough during rolling, permanently destroying the continuous lamination<sup>16</sup>. Conversely, if the SFC is too low (i.e., the butter is too warm), the fat transitions into a highly viscous liquid that will smear directly into the surrounding flour matrix<sup>25</sup>. This smearing chemically shortens the gluten strands, resulting in a crumbly, cookie-like texture rather than distinct, flaky layers<sup>25</sup>.

Furthermore, the butter must be tempered to stabilize its crystal polymorphism. Triglycerides can pack into several distinct crystalline lattice structures. For optimal lamination, the butter

must be maintained primarily in the intermediate  $\beta'$  (Beta-prime) polymorphic state<sup>16</sup>. The  $\beta'$  crystals pack into an orthorhombic sub-cell lattice, yielding smooth, tiny, needle-like crystals that allow the butter to bend, stretch, and deform seamlessly across vast surface areas without breaking<sup>16</sup>. The baker must therefore physically pound the cold cultured butter with a heavy rolling pin to generate mechanical friction. This kinetic energy renders the lipid highly plastic

and malleable, aligning the  $\beta'$  crystals, while strict ambient temperature control keeps the block's core temperature strictly below its complete melting point (maintaining the butter precisely at 13°C to 15°C)<sup>25</sup>.

### Geometric Folding and the Mechanics of Steam Leavening

Once the plasticized *beurrage* is encased within the highly extensible, sourdough-fermented *détrempe*, the entire mass is subjected to extreme compressive shear stress via rolling, followed by a sequence of sequential folding<sup>25</sup>. In a classical application, the dough is subjected to a series of "letter folds" (folding the dough into thirds, overlapping itself)<sup>25</sup>.

The mathematics of lamination dictate that the number of discrete fat layers ( $L$ ) generated follows an exponential function based on the number of folds ( $n$ ) and the type of fold ( $f$ ),

where a letter fold equals 3):

$$L = f^n$$

A traditional unfermented puff pastry protocol utilizing six sequential letter folds generates exactly 729 discrete layers of fat ( $3^6 = 729$ ), separating 730 layers of dough<sup>25</sup>. However, because the proteolytic action of the sourdough LAB significantly weakens the tensile strength of the gluten, pushing the lamination to 729 layers often results in the microscopic dough sheets tearing under their own extreme thinness, causing the adjacent fat layers to coalesce<sup>44</sup>. Therefore, for sourdough-leavened pastry, the geometry is typically restricted to three or four folds, generating between 27 and 81 robust layers capable of withstanding the massive internal pressure of baking without rupturing prematurely<sup>25</sup>.

The primary mechanism of volumetric expansion during baking is physical steam leavening, which vastly outpaces the biological  $CO_2$  expansion provided by the yeast<sup>34</sup>. As the cast-iron skillet transfers intense thermal energy to the pastry crust, the ambient moisture trapped within both the dough and the 16% aqueous plasma fraction of the cultured butter flashes into steam<sup>15</sup>. Because water expands volumetrically by a factor of roughly 1,600 when transitioning from a liquid to a gas at atmospheric pressure, massive internal vapor pressure is generated rapidly.

The continuous layers of solid fat melt, leaving behind microscopic voids, while the surrounding liquid lipid acts as an impermeable hydrophobic barrier<sup>25</sup>. This barrier temporarily prevents the steam from escaping, forcing the expanding gas to violently thrust the individual layers of dough upward in a massive, vertical thermodynamic expansion<sup>25</sup>. Simultaneously, the gluten proteins undergo irreversible thermal coagulation, permanently setting the airy, multi-layered architecture before the vapor eventually fractures the crust and vents safely into the oven cavity<sup>23</sup>.

## Biochemistry of Apple Caramelization and Pectin Gelation

Beneath the intricate architecture of the pastry, the apple and sugar matrix undergoes a violent, high-temperature sequence of dehydration, acid hydrolysis, and non-enzymatic browning. This process converts the raw fruit and unrefined beet sugar into a structurally coherent, intensely aromatic hydrocolloid gel.

### Acid-Catalyzed Sucrose Inversion and Arrhenius Kinetics

The caramelization phase initiates by melting the unrefined beet sugar in the presence of the localized cultured butter within the cast-iron skillet. As the thermal energy increases, the sucrose crystals dissolve and begin to thermally degrade. When the highly acidic 'Belle de Boskoop' and 'Berlepsch' apples are introduced, the thermal shock causes their cellular vacuoles to rupture, rapidly exuding massive quantities of cellular water and concentrated

malic acid directly into the boiling sugar matrix<sup>9</sup>.

The sudden introduction of malic acid into the boiling sucrose forces a massive chemical cascade known as acid-catalyzed sucrose inversion<sup>52</sup>. The malic acid donates free protons ( $H^+$ ) to the solution. These protons aggressively attack the glycosidic bonds linking the glucose and fructose monomers within the sucrose molecule<sup>54</sup>. Under extreme heat, the sucrose is forcefully hydrolyzed into a highly reactive, equimolar mixture of free D-glucose and D-fructose, a solution colloquially known as invert sugar<sup>55</sup>.

The thermodynamic kinetics of this inversion reaction are heavily temperature-dependent and can be mathematically modeled using the classical Arrhenius equation:

$$k = A \exp \left( -\frac{E_a}{RT} \right)$$

where  $k$  is the reaction rate constant,  $A$  is the pre-exponential frequency factor,  $E_a$  is the activation energy,  $R$  is the universal gas constant, and  $T$  is the absolute thermodynamic temperature in Kelvin<sup>52</sup>. Advanced kinetic research indicates that the activation energy ( $E_a$ ) for the degradation of sugars in an acidic fruit matrix typically ranges from 50 to 118 kJ/mol, depending heavily on the specific pH and moisture content of the system<sup>57</sup>. As the ambient water aggressively boils off and the temperature of the skillet climbs past 120°C, the reaction rate constant ( $k$ ) scales exponentially.

The newly liberated fructose molecules are exceptionally thermodynamically unstable compared to intact sucrose. They rapidly undergo multiple dehydration reactions to form highly reactive dicarbonyl intermediates, most notably 5-hydroxymethylfurfural (HMF)<sup>52</sup>. The HMF molecules further polymerize and degrade into a vast array of volatile furans (imparting sweet, caramel notes), diacetyl (reinforcing the cultured butter aroma), and massive, dark-brown melanoidin polymers<sup>52</sup>. Due to the unique trace impurities inherent in the unrefined beet sugar (such as betaine and specific amino acids), parallel early-stage Maillard reactions occur rapidly, generating a darker, more earthy, and highly complex aromatic profile that distinctively marks the regional terroir of the dish<sup>10</sup>.

## **Polyphenol Oxidase Inactivation and High Methoxyl Pectin (HMP) Gelation**

Concurrently, the structural chemistry of the apples undergoes rapid transformation. When the apples are peeled and cut, their cellular structure is damaged, exposing endogenous phenolic compounds (specifically chlorogenic acid, the primary phenolic acid in the 'Boskoop' variety) to ambient oxygen and the highly active enzyme polyphenol oxidase (PPO)<sup>61</sup>. This enzyme acts as a rapid catalyst, oxidizing the chlorogenic acid into reactive o-quinones, which quickly

polymerize into dark brown, bitter melanin pigments<sup>64</sup>. While enzymatic browning is generally undesirable and degrades flavor in fresh fruit, the intense thermal shock of the 150°C boiling caramel instantly denatures the PPO enzyme, permanently arresting the enzymatic browning<sup>64</sup>. This rapid heat inactivation preserves the integrity of the remaining unoxidized chlorogenic acid, which contributes heavily to the complex, tart, and slightly astringent flavor profile of the finished matrix<sup>63</sup>.

As the apples boil in the acidic syrup, their rigid cell walls further degrade, releasing massive quantities of High Methoxyl Pectin (HMP) directly into the surrounding caramel<sup>66</sup>. HMP is structurally defined by a Degree of Esterification (DE) greater than 50%<sup>66</sup>. The biophysics of HMP gelation are exceptionally strict: the polymer will only form a three-dimensional gel network in the presence of high concentrations of a co-solute (specifically, the beet sugar concentration must exceed 55-60% by weight to severely reduce water activity) and under highly acidic conditions (the pH must be strictly depressed below 3.5)<sup>66</sup>.

The extreme malic acid concentration natively provided by the Boskoop and Berlepsch apples forcefully drives the pH of the boiling syrup down to approximately 3.0. At this critical threshold, the massive carboxyl groups distributed along the pectin polysaccharide chains are protonated, completely neutralizing their natural electrostatic repulsion<sup>66</sup>. Without this repulsion, and facilitated by the massive reduction in water activity driven by the dense, dehydrating invert sugar syrup, the pectin molecules are forced into intimate proximity. They spontaneously self-assemble, forming highly structured "junction zones" stabilized by dense, overlapping networks of intermolecular hydrogen bonds and massive hydrophobic interactions between their methyl ester groups<sup>66</sup>. As the Tarte Tatin cools post-baking, this HMP network solidifies into a rigid, highly cohesive, transparent gel that permanently cements the apples and the caramel together into a unified, structural disc capable of supporting the weight of the pastry<sup>6</sup>.

## **Thermodynamics of the Cast-Iron Vessel and Demolding Mechanics**

The entire architecture of the Tarte Tatin relies on managing extreme thermal mass. The use of a heavy, thick-walled cast-iron skillet is not a rustic aesthetic choice; it is an absolute thermodynamic requirement<sup>69</sup>.

Cast iron possesses a profoundly high volumetric heat capacity and exceptionally high thermal emissivity. During the stovetop caramelization phase, the cast iron acts as a massive thermal reservoir. It absorbs thermal energy and radiates it highly uniformly into the viscous sugar matrix, preventing the localized, high-temperature hot spots that would otherwise instantly incinerate the sensitive beet sugar impurities and ruin the Maillard profile<sup>69</sup>.

Once the laminated sourdough pastry crust is secured over the boiling apple matrix, the entire skillet is transferred into a high-temperature convection environment, typically calibrated between 205°C and 220°C (400°F–430°F)<sup>71</sup>. The extreme thermal mass of the iron initiates a massive bottom-up heat transfer, forcefully driving the remaining water out of the apples and concentrating the pectin-sugar gel, while the ambient convection heat initiates the violent

steam expansion of the sourdough puff pastry from the top down<sup>72</sup>. The baking phase is complete strictly when the pastry demonstrates maximum volumetric expansion and irreversible Maillard browning across the crust surface, typically requiring 30 to 40 minutes<sup>75</sup>. The final, and physically most precarious, phase of the process is the demolding. A precise thermodynamic window governs this action. If the tart is inverted immediately upon removal from the oven, the HMP pectin gel remains entirely above its critical setting temperature; the matrix acts as a low-viscosity fluid, resulting in catastrophic structural collapse as the boiling caramel flows out across the plate, leaving the apples behind<sup>71</sup>. Conversely, if the skillet is allowed to cool completely to room temperature, the pectin junction zones fully solidify and form unbreakable adhesive bonds with the porous microscopic surface of the cast iron, permanently cementing the tart to the pan<sup>71</sup>.

The baker must carefully monitor the cooling curve, waiting exactly 15 to 20 minutes<sup>71</sup>. At this precise inflection point, the temperature of the caramel drops to roughly 50°C—cool enough for the pectin to undergo a phase transition into a cohesive, load-bearing gel, but warm enough to remain highly plastic and non-adhesive to the iron<sup>66</sup>. An inverted plate is applied, and the entire high-mass system is rapidly flipped, utilizing gravity to effortlessly detach the stabilized apple disc onto the serving platter, completing the architectural assembly<sup>71</sup>.

## The Synthesized Operational Protocol: A Berlin-Localized Tarte Tatin

The following formulation represents the complete biochemical and operational synthesis of the parameters discussed, standardized for a 25-centimeter (10-inch) cast-iron skillet, providing an exhaustive manual for the regionalized pastry.

### Phase 1: Fermentation and Rheological Conditioning of the Détrempe

1. **Autolyse and Sourdough Incorporation:** In a standardized mixing vessel, combine 200 g of Adler Mühle Type 550 flour, 50 g of Adler Mühle Type 812 flour, and 125 g of highly chilled water (4°C) to restrict premature enzymatic activity. Execute a brief mechanical homogenization until a shaggy mass forms. Subject the mass to a 45-minute autolyse phase to initiate passive flour hydration, starch swelling, and early, unstructured gluten formation<sup>35</sup>. Following the autolyse, incorporate 60 g of a mature, homofermentative lactic-dominant sourdough starter (100% hydration) and 5 g of sodium chloride<sup>78</sup>.
2. **Gluten Alignment and Bulk Fermentation:** Execute a series of high-tension coil folds at 30-minute intervals for the first two hours to mechanically align the gluten strands into an organized network<sup>35</sup>. Cover the vessel and maintain the ambient environment at a highly controlled 25°C to 26°C for approximately 3 to 4 hours to favor lactic acid production over acetic acid<sup>30</sup>. During this phase, closely monitor the dough for rheological relaxation—the physical manifestation of the LAB lowering the pH and activating the endogenous aspartic proteinases that depolymerize the Glutenin Macropolymer<sup>28</sup>. Once the dough exhibits a 50% volumetric increase and extreme extensibility (successfully passing the windowpane test), violently arrest the fermentation by transferring the mass into a 3°C refrigeration unit for a minimum of 8 to 12 hours<sup>25</sup>.

## Phase 2: Plasticization and Geometry of the Lamination

1. **Beurrage Preparation:** Extract 225 g of Gläserne Molkerei Sauerrahmbutter. Place the lipid between two sheets of parchment paper. Utilizing a heavy rolling pin, apply extreme mechanical force to pound the butter into a uniform 15-centimeter square block<sup>25</sup>. This mechanical friction forces the crystalline network into the highly plastic  $\beta'$  polymorphic state, without raising the internal temperature past 15°C<sup>16</sup>. If the lipid exceeds this threshold, return it immediately to refrigeration<sup>25</sup>.
2. **Encasing and Sheeting:** Extract the highly extensible, cold-retarded *détrempe* from the refrigerator. Sheet it into a rectangle precisely twice the length of the butter block. Encase the lipid squarely within the dough, folding the dough over and sealing the outer edges to create a hermetic boundary, preventing lipid leakage<sup>25</sup>.
3. **Geometric Folding Schedule:** Execute three sequential single folds (letter folds), rolling the dough to roughly 60 cm by 20 cm, folding it into thirds, and rotating the mass 90 degrees after each operation. This geometry yields exactly 27 discrete layers of lipid ( $3^3 = 27$ )<sup>25</sup>. Due to the severe proteolytic weakening of the gluten by the sourdough LAB, attempting standard commercial laminations (e.g., 729 layers) will result in microscopic tearing of the dough sheets<sup>44</sup>. Between each fold, the dough must be rested at 3°C for exactly 30 minutes to permit structural relaxation of the surviving disulfide bonds and to re-stabilize the Solid Fat Content of the butter<sup>25</sup>. Upon completion of the third fold, sheet the dough to a 4-millimeter thickness, cut a 26-centimeter disc, perforate the surface with four discrete holes to provide necessary steam exhaust vents during baking, and return it to refrigeration<sup>25</sup>.

## Phase 3: Caramelization Kinetics and Apple Matrix Assembly

1. **Apple Preparation:** Select 1.5 kilograms of cellar-aged 'Belle de Boskoop' and 'Berlepsch' apples. Peel, core, and precisely segment the fruit into roughly equal thirds<sup>72</sup>. To preemptively halt the enzymatic browning catalyzed by polyphenol oxidase before the thermal shock, submerge the segments briefly in a highly acidic ascorbic solution (lemon juice)<sup>72</sup>.
2. **Sucrose Inversion and Maillard Browning:** Place a heavy 25-centimeter cast-iron skillet over a high-output stovetop thermal source. Distribute 150 g of unrefined local beet sugar in an even monolayer<sup>75</sup>. Allow the sucrose to melt undisturbed until the edges display the initial phases of thermal degradation. Introduce 60 g of diced Sauerrahmbutter, initiating rapid emulsification and foaming<sup>70</sup>.
3. **Fruit Introduction and Dehydration:** Once the caramel achieves a dark amber hue (signaling advanced melanoidin formation and temperatures exceeding 160°C), rapidly arrange the apple segments within the skillet in a tightly nested, concentric geometry, overlapping each piece by one-third to account for shrinkage<sup>72</sup>. The apples will immediately release massive volumes of aqueous malic acid into the boiling lipid-sugar emulsion. Maintain high thermal output for approximately 15 to 20 minutes<sup>70</sup>. During this

phase, the malic acid violently catalyzes the inversion of the beet sucrose into glucose and fructose, while the cellular walls of the apples degrade, flooding the syrup with High Methoxyl Pectin<sup>9</sup>. The phase is complete strictly when the ambient water is entirely driven off, the bubbling becomes dense and viscous, and the apples are deeply saturated with the dark, inverted caramel<sup>72</sup>. Remove from the heat source and allow the matrix to cool for exactly 15 minutes, permitting the escaping steam to subside<sup>75</sup>.

## Phase 4: Thermodynamic Baking and Demolding

1. **Steam Expansion:** Preheat a convection oven to a highly aggressive 215°C (420°F)<sup>69</sup>. Retrieve the chilled, laminated sourdough disc and drape it tightly over the warm apple matrix, forcefully tucking the exterior perimeter down inside the inner wall of the cast-iron skillet to form a physical seal<sup>69</sup>.
2. **Thermal Coagulation:** Insert the entire heavy vessel into the thermal environment. The radiant and conductive heat will rapidly vaporize the 16% aqueous plasma trapped within the cultured butter layers<sup>15</sup>. This massive vapor pressure will violently force the 27 layers apart, generating the macroscopic lift of the pastry, while the severe oven temperature initiates the Maillard browning of the exterior crust<sup>25</sup>. Bake until the structural expansion halts and the surface achieves a profound, dark mahogany color, typically requiring 30 to 40 minutes depending on precise oven thermodynamics<sup>72</sup>.
3. **The Inversion:** Extract the skillet from the oven. To prevent the complete solidification of the pectin hydrocolloid network, the system must rest for exactly 15 to 20 minutes to allow the boiling caramel to drop to an optimal viscosity (approximately 50°C)<sup>71</sup>. Center a large, flat serving platter directly over the skillet. Utilizing heat-resistant gloves, apply immense pressure to sandwich the platter and skillet together, and rapidly, smoothly invert the entire mass<sup>69</sup>. The gravity-assisted demolding will cleanly release the stabilized apple and caramel matrix onto the plate, revealing the highly structured, caramelized geometry of the regional Tarte Tatin<sup>71</sup>.

## Conclusion

The construction of a Berlin-localized Tarte Tatin transcends rudimentary baking; it is a profound exercise in applied biochemistry, thermodynamics, and rheological engineering. By shunning highly standardized commercial inputs in favor of regional Brandenburg terroir, the baker must actively manage a cascade of complex, volatile reactions that are highly sensitive to temperature and time. The successful execution of this dish demands absolute mastery over the proteolytic depolymerization of the glutenin macropolymer via sourdough lactic acid bacteria, the precise geometric and polymorphic stabilization of cultured butter during lamination, and the violent, acid-catalyzed Arrhenius kinetics governing the caramelization of raw beet sugar and high-methoxyl pectin gelation. The resulting pastry is a highly sophisticated, structurally immaculate stabilization of chaos, firmly anchored in the unique physicochemical reality of its regional ingredients, offering a sensory profile that is entirely unreplicable with standardized, globalized commodities.

## Works cited

1. Old apple and pear varieties - Manufaktur Jörg Geiger, <https://www.manufaktur-joerg-geiger.de/en/about-Us/Old-apple-and-pear-varieties/>
2. Belle de Boskoop apple variety: taste & harvest - Plantura Magazin, <https://plantura.garden/uk/fruits/apple-varieties/belle-de-boskoop>
3. Berlepsch - Wikipedia, <https://de.wikipedia.org/wiki/Berlepsch>
4. Variety of apples | RAMSEIER, <https://www.ramseier.ch/en/variety-of-apples/>
5. Freiherr von Berlepsch Apple - General Fruit Growing, <https://growingfruit.org/t/freiherr-von-berlepsch-apple/41607>
6. Structure formation in sugar containing pectin gels | Request PDF, [https://www.researchgate.net/publication/257529539\\_Structure\\_formation\\_in\\_sugar\\_containing\\_pectin\\_gels\\_-\\_Influence\\_of\\_tartaric\\_acid\\_content\\_pH\\_and\\_cooling\\_rate\\_on\\_the\\_gelation\\_of\\_high-methoxylated\\_pectin](https://www.researchgate.net/publication/257529539_Structure_formation_in_sugar_containing_pectin_gels_-_Influence_of_tartaric_acid_content_pH_and_cooling_rate_on_the_gelation_of_high-methoxylated_pectin)
7. Postharvest quality of integrated and organically produced apple fruit, [https://www.researchgate.net/publication/222297773\\_Postharvest\\_quality\\_of\\_integrated\\_and\\_organically\\_produced\\_apple\\_fruit](https://www.researchgate.net/publication/222297773_Postharvest_quality_of_integrated_and_organically_produced_apple_fruit)
8. Monselise, Shaul P - CRC Handbook of Fruit Set and Development, <https://www.scribd.com/document/492014612/Monselise-Shaul-P-CRC-Handbook-of-Fruit-Set-and-Development-1986>
9. Compositional Characteristics of Fruits of several Apple (Malus, [https://www.researchgate.net/publication/49591548\\_Compositional\\_Characteristics\\_of\\_Fruits\\_of\\_several\\_Apple\\_Malus\\_domestica\\_Borkh\\_Cultivars](https://www.researchgate.net/publication/49591548_Compositional_Characteristics_of_Fruits_of_several_Apple_Malus_domestica_Borkh_Cultivars)
10. Cane vs. Beet Sugar: A Difference? - WhatSugar, <https://www.whatsugar.com/post/difference-between-cane-and-beet-sugar>
11. Sugar Cane vs. Sugar Beets: Examining Their Differences, <https://www.rockymountainsoda.com/blogs/soda-news/sugar-cane-vs-sugar-beets-examining-their-differences>
12. Sensory differences between product matrices made with beet and, <https://pubmed.ncbi.nlm.nih.gov/25308166/>
13. Beet Sugar vs. Cane Sugar: Which Is Healthier? - Healthline, <https://www.healthline.com/nutrition/beet-sugar-vs-cane-sugar>
14. Die Gläserne Molkerei, <https://glaeserne-molkerei.de/>
15. Fass Butter – Gläserne Molkerei – 250g - Open Food Facts, <https://de.openfoodfacts.org/produkt/4250017300914/fass-butter-gl%C3%A4serne-molkerei>
16. 83 Biochemistry Of Butter And Ghee.pdf
17. Why does butter from Europe taste so much better than American, <https://www.quora.com/Why-does-butter-from-Europe-taste-so-much-better-than-American-butter-Is-it-the-cream-content>
18. Fass Butter – Gläserne Molkerei – 250g - Open Food Facts, <https://world.openfoodfacts.org/product/4250017300914/fass-butter-glaserne-molkerei>
19. Mehlnummern - Adler Muehle,

- <https://www.adler-muehle.de/info/frag/Nummern.html>
20. Mehltest | der brotdoc, <https://brotdoc.com/tag/mehltest/>
  21. Mehltypen und ihre Verwendung - Adler Muehle, <https://www.adler-muehle.de/info/frag/mehltypen.html>
  22. Weizenmehl Type 550 - Stadtmühle Waldenbuch, <https://xn--stadtmhle-waldenbuch-uec.de/p/1459/weizenmehl-type-550/>
  23. DIFFERENT LEAVENING AGENTS & THEIR ROLES IN BAKING, <https://bakewithshivesh.com/different-leavening-agents-their-roles-in-baking/>
  24. Weizenmehl Type 812 - 2,5kg | Futtermittel Online Shop Mühle Gladen, <https://www.muehle-gladen.de/Naturkost/Backmehle-und-Schrot/Weizenmehl-Type-812-2-5kg.html>
  25. Laminated Dough Technique: Croissants, Puff Pastry, and Danish, <https://bakingtechniquesauthority.com/laminated-dough-technique/>
  26. Types of Yeast - Baking With Theory, <https://www.bakingwiththeory.com/theory/types-of-yeast/>
  27. Wheat Sourdough Fermentation: Effects of Time and Acidification on, [https://www.researchgate.net/publication/237328370\\_Wheat\\_Sourdough\\_Fermentation\\_Effects\\_of\\_Time\\_and\\_Acidification\\_on\\_Fundamental\\_Rheological\\_Properties](https://www.researchgate.net/publication/237328370_Wheat_Sourdough_Fermentation_Effects_of_Time_and_Acidification_on_Fundamental_Rheological_Properties)
  28. Gluten Hydrolysis and Depolymerization during Sourdough, <http://microsour.com/pdf/gluten-hydrolysis.pdf>
  29. Sourdough Fermentation and Gluten Reduction - MDPI, <https://www.mdpi.com/2036-7481/16/7/161>
  30. Bulk fermentation in depth: yeast activity, enzymatic reactions, gluten, <https://domson.co.uk/gb/academy/bread-technology/bulk-fermentation-in-depth>
  31. Proteolysis by Sourdough Lactic Acid Bacteria: Effects on Wheat, <https://pmc.ncbi.nlm.nih.gov/articles/PMC126681/>
  32. Effect of dominant Lactobacillus and yeast in sourdough on glutenin, <https://www.sciopen.com/article/10.11882/j.issn.0254-5071.2022.04.024>
  33. Sourdough Glossary: 100+ Terms & Process Definitions (Science, <https://thesourdoughcompanion.com/blog/sourdough-glossary/>
  34. Sugar, yeast, and laminated pastries: mastering the art of perfect, <https://bakeryinsider.com/sugar-yeast-laminated-pastries/>
  35. Sourdough Glossary | Sourdough Terminology You Need to Know, <https://kansascitysourdough.com/sourdough-glossary/>
  36. Sourdough Glossary, <https://celticsourdough.com/sourdough-glossary/>
  37. Influence of processing on the technological and sensory quality of, <https://www.itjfs.com/index.php/ijfs/article/view/2941/1587>
  38. Depolymerisation and re-polymerisation of wheat glutenin during, <https://repository.tno.nl/SingleDoc?docId=12571>
  39. (PDF) Structural Changes of Gliadins during Sourdough Fermentation, [https://www.researchgate.net/publication/355720024\\_Structural\\_Changes\\_of\\_Gliadins\\_during\\_Sourdough\\_Fermentation](https://www.researchgate.net/publication/355720024_Structural_Changes_of_Gliadins_during_Sourdough_Fermentation)
  40. Gliadin Degradation Ability of Artisanal Lactic Acid Bacteria, The, <https://lifescienceglobal.com/pms/index.php/jnt/article/view/1419>

41. Microstructural and conformational changes of gluten proteins in, [https://www.researchgate.net/publication/331712711\\_Microstructural\\_and\\_conformational\\_changes\\_of\\_gluten\\_proteins\\_in\\_wheat-rye\\_sourdough](https://www.researchgate.net/publication/331712711_Microstructural_and_conformational_changes_of_gluten_proteins_in_wheat-rye_sourdough)
42. Change in Glutenin Macropolymer Secondary Structure in Wheat, <https://pubmed.ncbi.nlm.nih.gov/28013448/>
43. Prolamin degradation in sourdoughs - Helda - University of Helsinki, <https://helda.helsinki.fi/bitstreams/fce78ed4-e7bc-4002-ad73-7943784ed766/download>
44. The impact of redox agents on further dough development, [https://www.researchgate.net/publication/315784576\\_The\\_impact\\_of\\_redox\\_agents\\_on\\_further\\_dough\\_development\\_relaxation\\_and\\_elastic\\_recoil\\_during\\_lamination\\_and\\_fermentation\\_of\\_multi-layered\\_pastry\\_dough](https://www.researchgate.net/publication/315784576_The_impact_of_redox_agents_on_further_dough_development_relaxation_and_elastic_recoil_during_lamination_and_fermentation_of_multi-layered_pastry_dough)
45. What does cold proof do for your dough? : r/Sourdough - Reddit, [https://www.reddit.com/r/Sourdough/comments/1cyr7uo/what\\_does\\_cold\\_proof\\_do\\_for\\_your\\_dough/](https://www.reddit.com/r/Sourdough/comments/1cyr7uo/what_does_cold_proof_do_for_your_dough/)
46. A Spin on the Standard Apple Cheese Tart - Bubbling Starter, <https://bubblingstarter.com/a-spin-on-the-standard-apple-cheese-tart/>
47. Jalapeno, Honey, & Brie Sourdough Turnovers - Bubbling Starter, <https://bubblingstarter.com/jalapeno-honey-brie-sourdough-turnovers/>
48. Lighten up: let's talk leavening, <https://hungryhungryhippocampus.com/2020/08/21/lighten-up-lets-talk-leavening/>
49. Effect of mixing on puff pastry dough microstructure and rheology, <https://www.longdom.org/proceedings/effect-of-mixing-on-puff-pastry-dough-microstructure-and-rheology-61242.html>
50. The complete guide to leavening agents and chemical leaveners, <https://bakeschool.com/leavening-agents-in-baking/>
51. Sustainable Cider Apple Production Eleni Vysini, Jim Dunwell , Bob, <http://archiveofciderpomology.co.uk/ArchiveReports/SustainableCiderAppleProduction7.2.4.pdf>
52. Quality Changes and Nonenzymic Browning Intermediates in, [https://www.researchgate.net/publication/229511220\\_Quality\\_Changes\\_and\\_Nonenzymic\\_Browning\\_Intermediates\\_in\\_Grapefruit\\_Juice\\_During\\_Storage](https://www.researchgate.net/publication/229511220_Quality_Changes_and_Nonenzymic_Browning_Intermediates_in_Grapefruit_Juice_During_Storage)
53. (PDF) Sugar - ResearchGate, [https://www.researchgate.net/publication/229503721\\_Sugar](https://www.researchgate.net/publication/229503721_Sugar)
54. Chemistry and Processing of Sugarbeet and Sugarcane, Volume 9, <https://www.scribd.com/document/466386531/Chemistry-and-Processing-of-Sugarbeet-and-Sugarcane-Volume-9-pdf>
55. Flashcard - Food Chemistry | LMS - MemoCard, <https://www.memocard.ch/EN/Lernkarten/LernkartenInfo/Info/food-chemistry/f3d17b39-42ce-4cda-90c3-cf067925c7ca>
56. Hard Candy Production and Quality Parameters: A review - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11214042/>
57. Thermal behaviour and degradation kinetics of apple pomace flours, [https://www.researchgate.net/publication/330319430\\_Thermal\\_behaviour\\_and\\_de](https://www.researchgate.net/publication/330319430_Thermal_behaviour_and_de)

[gradation\\_kinetics\\_of\\_apple\\_pomace\\_flours](#)

58. Effects of sugars in batter formula and baking conditions on 5, [https://www.researchgate.net/publication/257422764\\_Effects\\_of\\_sugars\\_in\\_batter\\_formula\\_and\\_baking\\_conditions\\_on\\_5-hydroxymethylfurfural\\_and\\_furfural\\_formation\\_in\\_sponge\\_cake\\_models](https://www.researchgate.net/publication/257422764_Effects_of_sugars_in_batter_formula_and_baking_conditions_on_5-hydroxymethylfurfural_and_furfural_formation_in_sponge_cake_models)
59. Changes in physical and chemical properties during the production, [https://www.researchgate.net/publication/285697394\\_Changes\\_in\\_physical\\_and\\_chemical\\_properties\\_during\\_the\\_production\\_of\\_palm\\_sugar\\_syrup\\_by\\_open\\_pan\\_and\\_vacuum\\_evaporator](https://www.researchgate.net/publication/285697394_Changes_in_physical_and_chemical_properties_during_the_production_of_palm_sugar_syrup_by_open_pan_and_vacuum_evaporator)
60. Chlorogenic Acid Oxidation-Induced Greening of Sunflower Butter, [https://digitalcommons.chapman.edu/cgi/viewcontent.cgi?article=1024&context=food\\_science\\_articles](https://digitalcommons.chapman.edu/cgi/viewcontent.cgi?article=1024&context=food_science_articles)
61. Cross-linking of Peanut Allergen Ara h 2 by Polyphenol Oxidase, [https://www.researchgate.net/publication/284714079\\_Cross-linking\\_of\\_Peanut\\_Allergen\\_Ara\\_h\\_2\\_by\\_Polyphenol\\_Oxidase\\_Digestibility\\_and\\_Potential\\_Allergenicity\\_Assessment](https://www.researchgate.net/publication/284714079_Cross-linking_of_Peanut_Allergen_Ara_h_2_by_Polyphenol_Oxidase_Digestibility_and_Potential_Allergenicity_Assessment)
62. Recent Development on the Chemical Composition and Phenolic, [https://www.researchgate.net/publication/374683094\\_Recent\\_Development\\_on\\_the\\_Chemical\\_Composition\\_and\\_Phenolic\\_Extraction\\_Methods\\_of\\_Apple\\_Malus\\_domestica-A\\_Review](https://www.researchgate.net/publication/374683094_Recent_Development_on_the_Chemical_Composition_and_Phenolic_Extraction_Methods_of_Apple_Malus_domestica-A_Review)
63. Polyphenolic Compounds Analysis of Old and New Apple Cultivars, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5789330/>
64. Comparison of Soluble and Plastidal Polyphenol Oxidase in Mature, [https://www.researchgate.net/publication/250193327\\_Comparison\\_of\\_Soluble\\_and\\_d\\_Plastidal\\_Polyphenol\\_Oxidase\\_in\\_Mature\\_Apples](https://www.researchgate.net/publication/250193327_Comparison_of_Soluble_and_d_Plastidal_Polyphenol_Oxidase_in_Mature_Apples)
65. Effects of pulsed electric field treatment of apple mash on juice yield, [https://www.researchgate.net/publication/222708110\\_Effects\\_of\\_pulsed\\_electric\\_field\\_treatment\\_of\\_apple\\_mash\\_on\\_juice\\_yield\\_and\\_quality\\_attributes\\_of\\_apple\\_juice](https://www.researchgate.net/publication/222708110_Effects_of_pulsed_electric_field_treatment_of_apple_mash_on_juice_yield_and_quality_attributes_of_apple_juice)
66. (PDF) Role of cosolutes in gelation of high-methoxy pectin. Part 1, [https://www.researchgate.net/publication/248425019\\_Role\\_of\\_cosolutes\\_in\\_gelation\\_of\\_high-methoxy\\_pectin\\_Part\\_1\\_Comparison\\_of\\_sugars\\_and\\_polyols](https://www.researchgate.net/publication/248425019_Role_of_cosolutes_in_gelation_of_high-methoxy_pectin_Part_1_Comparison_of_sugars_and_polyols)
67. Synergistic gelling mechanism of RG-I rich citrus pectic, [https://eprints.whiterose.ac.uk/id/eprint/171496/3/Zheng\\_FOODCHEM\\_2021.pdf](https://eprints.whiterose.ac.uk/id/eprint/171496/3/Zheng_FOODCHEM_2021.pdf)
68. Development of low-calorie gels from sunflower pectin extracted by, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11953053/>
69. French Tarte Tatin - Persnickety Panhandler, <https://www.persnicketypanhandler.com/blog/french-tarte-tatin>
70. Apple Tarte Tatin - Riegl Palate, <https://www.rieglpalate.com/apple-tarte-tatin/>
71. Tarte Tatin Recipe | Food Network, <https://www.foodnetwork.com/recipes/tarte-tatin-recipe3-1940720>
72. Classic Tarte Tatin - Pardon Your French, <https://www.pardonyourfrench.com/classic-tarte-tatin/>
73. perfect apple tarte tatin – smitten kitchen, <https://smittenkitchen.com/2019/11/perfect-apple-tarte-tatin/>

74. A Baker's Guide to Leavening Agents | Suprima Bakeries,  
<https://www.suprima.com.au/blog/leavening-agents/>
75. Classic Apple Tarte Tatin Recipe | King Arthur Baking,  
<https://www.kingarthurbaking.com/recipes/classic-apple-tarte-tatin-recipe>
76. How to Make Tarte Tatin - Joe Pastry,  
[https://joepastry.com/2008/how\\_to\\_make\\_tarte\\_tatin/](https://joepastry.com/2008/how_to_make_tarte_tatin/)
77. Peach Tarte Tatin - Cooking with Cas,  
<https://www.cookingwithcas.com/peach-tarte-tatin.html>
78. Experiments with Laminating Lean Dough - Breadtopia,  
<https://breadtopia.com/experiments-with-laminating-lean-dough/>